



## MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

# Medical Policy Lysis of Epidural Adhesions

### Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

### Policy Number: 598

BCBSA Reference Number: 8.01.18

NCD/LCD: N/A

### Related Policies

None

### Policy

#### Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO Blue<sup>SM</sup> and Medicare PPO Blue<sup>SM</sup> Members

Catheter-based techniques for lysis of epidural adhesions, with or without endoscopic guidance, are considered **INVESTIGATIONAL**. Techniques used either alone or in combination include mechanical disruption with a catheter and/or injection of hypertonic solutions with corticosteroids, analgesics, or hyaluronidase.

### Prior Authorization Information

#### Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

#### Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	This is <b>not</b> a covered service.
Commercial PPO and Indemnity	This is <b>not</b> a covered service.
Medicare HMO Blue <sup>SM</sup>	This is <b>not</b> a covered service.
Medicare PPO Blue <sup>SM</sup>	This is <b>not</b> a covered service.

## CPT Codes / HCPCS Codes / ICD Codes

*Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.*

*Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.*

### CPT Codes

CPT codes:	Code Description
62263	Percutaneous lysis of epidural adhesions using solution injection (eg, hypertonic saline, enzyme) or mechanical means (eg, catheter) including radiologic localization (includes contrast when administered), multiple adhesiolysis sessions; 2 or more days
62264	Percutaneous lysis of epidural adhesions using solution injection (eg, hypertonic saline, enzyme) or mechanical means (eg, catheter) including radiologic localization (includes contrast when administered), multiple adhesiolysis sessions; 1 day

### Description

#### Epidural Fibrosis and Adhesive Arachnoiditis

Epidural fibrosis with or without adhesive arachnoiditis most commonly occurs as a complication of spinal surgery and may be included under the diagnosis of "failed back surgery syndrome". Both conditions result from the manipulation of the supporting structures of the spine. Epidural fibrosis can occur in isolation, but adhesive arachnoiditis is rarely present without associated epidural fibrosis. Arachnoiditis is most frequently seen in patients who have undergone multiple surgical procedures.

Epidural fibrosis and adhesive arachnoiditis are related to inflammatory reactions that result in the entrapment of nerves within dense scar tissue, increasing the susceptibility of the nerve root to compression or tension. The condition most frequently involves the nerves within the lumbar spine and cauda equina. Signs and symptoms indicate the involvement of multiple nerve roots and include low back pain, radicular pain, tenderness, sphincter disturbances, limited trunk mobility, muscular spasm or contracture, and motor-sensory and reflex changes. Typically, pain is characterized as constant and burning. In some cases, pain and disability are severe, leading to analgesic dependence and chronic invalidism.

#### Treatment

Lysis of epidural adhesions, also called the Racz procedure, has been investigated as a treatment option. The Racz procedure involves the passage of a fluoroscopically guided catheter (the Racz catheter), inserted either endoscopically or percutaneously, and the use of epidural injections of hypertonic saline in conjunction with corticosteroids and analgesics. Theoretically, the use of hypertonic saline results in mechanical disruption of the adhesions. The saline may also function to reduce edema within previously scarred and/or inflamed nerves. Finally, manipulating the catheter at the time of the injection may disrupt adhesions. Spinal endoscopy has been used to guide the lysis procedure, but the procedure is more commonly performed percutaneously using epidurography to guide catheter placement and identify nonfilling adhesions that indicate epidural scarring. Using endoscopy guidance, a flexible fiberoptic catheter is inserted into the sacral hiatus, providing 3-dimensional visualization to steer the catheter toward the adhesions. With the increased visualization, the catheter is more apt to precisely place the injectate in the epidural space and onto the nerve root. Various protocols for lysis have been described; in some situations, the catheter may remain in place for several days for serial treatment sessions.

Endoscopic epidurolysis is also being investigated to treat degenerative chronic low back pain, including spondylolisthesis, stenosis, and hernia associated with radiculopathy. Along with mechanical adhesiolysis, hyaluronidase, ciprofloxacin, and ozone have been applied.

## Summary

Lysis of epidural adhesions involves passing a catheter, either endoscopically or percutaneously, under fluoroscopic guidance into the epidural space to break up adhesions and reduce pain and inflammation.

For individuals who have epidural adhesions who receive lysis, the evidence includes randomized controlled trials. The relevant outcomes are symptoms, functional outcomes, quality of life, medication use, and treatment-related morbidity. Several randomized controlled trials have reported benefits for epidural lysis of adhesions compared with placebo treatment. Many of these trials were conducted at the same center. The interpretation of these trials is limited by differences in patients, populations, and treatment protocols. The treatment for lysis of adhesions varied in the use of mechanical disruption, the type of lytic medications used, and the number of injections given. There was also a large effect in the placebo group, raising questions whether some component of the placebo treatment may be therapeutic. Larger trials with standardized treatment protocols would help determine whether specific treatment protocols have beneficial effects in specific patient populations. The evidence is insufficient to determine the effects of the technology on health outcomes.

## Policy History

Date	Action
1/2020	BCBSA National medical policy review. Description, summary and references updated. Policy statements unchanged.
1/2019	BCBSA National medical policy review. Description, summary and references updated. Policy statements unchanged.
3/2014	New references added from BCBSA National medical policy.
11/2011-4/2012	Medical policy ICD 10 remediation: Formatting, editing and coding updates. No changes to policy statements.
1/2012	Reviewed - Medical Policy Group - Neurology and Neurosurgery. No changes to policy statements.
12/2011	New policy, effective 12/2011, describing ongoing non-coverage.

## Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

## References

1. Chopra P, Smith HS, Deer TR, et al. Role of adhesiolysis in the management of chronic spinal pain: a systematic review of effectiveness and complications. *Pain Physician*. Jan 2005;8(1):87-100. PMID 16850047.
2. Trescot AM, Chopra P, Abdi S, et al. Systematic review of effectiveness and complications of adhesiolysis in the management of chronic spinal pain: an update. *Pain Physician*. Jan 2007;10(1):129-146. PMID 17256027.
3. Helm li S, Benyamin RM, Chopra P, et al. Percutaneous adhesiolysis in the management of chronic low back pain in post lumbar surgery syndrome and spinal stenosis: a systematic review. *Pain Physician*. Jul-Aug 2012;15(4):E435-462. PMID 22828693.
4. Racz GB, Heavner JE, Trescot A. Percutaneous lysis of epidural adhesions--evidence for safety and efficacy. *Pain Pract*. Jul-Aug 2008;8(4):277-286. PMID 18503627.
5. Hayek SM, Helm S, Benyamin RM, et al. Effectiveness of spinal endoscopic adhesiolysis in post lumbar surgery syndrome: a systematic review. *Pain Physician*. Mar-Apr 2009;12(2):419-435. PMID 19305488.

6. Epter RS, Helm S, 2nd, Hayek SM, et al. Systematic review of percutaneous adhesiolysis and management of chronic low back pain in post lumbar surgery syndrome. *Pain Physician*. Mar-Apr 2009;12(2):361-378. PMID 19305485.
7. Helm S, Hayek SM, Colson J, et al. Spinal endoscopic adhesiolysis in post lumbar surgery syndrome: an update of assessment of the evidence. *Pain Physician*. Apr 2013;16(2 Suppl):SE125-150. PMID 23615889.
8. Gerdesmeyer L, Wagenpfeil S, Birkenmaier C, et al. Percutaneous epidural lysis of adhesions in chronic lumbar radicular pain: a randomized, double-blind, placebo-controlled trial. *Pain Physician*. May-Jun 2013;16(3):185- 196. PMID 23703406.
9. Manchikanti L, Cash KA, McManus CD, et al. The preliminary results of a comparative effectiveness evaluation of adhesiolysis and caudal epidural injections in managing chronic low back pain secondary to spinal stenosis: a randomized, equivalence controlled trial. *Pain Physician*. Nov-Dec 2009;12(6):E341-354. PMID 19935991.
10. Manchikanti L, Singh V, Cash KA, et al. A comparative effectiveness evaluation of percutaneous adhesiolysis and epidural steroid injections in managing lumbar post surgery syndrome: a randomized, equivalence controlled trial. *Pain Physician*. Nov-Dec 2009;12(6):E355-368. PMID 19935992.
11. Manchikanti L, Singh V, Cash KA, et al. Assessment of effectiveness of percutaneous adhesiolysis and caudal epidural injections in managing post lumbar surgery syndrome: 2-year follow-up of a randomized, controlled trial. *J Pain Res*. Jan 2012;5:597-608. PMID 23293536.
12. Manchikanti L, Rivera JJ, Pampati V, et al. One day lumbar epidural adhesiolysis and hypertonic saline neurolysis in treatment of chronic low back pain: a randomized, double-blind trial. *Pain Physician*. Apr 2004;7(2):177-186. PMID 16868590.
13. Manchikanti L, Pampati V, Fellows B, et al. Role of one day epidural adhesiolysis in management of chronic low back pain: a randomized clinical trial. *Pain Physician*. Apr 2001;4(2):153-166. PMID 16902688.
14. Wagner KJ, Sprenger T, Pecho C, et al. [Risks and complications of epidural neurolysis -- a review with case report] [German]. *Anesthesiol Intensivmed Notfallmed Schmerzther*. Apr 2006;41(4):213-222. PMID 16636945.
15. Manchikanti L, Malla Y, Wargo BW, et al. A prospective evaluation of complications of 10,000 fluoroscopically directed epidural injections. *Pain Physician*. Mar-Apr 2012;15(2):131-140. PMID 22430650.
16. Manchikanti L, Rivera JJ, Pampati V, et al. Spinal endoscopic adhesiolysis in the management of chronic low back pain: a preliminary report of a randomized, double-blind trial. *Pain Physician*. Jul 2003;6(3):259-267. PMID 16880869.
17. Donato AD, Fontana C, Pinto R, et al. The effectiveness of endoscopic epidurolysis in treatment of degenerative chronic low back pain: a prospective analysis and follow-up at 48 months. *Acta Neurochir Suppl*. Nov 2011;108:67-73. PMID 21107940.
18. Manchikanti L, Pampati V, Bakhit CE, et al. Non-endoscopic and endoscopic adhesiolysis in post-lumbar laminectomy syndrome: a one-year outcome study and cost effectiveness analysis. *Pain Physician*. Oct 1999;2(3):52-58. PMID 16906216.
19. Manchikanti L, Pakanati RR, Pampati V. The value and safety of epidural endoscopic adhesiolysis. *Am J Anesthesiol*. 2000;27(5):275-279. PMID.
20. Manchikanti L, Abdi S, Atluri S, et al. An update of comprehensive evidence-based guidelines for interventional techniques in chronic spinal pain. Part II: guidance and recommendations. *Pain Physician*. Apr 2013;16(2 Suppl):S49-283. PMID 23615883.
21. Chou R, Loeser JD, Owens DK, et al. Interventional therapies, surgery, and interdisciplinary rehabilitation for low back pain: an evidence-based clinical practice guideline from the American Pain Society. *Spine (Phila Pa 1976)*. May 1 2009;34(10):1066-1077. PMID 19363457.