

Blue Cross Blue Shield of Massechusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

Medical Policy Optical Coherence Tomography for Imaging of Coronary Arteries

Table of Contents

- Policy: Commercial
- Coding Information
- Information Pertaining to All Policies

References

- Policy: Medicare
- <u>Authorization Information</u>
- Description
- Policy History
- Policy Number: 915

BCBSÁ Reference Number: 2.02.29A NCD/LCD: N/A

Related Policies

Anterior Eye Segment Optical Imaging, #084

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO BlueSM and Medicare PPO BlueSM Members

Optical coherence tomography is considered **INVESTIGATIONAL** when used as an adjunct to percutaneous coronary interventions with stenting.

Optical coherence tomography is considered **INVESTIGATIONAL** in all other situations, including but not limited to, risk stratification of intracoronary atherosclerotic plaques and follow-up evaluation of stenting.

Prior Authorization Information

Inpatient

 For services described in this policy, precertification/preauthorization <u>IS REQUIRED</u> for all products if the procedure is performed <u>inpatient</u>.

Outpatient

 For services described in this policy, see below for products where prior authorization <u>might be</u> <u>required</u> if the procedure is performed <u>outpatient</u>.

	Outpatient
Commercial Managed Care (HMO and POS)	This is not a covered service.
Commercial PPO and Indemnity	This is not a covered service.
Medicare HMO Blue sM	This is not a covered service.
Medicare PPO Blue SM	This is not a covered service.

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

CPT Codes

There is no specific CPT code for this service.

Description

Optical coherence tomography (OCT) is an imaging technique that uses near-infrared light to image the coronary arteries. Potential applications in cardiology include evaluating the characteristics of coronary artery plaques for the purpose of risk stratification and following coronary stenting to determine the success of the procedure.

Summary

Optical coherence tomography (OCT) is an imaging technique that has some advantages over intravascular ultrasound (IVUS) for imaging coronary arteries. It has a higher resolution and provides images with greater detail for accessible structures compared to IVUS. Case series have demonstrated that OCT can be performed with a high success rate and few complications. Head-to-head comparisons of OCT and IVUS report that OCT picks up additional abnormalities that are not detected by IVUS, implying that OCT is a more sensitive test compared to IVUS.

As an adjunct to PCI, OCT may improve upon the ability to pick up abnormalities compared to IVUS, and this may lead to changes in management. However, the current evidence is limited and includes relatively small numbers of patients who have received OCT. As a result, it is not possible to determine the degree of improvement with OCT, or the clinical significance of this improvement. Therefore, the use of OCT as an adjunct to PCI is considered investigational.

For the indications of risk stratification of coronary plaques and follow-up of stenting, OCT may also be more accurate than IVUS for imaging of superficial structures. However, the clinical utility of IVUS has not been demonstrated for these indications, since test results do not lead to changes in management that improve outcomes. Therefore, clinical utility has not been demonstrated for OCT for the same reasons. As a result, OCT is considered investigational for risk stratification of coronary plaques and for follow-up post-stent implantation.

Policy History

Date	Action
2/2020	Policy updated with literature review through February 1, 2020, references added. Policy statements unchanged.
1/2017	Clarified coding information for the 2017 code changes.
3/2015	New references added from BCBSA National medical policy.
5/2014	New references from BCBSA National medical policy.
2/2013	New policy describing non-coverage. Effective 02/04/13

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information: <u>Medical Policy Terms of Use</u> <u>Managed Care Guidelines</u> <u>Indemnity/PPO Guidelines</u> <u>Clinical Exception Process</u> <u>Medical Technology Assessment Guidelines</u>

References

- Prati F, Regar E, Mintz GS, et al. Expert review document on methodology, terminology, and clinical applications of optical coherence tomography: physical principles, methodology of image acquisition, and clinical application for assessment of coronary arteries and atherosclerosis. Eur Heart J. Feb 2010;31(4):401-415. PMID 19892716
- Lindsay AC, Viceconte N, Di Mario C. Optical coherence tomography: has its time come? Heart. Sep 2011;97(17):1361-1362. PMID 21730261
- Low AF, Tearney GJ, Bouma BE, et al. Technology Insight: optical coherence tomography--current status and future development. Nat Clin Pract Cardiovasc Med. Mar 2006;3(3):154-162; quiz 172. PMID 16505861
- 4. Gonzalo N, Tearney GJ, Serruys PW, et al. Second-generation optical coherence tomography in clinical practice. High-speed data acquisition is highly reproducible in patients undergoing percutaneous coronary intervention. Rev Esp Cardiol. Aug 2010;63(8):893-903. PMID 20738934
- Fedele S, Biondi-Zoccai G, Kwiatkowski P, et al. Reproducibility of coronary optical coherence tomography for lumen and length measurements in humans (The CLI-VAR [Centro per la Lotta contro l'Infarto-VARiability] study). The American journal of cardiology. Oct 15 2012;110(8):1106-1112. PMID 22748353
- Jamil Z, Tearney G, Bruining N, et al. Interstudy reproducibility of the second generation, Fourier domain optical coherence tomography in patients with coronary artery disease and comparison with intravascular ultrasound: a study applying automated contour detection. Int J Cardiovasc Imaging. Jan 2013;29(1):39-51. PMID 22639296
- Liu Y, Shimamura K, Kubo T, et al. Comparison of longitudinal geometric measurement in human coronary arteries between frequency-domain optical coherence tomography and intravascular ultrasound. Int J Cardiovasc Imaging. Feb 2014;30(2):271-277. PMID 24272334
- Brugaletta S, Garcia-Garcia HM, Gomez-Lara J, et al. Reproducibility of qualitative assessment of stent struts coverage by optical coherence tomography. Int J Cardiovasc Imaging. Jan 2013;29(1):5-11. PMID 22415543
- Jang IK, Bouma BE, Kang DH, et al. Visualization of coronary atherosclerotic plaques in patients using optical coherence tomography: comparison with intravascular ultrasound. J Am Coll Cardiol. Feb 20 2002;39(4):604- 609. PMID 11849858
- Kubo T, Nakamura N, Matsuo Y, et al. Virtual histology intravascular ultrasound compared with optical coherence tomography for identification of thin-cap fibroatheroma. Int Heart J. 2011;52(3):175-179. PMID 21646741
- 11. Miyamoto Y, Okura H, Kume T, et al. Plaque characteristics of thin-cap fibroatheroma evaluated by OCT and IVUS. JACC Cardiovasc Imaging. Jun 2011;4(6):638-646. PMID 21679899
- 12. Uemura S, Ishigami KI, Soeda T, et al. Thin-cap fibroatheroma and microchannel findings in optical coherence tomography correlate with subsequent progression of coronary atheromatous plaques. Eur Heart J. Aug 10 2011. PMID 21831910
- Yonetsu T, Kakuta T, Lee T, et al. In vivo critical fibrous cap thickness for rupture-prone coronary plaques assessed by optical coherence tomography. Eur Heart J. May 2011;32(10):1251-1259. PMID 21273202
- 14. Guo J, Chen YD, Tian F, et al. Thrombosis and morphology of plaque rupture using optical coherence tomography. Chin Med J (Engl). Mar 2013;126(6):1092-1095. PMID 23506584
- Jia H, Abtahian F, Aguirre AD, et al. In vivo diagnosis of plaque erosion and calcified nodule in patients with acute coronary syndrome by intravascular optical coherence tomography. J Am Coll Cardiol. Nov 5 2013;62(19):1748-1758. PMID 23810884
- 16. Gamou T, Sakata K, Matsubara T, et al. Impact of thin-cap fibroatheroma on predicting deteriorated coronary flow during interventional procedures in acute as well as stable coronary syndromes: insights from optical coherence tomography analysis. Heart Vessels. Jul 19 2014. PMID 25037112
- 17. Galon MZ, Wang Z, Bezerra HG, et al. Differences determined by optical coherence tomography volumetric analysis in non-culprit lesion morphology and inflammation in ST-segment elevation myocardial infarction and stable angina pectoris patients. Catheter Cardiovasc Interv. Sep 1 2014. PMID 25178981
- 18. Wykrzykowska JJ, Diletti R, Gutierrez-Chico JL, et al. Plaque sealing and passivation with a mechanical selfexpanding low outward force nitinol vShield device for the treatment of IVUS and OCT-derived thin cap fibroatheromas (TCFAs) in native coronary arteries: report of the pilot study

vShield Evaluated at Cardiac hospital in Rotterdam for Investigation and Treatment of TCFA (SECRITT). EuroIntervention. Dec 20 2012;8(8):945-954. PMID 22669133

- 19. Fitzgerald PJ, Oshima A, Hayase M, et al. Final results of the Can Routine Ultrasound Influence Stent Expansion (CRUISE) study. Circulation. Aug 1 2000;102(5):523-530. PMID 10920064
- Jakabcin J, Spacek R, Bystron M, et al. Long-term health outcome and mortality evaluation after invasive coronary treatment using drug eluting stents with or without the IVUS guidance. Randomized control trial. HOME DES IVUS. Catheter Cardiovasc Interv. Mar 1 2010;75(4):578-583. PMID 19902491
- Roy P, Steinberg DH, Sushinsky SJ, et al. The potential clinical utility of intravascular ultrasound guidance in patients undergoing percutaneous coronary intervention with drug-eluting stents. Eur Heart J. Aug 2008;29(15):1851-1857. PMID 18550555
- 22. Smith SC, Jr., Dove JT, Jacobs AK, et al. ACC/AHA guidelines for percutaneous coronary intervention (revision of the 1993 PTCA guidelines)-executive summary: a report of the American College of Cardiology/American Heart Association task force on practice guidelines (Committee to revise the 1993 guidelines for percutaneous transluminal coronary angioplasty) endorsed by the Society for Cardiac Angiography and Interventions. Circulation. Jun 19 2001;103(24):3019-3041. PMID 11413094
- 23. Habara M, Nasu K, Terashima M, et al. Impact of frequency-domain optical coherence tomography guidance for optimal coronary stent implantation in comparison with intravascular ultrasound guidance. Circ Cardiovasc Interv. Apr 2012;5(2):193-201. PMID 22456026
- 24. Prati F, Di Vito L, Biondi-Zoccai G, et al. Angiography alone versus angiography plus optical coherence tomography to guide decision-making during percutaneous coronary intervention: the Centro per la Lotta contro l'Infarto-Optimisation of Percutaneous Coronary Intervention (CLI-OPCI) study. EuroIntervention. Nov 22 2012;8(7):823-829. PMID 23034247
- Yamaguchi T, Terashima M, Akasaka T, et al. Safety and feasibility of an intravascular optical coherence tomography image wire system in the clinical setting. Am J Cardiol. Mar 1 2008;101(5):562-567. PMID 18307999
- 26. Kawamori H, Shite J, Shinke T, et al. The ability of optical coherence tomography to monitor percutaneous coronary intervention: detailed comparison with intravascular ultrasound. J Invasive Cardiol. Nov 2010;22(11):541-545. PMID 21041851
- 27. Bezerra HG, Attizzani GF, Sirbu V, et al. Optical coherence tomography versus intravascular ultrasound to evaluate coronary artery disease and percutaneous coronary intervention. JACC Cardiovasc Interv. Mar 2013;6(3):228-236. PMID 23517833
- Sohn J, Hur SH, Kim IC, et al. A comparison of tissue prolapse with optical coherence tomography and intravascular ultrasound after drug-eluting stent implantation. Int J Cardiovasc Imaging. Oct 2 2014. PMID 25273918
- 29. Sugiyama T, Kimura S, Akiyama D, et al. Quantitative assessment of tissue prolapse on optical coherence tomography and its relation to underlying plaque morphologies and clinical outcome in patients with elective stent implantation. Int J Cardiol. Sep 2014;176(1):182-190. PMID 25042663
- 30. Ann SH, Lim KH, De Jin C, et al. Multi-modality imaging for stent edge assessment. Heart Vessels. Jan 31 2014. PMID 24481539
- 31. Souteyrand G, Amabile N, Combaret N, et al. Invasive management without stents in selected acute coronary syndrome patients with a large thrombus burden: a prospective study of optical coherence tomography guided treatment decisions. EuroIntervention. Jul 19 2014. PMID 25033106
- Cervinka P, Spacek R, Bystron M, et al. Optical coherence tomography-guided primary percutaneous coronary intervention in ST-segment elevation myocardial infarction patients: a pilot study. Can J Cardiol. Apr 2014;30(4):420-427. PMID 24680171
- Radu MD, Raber L, Heo J, et al. Natural history of optical coherence tomography-detected non-flowlimiting edge dissections following drug-eluting stent implantation. EuroIntervention. Jan 22 2014;9(9):1085-1094. PMID 24064426
- 34. Capodanno D, Prati F, Pawlowsky T, et al. Comparison of optical coherence tomography and intravascular ultrasound for the assessment of in-stent tissue coverage after stent implantation. EuroIntervention. Nov 2009;5(5):538-543. PMID 20142173
- 35. Inoue T, Shite J, Yoon J, et al. Optical coherence evaluation of everolimus-eluting stents 8 months after implantation. Heart. Sep 2011;97(17):1379-1384. PMID 21051456

- Harris KC, Manouzi A, Fung AY, et al. Feasibility of optical coherence tomography in children with Kawasaki disease and pediatric heart transplant recipients. Circ Cardiovasc Imaging. Jul 2014;7(4):671-678. PMID 24874056
- 37. Barlis P, Gonzalo N, Di Mario C, et al. A multicentre evaluation of the safety of intracoronary optical coherence tomography. EuroIntervention. May 2009;5(1):90-95. PMID 19577988
- Lehtinen T, Nammas W, Airaksinen JK, et al. Feasibility and safety of frequency-domain optical coherence tomography for coronary artery evaluation: a single-center study. Int J Cardiovasc Imaging. Jun 2013;29(5):997- 1005. PMID 23417516
- Meneveau N, Ecarnot F, Souteyrand G, et al. Does optical coherence tomography optimize results of stenting? Rationale and study design. Am Heart J. Aug 2014;168(2):175-181 e171-172. PMID 25066556
- Lotfi A, Jeremias A, Fearon WF, et al. Expert consensus statement on the use of fractional flow reserve, intravascular ultrasound, and optical coherence tomography: a consensus statement of the Society of Cardiovascular Angiography and Interventions. Catheter Cardiovasc Interv. Mar 1 2014;83(4):509-518. PMID 24227282
- 41. Tearney GJ, Regar E, Akasaka T, et al. Consensus standards for acquisition, measurement, and reporting of intravascular optical coherence tomography studies: a report from the International Working Group for Intravascular Optical Coherence Tomography Standardization and Validation. J Am Coll Cardiol. Mar 20 2012;59(12):1058-1072. PMID 22421299
- 42. Ali ZA, et al. Optical coherence tomography compared with intravascular ultrasound and with angiography to guide coronary stent implantation (ILUMIEN III: OPTIMIZE PCI): a randomised controlled trial. Lancet 2016 Clinical Trial. PMID 27806900
- 43. Jaguszewski M, et al. The REMEDEE-OCT Study: An Evaluation of the Bioengineered COMBO Dual-Therapy CD34 Antibody-Covered Sirolimus-Eluting Coronary Stent Compared with a Cobalt-Chromium Everolimus-Eluting Stent in Patients with Acute Coronary Syndromes: Insights From Optical Coherence Tomography Imaging Analysis. JACC Cardiovasc Interv 2017 - Clinical Trial. PMID 28279316