



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

## Medical Policy

# Transcutaneous Electrical Nerve Stimulation

### Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

### Policy Number: 003

BCBSA Reference Number: 1.01.09

NCD/LCD:

- National Coverage Determination (NCD) for Assessing Patient's Suitability for Electrical Nerve Stimulation Therapy (160.7.1)
- National Coverage Determination for Transcutaneous Electrical Nerve Stimulation (TENS) for Acute Post-Operative Pain (10.2)
- National Coverage Determination for Supplies Used in the Delivery of Transcutaneous Electrical Nerve Stimulation (TENS) and Neuromuscular Electrical Stimulation (NMES) (160.13)

### Related Policies

- Interferential Stimulation for Treatment of Pain, #[509](#)
- Temporomandibular Joint Dysfunction, #[035](#)
- Percutaneous Electrical Nerve Stimulation or Percutaneous Neuromodulation Therapy, #[172](#)

### Policy

#### Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity

A trial of transcutaneous electrical nerve stimulation (TENS) of at least 30 days may be considered **MEDICALLY NECESSARY** to establish efficacy for the management of refractory chronic pain (e.g., chronic musculoskeletal or neuropathic pain) that causes significant disruption of function when the following conditions have been met:

- The pain is unresponsive to at least 3 months of conservative medical therapy, AND
- The trial is monitored by a physician.

Refractory chronic pain is defined in this policy as pain that causes significant disruption of function and has not responded to at least 3 months of conservative therapy, including nonsteroidal anti-inflammatory medications, ice, rest, and/or physical therapy.

Documentation for the trial should include:

- Initial assessment/evaluation of the nature, duration, and perceived intensity of pain;
- The types and duration of prior treatments;
- Treatment plan including ongoing medications and proposed use of TENS unit, including the frequency and duration of treatment.

Clinical summary of the trial to determine efficacy should include:

- Perceived intensity of pain with and without TENS (e.g., 2 point or 30% improvement in visual analog scale [VAS]),
- Ongoing medication requirements for pain relief (if any),
- Other modalities (if any) in use for pain control, and
- Actual use of TENS on a daily basis (frequency and duration of application).

Continued use of TENS may be considered **MEDICALLY NECESSARY** for treatment of refractory chronic pain (e.g., chronic musculoskeletal or neuropathic pain) that causes significant disruption of function when the following conditions have been met:

- Efficacy has been demonstrated in an initial therapeutic trial; AND
- Compliance has been demonstrated in the therapeutic trial with the device used on a regular basis (eg, daily or near daily use) throughout the trial period.

*Note: A TENS billed as a purchased unit (modifier NU) must meet above criteria for continued use.*

TENS is **INVESTIGATIONAL** for the management of acute pain (e.g., postoperative or during labor and delivery).

The use of TENS for any other condition, including but not limited to the treatment of dementia and prevention of migraine headaches, is **INVESTIGATIONAL**.

## Medicare HMO Blue<sup>SM</sup> and Medicare PPO Blue<sup>SM</sup> Members

Medical necessity criteria and coding guidance can be found through the link below.

### [National Coverage Determinations \(NCDs\)](#)

- National Coverage Determination (NCD) for Assessing Patient's Suitability for Electrical Nerve Stimulation Therapy (160.7.1)
- National Coverage Determination for Transcutaneous Electrical Nerve Stimulation (TENS) for Acute Post-Operative Pain (10.2)
- National Coverage Determination for Supplies Used in the Delivery of Transcutaneous Electrical Nerve Stimulation (TENS) and Neuromuscular Electrical Stimulation (NMES) (160.13)

**Note:** To review the specific NCD, please remember to click "accept" on the CMS licensing agreement at the bottom of the CMS webpage.

## Prior Authorization Information

### Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

### Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	Prior authorization is <b>not required</b> .
Commercial PPO and Indemnity	Prior authorization is <b>not required</b> .
Medicare HMO Blue <sup>SM</sup>	Prior authorization is <b>not required</b> .
Medicare PPO Blue <sup>SM</sup>	Prior authorization is <b>not required</b> .

## CPT Codes / HCPCS Codes / ICD Codes

*Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.*

*Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.*

**The above medical necessity criteria MUST be met for the following codes to be covered for Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity:**

### HCPCS Codes

HCPCS codes:	Code Description
A4595	Electrical stimulator supplies, 2 lead, per month, (e.g., TENS, NMES)
A4630	Replacement batteries, medically necessary, transcutaneous electrical stimulator, owned by patient
E0720	Transcutaneous electrical nerve stimulator (TENS) device, two lead, localized stimulation
E0730	Transcutaneous electrical nerve stimulation (TENS) device, four or more leads, for multiple nerve stimulation
E0731	Form-fitting conductive garment for delivery of TENS or NMES (with conductive fibers separated from the patient's skin by layers of fabric)

**The following CPT code is considered investigational for Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:**

### CPT Codes

CPT codes:	Code Description
0278T	Transcutaneous electrical modulation pain reprocessing (eg, scrambler therapy), each treatment session (includes placement of electrodes)

### Description

Transcutaneous electrical nerve stimulation (TENS) has been used to treat chronic intractable pain, postsurgical pain, and pain associated with active or posttrauma injury unresponsive to other standard pain therapies. It has been proposed that TENS may provide pain relief through the release of endorphins in addition to potential blockade of local pain pathways. TENS has also been used to treat dementia by altering neurotransmitter activity and increasing brain activity that is thought to reduce neural degeneration and stimulate regenerative processes.

Percutaneous electrical nerve stimulation (see policy #[172](#)) is similar to TENS but uses microneedles that penetrate the skin instead of surface electrodes. Interferential stimulation (see policy #[509](#)) uses a modulated waveform for deeper tissue stimulation, and the stimulation is believed to improve blood flow to the affected area.

### Summary

Transcutaneous electrical nerve stimulation (TENS) describes the application of electrical stimulation to the surface of the skin at the site of pain. In addition to more traditional settings such as a physician's office or an outpatient clinic, TENS can be self-administered in a patient's home.

For individuals who have chronic pain (eg, musculoskeletal, neuropathic, and mixed pain conditions) who receive TENS, the evidence includes numerous randomized controlled trials (RCTs) and systematic reviews. Relevant outcomes are symptoms, functional outcomes, quality of life, and medication use. The overall strength of the evidence is weak. The best evidence exists for treatment of chronic, intractable pain. Available evidence indicates that TENS can improve chronic intractable pain in some patients, and

there is support for its use in clinical guidelines by specialty societies. To best direct TENS toward patients who will benefit, a short-term trial of TENS is appropriate, with continuation only in patients who show an initial improvement. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

Clinical input obtained in 2009 and 2011 was, for the most part, in agreement that TENS is a generally accepted treatment modality and can be beneficial for the management of chronic pain in some patients. A trial period, similar to that listed in Medicare Coverage guidelines, was recommended by some.

For individuals who have acute pain (eg, surgical, musculoskeletal, labor, and mixed pain conditions) who receive TENS, the evidence includes RCTs and systematic reviews. Relevant outcomes are symptoms and medication use. Overall, evidence for the use of TENS from high-quality trials remains inconclusive for most indications. A Cochrane review of TENS for acute pain (eg, cervical laser treatment, venipuncture, screening flexible sigmoidoscopy, postpartum uterine contractions, rib fractures) found some evidence that TENS reduces pain intensity over and above that seen with placebo, but the high risk of bias made definitive conclusions impossible. For the treatment of pain after total knee arthroplasty, 2 large RCTs found no benefit of TENS compared with sham TENS. A subsequent systematic review found that TENS reduced pain in the immediate postoperative period (24 hours) after total knee arthroplasty compared with control intervention, however, neither the intensity nor optimal duration time for TENS have been established. For the prevention of migraine headaches, a small RCT reported a greater proportion of patients achieving at least a 50% reduction in migraines with TENS than with sham placebo, and modest reductions in the number of total headache and migraine days. This manufacturer-sponsored trial needs corroboration before conclusions can be made about the efficacy of TENS for preventing migraine headaches. For the relief of pain during office-based hysteroscopy, an RCT found decreased pain and higher patient satisfaction in patients receiving TENS compared with placebo or control. The evidence is insufficient to determine the effects of the technology on health outcomes.

Clinical input obtained in 2009 was generally in agreement that TENS is investigational for the management of acute pain and for other conditions such as dementia.

## Policy History

Date	Action
1/2019	BCBSA National medical policy review. Description, summary and references updated. Policy statements unchanged. Clarified coding information.
1/2018	New references added from BCBSA National medical policy.
1/2018	Clarified coding information.
8/2016	Clarified coding information.
12/2015	New references added from BCBSA National medical policy.
12/2015	Added coding language.
6/2015	New references added from BCBSA National medical policy.
9/2014	BCBSA National medical policy review. New investigational indications described. Coding information clarified. Effective 9/1/2014.
5/2014	Updated Coding section with ICD10 procedure and diagnosis codes. Effective 10/2015.
12/2013	Medically necessary indications clarified.
10/2013	New references from BCBSA National medical policy.
11/2011-4/2012	Medical policy ICD 10 remediation: Formatting, editing and coding updates. No changes to policy statements.
1/2012	Reviewed - Medical Policy Group - Neurology and Neurosurgery. No changes to policy statements.
6/2011	Reviewed - Medical Policy Group - Orthopedics, Rehabilitation Medicine and Rheumatology. No changes to policy statements.
1/2011	Reviewed - Medical Policy Group - Neurology and Neurosurgery. No changes to policy statements.

7/2010	Reviewed - Medical Policy Group - Orthopedics, Rehabilitation Medicine and Rheumatology. No changes to policy statements.
3/2010	BCBSA National medical policy review. Changes to policy statements.
2/2010	BCBSA National medical policy review. Changes to policy statements.
7/2009	BCBSA National medical policy review. Changes to policy statements.

## Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

## References

1. Food and Drug Administration. De Novo Classification Request for Cefaly Device. 2012; [https://www.accessdata.fda.gov/cdrh\\_docs/reviews/K122566.pdf](https://www.accessdata.fda.gov/cdrh_docs/reviews/K122566.pdf). Accessed October 26, 2018.
2. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). TENS or PENS in the treatment of chronic and postoperative pain. *TEC Assessments*. 1996;Volume 11, Tab 21.
3. Bronfort G, Nilsson N, Haas M, et al. Non-invasive physical treatments for chronic/recurrent headache. *Cochrane Database Syst Rev*. Jul 2004(3):CD001878. PMID 15266458
4. Brosseau L, Judd MG, Marchand S, et al. Transcutaneous electrical nerve stimulation (TENS) for the treatment of rheumatoid arthritis in the hand. *Cochrane Database Syst Rev*. Aug 2003(3):CD004377. PMID 12918009
5. Brosseau LU, Pelland LU, Casimiro LY, et al. Electrical stimulation for the treatment of rheumatoid arthritis. *Cochrane Database Syst Rev*. Jun 2002(2):CD003687. PMID 12076504
6. Cameron M, Lonergan E, Lee H. Transcutaneous electrical nerve stimulation (TENS) for dementia. *Cochrane Database Syst Rev*. Aug 2003(3):CD004032. PMID 12917999
7. Carroll D, Moore RA, McQuay HJ, et al. Transcutaneous electrical nerve stimulation (TENS) for chronic pain. *Cochrane Database Syst Rev*. Nov 2001(3):CD003222. PMID 11687055
8. Dowswell T, Bedwell C, Lavender T, et al. Transcutaneous electrical nerve stimulation (TENS) for pain relief in labour. *Cochrane Database Syst Rev*. Apr 15 2009(2):CD007214. PMID 19370680
9. Hurlow A, Bennett MI, Robb KA, et al. Transcutaneous electric nerve stimulation (TENS) for cancer pain in adults. *Cochrane Database Syst Rev*. Mar 14 2012;3(3):CD006276. PMID 22419313
10. Khadilkar A, Milne S, Brosseau L, et al. Transcutaneous electrical nerve stimulation (TENS) for chronic low-back pain. *Cochrane Database Syst Rev*. Jul 20 2005(3):CD003008. PMID 16034883
11. Khadilkar A, Odebiyi DO, Brosseau L, et al. Transcutaneous electrical nerve stimulation (TENS) versus placebo for chronic low-back pain. *Cochrane Database Syst Rev*. Oct 08 2008(4):CD003008. PMID 18843638
12. Kroeling P, Gross A, Goldsmith CH, et al. Electrotherapy for neck pain. *Cochrane Database Syst Rev*. Oct 07 2009(4):CD004251. PMID 19821322
13. Kroeling P, Gross A, Graham N, et al. Electrotherapy for neck pain. *Cochrane Database Syst Rev*. Aug 26 2013;8(8):CD004251. PMID 23979926
14. Kroeling P, Gross A, Houghton PE, et al. Electrotherapy for neck disorders. *Cochrane Database Syst Rev*. Apr 18 2005(2):CD004251. PMID 15846703
15. Milne S, Welch V, Brosseau L, et al. Transcutaneous electrical nerve stimulation (TENS) for chronic low back pain. *Cochrane Database Syst Rev*. Jun 2001(2):CD003008. PMID 11406059
16. Mulvey MR, Bagnall AM, Johnson MI, et al. Transcutaneous electrical nerve stimulation (TENS) for phantom pain and stump pain following amputation in adults. *Cochrane Database Syst Rev*. May 12 2010(5):CD007264. PMID 20464749
17. Nnoaham KE, Kumbang J. Transcutaneous electrical nerve stimulation (TENS) for chronic pain. *Cochrane Database Syst Rev*. Jul 16 2008(3):CD003222. PMID 18646088
18. Osiri M, Welch V, Brosseau L, et al. Transcutaneous electrical nerve stimulation for knee osteoarthritis. *Cochrane Database Syst Rev*. Oct 2000(4):CD002823. PMID 11034768
19. Price CI, Pandyan AD. Electrical stimulation for preventing and treating post-stroke shoulder pain. *Cochrane Database Syst Rev*. Oct 2000(4):CD001698. PMID 11034725

20. Proctor ML, Smith CA, Farquhar CM, et al. Transcutaneous electrical nerve stimulation and acupuncture for primary dysmenorrhoea. *Cochrane Database Syst Rev*. Mar 2002(1):CD002123. PMID 11869624
21. Robb KA, Bennett MI, Johnson MI, et al. Transcutaneous electric nerve stimulation (TENS) for cancer pain in adults. *Cochrane Database Syst Rev*. Jul 16 2008(3):CD006276. PMID 18646140
22. Rutjes AW, Nuesch E, Sterchi R, et al. Transcutaneous electrostimulation for osteoarthritis of the knee. *Cochrane Database Syst Rev*. Oct 07 2009(4):CD002823. PMID 19821296
23. Walsh DM, Howe TE, Johnson MI, et al. Transcutaneous electrical nerve stimulation for acute pain. *Cochrane Database Syst Rev*. Apr 15 2009(2):CD006142. PMID 19370629
24. Dubinsky RM, Miyasaki J. Assessment: efficacy of transcutaneous electric nerve stimulation in the treatment of pain in neurologic disorders (an evidence-based review): report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology*. Jan 12 2010;74(2):173-176. PMID 20042705
25. Wu LC, Weng PW, Chen CH, et al. Literature review and meta-analysis of transcutaneous electrical nerve stimulation in treating chronic back pain. *Reg Anesth Pain Med*. May 2018;43(4):425-433. PMID 29394211
26. Keskin EA, Onur O, Keskin HL, et al. Transcutaneous electrical nerve stimulation improves low back pain during pregnancy. *Gynecol Obstet Invest*. Jun 2012;74(1):76-83. PMID 22722614
27. Chen FC, Jin ZL, Wang DF. A retrospective study of transcutaneous electrical nerve stimulation for chronic pain following ankylosing spondylitis. *Medicine (Baltimore)*. Jul 2018;97(27):e11265. PMID 29979392
28. Kong X, Gozani SN. Effectiveness of fixed-site high-frequency transcutaneous electrical nerve stimulation in chronic pain: a large-scale, observational study. *J Pain Res*. 2018;11:703-714. PMID 29670397
29. Gossrau G, Wahner M, Kuschke M, et al. Microcurrent transcutaneous electric nerve stimulation in painful diabetic neuropathy: a randomized placebo-controlled study. *Pain Med*. Jun 2011;12(6):953-960. PMID 21627767
30. Dailey DL, Rakel BA, Vance CG, et al. Transcutaneous electrical nerve stimulation reduces pain, fatigue and hyperalgesia while restoring central inhibition in primary fibromyalgia. *Pain*. Nov 2013;154(11):2554-2562. PMID 23900134
31. Lauretti GR, Chubaci EF, Mattos AL. Efficacy of the use of two simultaneously TENS devices for fibromyalgia pain. *Rheumatol Int*. Aug 2013;33(8):2117-2122. PMID 23423539
32. Schneider MP, Tellenbach M, Mordasini L, et al. Refractory chronic pelvic pain syndrome in men: can transcutaneous electrical nerve stimulation help? *BJU Int*. Jul 2013;112(2):E159-163. PMID 23433012
33. Cherian JJ, Harrison PE, Benjamin SA, et al. Do the effects of transcutaneous electrical nerve stimulation on knee osteoarthritis pain and function last? *J Knee Surg*. Aug 2016;29(6):497-501. PMID 26540652
34. Palmer S, Domaille M, Cramp F, et al. Transcutaneous electrical nerve stimulation as an adjunct to education and exercise for knee osteoarthritis: a randomized controlled trial. *Arthritis Care Res*. Mar 2014;66(3):387-394. PMID 23983090
35. Vance CG, Rakel BA, Blodgett NP, et al. Effects of transcutaneous electrical nerve stimulation on pain, pain sensitivity, and function in people with knee osteoarthritis: a randomized controlled trial. *Phys Ther*. Jul 2012;92(7):898-910. PMID 22466027
36. Chen WL, Hsu WC, Lin YJ, et al. Comparison of intra-articular hyaluronic acid injections with transcutaneous electric nerve stimulation for the management of knee osteoarthritis: a randomized controlled trial. *Arch Phys Med Rehabil*. Aug 2013;94(8):1482-1489. PMID 23628378
37. Sawant A, Dadurka K, Overend T, et al. Systematic review of efficacy of TENS for management of central pain in people with multiple sclerosis. *Mult Scler Relat Disord*. May 2015;4(3):219-227. PMID 26008938
38. Johnson MI, Mulvey MR, Bagnall AM. Transcutaneous electrical nerve stimulation (TENS) for phantom pain and stump pain following amputation in adults. *Cochrane Database Syst Rev*. Aug 18 2015;8:CD007264. PMID 26284511
39. Boldt I, Eriks-Hoogland I, Brinkhof MW, et al. Non-pharmacological interventions for chronic pain in people with spinal cord injury. *Cochrane Database Syst Rev*. 2014;11:CD009177. PMID 25432061

40. Schoenen J, Vandersmissen B, Jeanette S, et al. Migraine prevention with a supraorbital transcutaneous stimulator: a randomized controlled trial. *Neurology*. Feb 19 2013;80(8):697-704. PMID 23390177
41. Magis D, Sava S, d'Elia TS, et al. Safety and patients' satisfaction of transcutaneous supraorbital neurostimulation (tSNS) with the Cefaly(R) device in headache treatment: a survey of 2,313 headache sufferers in the general population. *J Headache Pain*. Dec 2013;14:95. PMID 24289825
42. De Giorgi I, Castroflorio T, Sartoris B, et al. The use of conventional transcutaneous electrical nerve stimulation in chronic facial myalgia patients. *Clin Oral Investig*. Jan 2017;21(1):275-280. PMID 27000071
43. Ferreira AP, Costa DR, Oliveira AI, et al. Short-term transcutaneous electrical nerve stimulation reduces pain and improves the masticatory muscle activity in temporomandibular disorder patients: a randomized controlled trial. *J Appl Oral Sci*. Mar-Apr 2017;25(2):112-120. PMID 28403351
44. Johnson M, Martinson M. Efficacy of electrical nerve stimulation for chronic musculoskeletal pain: a meta-analysis of randomized controlled trials. *Pain*. Jul 2007;130(1-2):157-165. PMID 17383095
45. Deyo RA, Walsh NE, Martin DC, et al. A controlled trial of transcutaneous electrical nerve stimulation (TENS) and exercise for chronic low back pain. *N Engl J Med*. Jun 7 1990;322(23):1627-1634. PMID 2140432
46. Machin D, Lewith GT, Wylson S. Pain measurement in randomized clinical trials: A comparison of two pain scales. *Clin J Pain*. 1988;4:161-168.
47. Oosterhof J, De Boo TM, Oostendorp RA, et al. Outcome of transcutaneous electrical nerve stimulation in chronic pain: short-term results of a double-blind, randomised, placebo-controlled trial. *J Headache Pain*. Sep 2006;7(4):196-205. PMID 16897618
48. Oosterhof J, Samwel HJ, de Boo TM, et al. Predicting outcome of TENS in chronic pain: a prospective, randomized, placebo controlled trial. *Pain*. May 2008;136(1-2):11-20. PMID 17659838
49. Oosterhof J, Wilder-Smith OH, de Boo T, et al. The long-term outcome of transcutaneous electrical nerve stimulation in the treatment for patients with chronic pain: a randomized, placebo-controlled trial. *Pain Pract*. Sep 2012;12(7):513-522. PMID 22304690
50. Lang T, Barker R, Steinlechner B, et al. TENS relieves acute posttraumatic hip pain during emergency transport. *J Trauma*. Jan 2007;62(1):184-188; discussion 188. PMID 17215752
51. Zhu Y, Feng Y, Peng L. Effect of transcutaneous electrical nerve stimulation for pain control after total knee arthroplasty: A systematic review and meta-analysis. *J Rehabil Med*. Nov 21 2017;49(9):700-704. PMID 28933513
52. Rakel BA, Zimmerman MB, Geasland K, et al. Transcutaneous electrical nerve stimulation for the control of pain during rehabilitation after total knee arthroplasty: A randomized, blinded, placebo-controlled trial. *Pain*. Dec 2014;155(12):2599-2611. PMID 25270585
53. Ramanathan D, Saleh A, Klika AK, et al. The use of transcutaneous electrical nerve stimulation after total knee arthroplasty: a prospective randomized controlled trial. *Surg Technol Int*. Jul 25 2017;30:425-434. PMID 28537354
54. Galli TT, Chiavegato LD, Lievano RE. Effects of TENS in living kidney donors submitted to open nephrectomy: a randomized placebo-controlled trial. *Eur J Pain*. Jan 2015;19(1):67-76. PMID 24831862
55. Tokuda M, Tabira K, Masuda T, et al. Effect of modulated-frequency and modulated-intensity transcutaneous electrical nerve stimulation after abdominal surgery: a randomized controlled trial. *Clin J Pain*. Jul 2014;30(7):565-570. PMID 24901753
56. Silva MB, de Melo PR, de Oliveira NM, et al. Analgesic effect of transcutaneous electrical nerve stimulation after laparoscopic cholecystectomy. *Am J Phys Med Rehabil*. Aug 2012;91(8):652-657. PMID 22311059
57. DeSantana JM, Walsh DM, Vance C, et al. Effectiveness of transcutaneous electrical nerve stimulation for treatment of hyperalgesia and pain. *Curr Rheumatol Rep*. Dec 2008;10(6):492-499. PMID 19007541
58. Tucker DL, Rockett M, Hasan M, et al. Does transcutaneous electrical nerve stimulation (TENS) alleviate the pain experienced during bone marrow sampling in addition to standard techniques? A randomised, double-blinded, controlled trial. *J Clin Pathol*. Jun 2015;68(6):479-483. PMID 25759407
59. Lison JF, Amer-Cuenca JJ, Piquer-Marti S, et al. Transcutaneous nerve stimulation for pain relief during office hysteroscopy: a randomized controlled trial. *Obstet Gynecol*. Feb 2017;129(2):363-370. PMID 28079781

60. Kayman-Kose S, Arioz DT, Toktas H, et al. Transcutaneous electrical nerve stimulation (TENS) for pain control after vaginal delivery and cesarean section. *J Matern Fetal Neonatal Med.* Oct 2014;27(15):1572-1575. PMID 24283391
61. Johnson MI, Paley CA, Howe TE, et al. Transcutaneous electrical nerve stimulation for acute pain. *Cochrane Database Syst Rev.* Jun 15 2015;6(6):CD006142. PMID 26075732
62. Simpson PM, Fouche PF, Thomas RE, et al. Transcutaneous electrical nerve stimulation for relieving acute pain in the prehospital setting: a systematic review and meta-analysis of randomized-controlled trials. *Eur J Emerg Med.* Feb 2014;21(1):10-17. PMID 23839103
63. Butera KA, George SZ, Borsa PA, et al. Prolonged reduction in shoulder strength after transcutaneous electrical nerve stimulation treatment of exercise-induced acute muscle pain. *Pain Pract.* Mar 5 2018. PMID 29505689
64. Chesterton LS, Lewis AM, Sim J, et al. Transcutaneous electrical nerve stimulation as adjunct to primary care management for tennis elbow: pragmatic randomised controlled trial (TATE trial). *BMJ.* Sep 02 2013;347:f5160. PMID 23999980
65. Martelletti P, Jensen RH, Antal A, et al. Neuromodulation of chronic headaches: position statement from the European Headache Federation. *J Headache Pain.* Oct 21 2013;14(1):86. PMID 24144382
66. McAlindon TE, Bannuru RR, Sullivan MC, et al. OARSI guidelines for the non-surgical management of knee osteoarthritis. *Osteoarthritis Cartilage.* Mar 2014;22(3):363-388. PMID 24462672
67. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Adult Cancer Pain. Version 1.2018. [https://www.nccn.org/professionals/physician\\_gls/pdf/pain.pdf](https://www.nccn.org/professionals/physician_gls/pdf/pain.pdf). Accessed October 26, 2018.
68. National Cancer Institute. Pain (PDQ®)-Health Professional Version. 2018; <https://www.cancer.gov/about-cancer/treatment/side-effects/pain/pain-hp-pdq#section/all?redirect=true>. Accessed October 26, 2018.
69. Bono CM, Ghiselli G, Gilbert TJ, et al. An evidence-based clinical guideline for the diagnosis and treatment of cervical radiculopathy from degenerative disorders. *Spine J.* Jan 2011;11(1):64-72. PMID 21168100
70. American Society of Anesthesiologists Task Force on Chronic Pain Management, American Society of Regional Anesthesia Pain Medicine. Practice guidelines for chronic pain management: an updated report by the American Society of Anesthesiologists Task Force on Chronic Pain Management and the American Society of Regional Anesthesia and Pain Medicine. *Anesthesiology.* Apr 2010;112(4):810-833. PMID 20124882
71. National Institute for Health and Care Excellence (NICE). Low back pain and sciatica in over 16s: assessment and management [NG59]. 2016; <https://www.nice.org.uk/guidance/NG59>. Accessed October 26, 2018.
72. National Institute for Health and Care Excellence (NICE). Osteoarthritis: care and management [CG177]. 2014; <https://www.nice.org.uk/guidance/cg177>. Accessed October 26, 2018.
73. National Institute for Health and Care Excellence (NICE). Intrapartum care for healthy women and babies [CP190]. 2017; <https://www.nice.org.uk/guidance/cg190>. Accessed October 26, 2018.
74. American Congress of Obstetricians and Gynecologists (ACOG). Women's Health Care Physicians. 2007; <http://www.acog.org/Resources-And-Publications/>. Accessed October 26, 2018.
75. ACOG Committee on Practice Bulletins--Gynecology. ACOG Practice Bulletin No. 51. Chronic pelvic pain. *Obstet Gynecol.* Mar 2004;103(3):589-605. PMID 14990428
76. Committee on Obstetric Practice. Committee Opinion No. 687: Approaches to limit intervention during labor and birth. *Obstet Gynecol.* Feb 2017;129(2):e20-e28. PMID 28121831
77. Qaseem A, Wilt TJ, McLean RM, et al. Noninvasive treatments for acute, subacute, and chronic low back pain: a clinical practice guideline from the American College of Physicians. *Ann Intern Med.* Apr 04 2017;166(7):514-530. PMID 28192789
78. Cruccu G, Aziz TZ, Garcia-Larrea L, et al. EFNS guidelines on neurostimulation therapy for neuropathic pain. *Eur J Neurol.* Sep 2007;14(9):952-970. PMID 17718686
79. Centers for Medicare & Medicaid. National Coverage Determination (NCD) for Supplies Used in the Delivery of Transcutaneous Electrical Nerve Stimulation (TENS) and Neuromuscular Electrical Stimulation (NMES) (160.13). 1988; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=151&ncdver=1&DocID=160.13&bc=gAAAABAAAAAAA%3d%3d&>. Accessed October 26, 2018.
80. Centers for Medicare & Medicaid. National Coverage Determination (NCD) for Transcutaneous Electrical Nerve Stimulators (TENS) (280.13). 1995; <https://www.cms.gov/medicare-coverage->



database/details/ncd-  
details.aspx?NCDId=273&ncdver=1&DocID=280.13&bc=gAAAAABAAAAAAAA%3d%3d&. Accessed  
October 26, 2018.

81. Centers for Medicare & Medicaid. National Coverage Determination (NCD) for Transcutaneous Electrical Nerve Stimulation (TENS) for Acute Post-Operative Pain (10.2). 1995; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?ncdid=145&ver=1>. Accessed October 26, 2018.
82. Centers for Medicare & Medicaid. National Coverage Determination (NCD) for Assessing Patient's Suitability for Electrical Nerve Stimulation Therapy (160.7.1). 2006; [https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?TAId=4&MEDCACId=15&MCDId=10&CALId=97&NCDId=63&ncdver=2&CoverageSelection=National&ncd\\_id=10.1&ncd\\_version=1&basket=ncd%25253A10%25252E1%25253A1%25253AUse+of+Visual+Tests+Prior+to+and+General+Anesthesia+during+Cataract+Surgery&bc=gAAAAABAAAAAAAA%3D%3D&](https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?TAId=4&MEDCACId=15&MCDId=10&CALId=97&NCDId=63&ncdver=2&CoverageSelection=National&ncd_id=10.1&ncd_version=1&basket=ncd%25253A10%25252E1%25253A1%25253AUse+of+Visual+Tests+Prior+to+and+General+Anesthesia+during+Cataract+Surgery&bc=gAAAAABAAAAAAAA%3D%3D&). Accessed October 26, 2018.
83. Centers for Medicare & Medicaid. Decision Memo for Transcutaneous Electrical Nerve Stimulation for Chronic Low Back Pain (CAG-00429N). 2012; <https://www.cms.gov/medicare-coverage-database/details/nca-decision-memo.aspx?NCAId=256&NcaName=Transcutaneous+Electrical+Nerve+Stimulation+for+Chronic+Low+Back+Pain&TimeFrame=7&DocType=All&bc=AQAAlAAAlAAA&>. Accessed October 26, 2018