



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

Pharmacy Medical Policy Botulinum Toxin Injections

Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Coding Information](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)
- [Endnotes](#)
- [Forms](#)

Policy Number: 006

BCBSA Reference Number: 5.01.05 & 8.01.19

Related Policies

Treatment of Hyperhidrosis, #406

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity

Note: All requests for outpatient retail pharmacy for indications listed and not listed on the medical policy guidelines may be submitted to BCBSMA Clinical Pharmacy Operations by completing the Prior Authorization Form on the last page of this document. Physicians may also call BCBSMA Pharmacy Operations department at (800)366-7778 to request a prior authorization/formulary exception verbally. Physicians may also submit requests for retail pharmacy exceptions via the web using Express PAtH which can be found on the BCBSMA provider website or directly on the web at <https://provider.express-path.com>. Patients must have pharmacy benefits under their subscriber certificates.

Please refer to the chart below for the formulary status of the medications affected by this policy.

Drug	Formulary Information
	Standard
	Formulary Status
Botox™ [^] (onabotulinumtoxin a)	Preferred with PA
Dysport™ [^] (botulinum toxin type a)	Preferred with PA
Myobloc™ [^] (rimabotulinumtoxin b)	Non-Preferred with PA
Xeomin® [^] (incobotulinumtoxin a)	Non-Preferred with PA

[^] - This Drug is part of Medications covered only under the pharmacy benefit program.

Note: To obtain a Non-Preferred toxin one must try and fail at least one preferred toxin.

Dysport™ is required to be used prior to Botox™ for the following indications only:

- For upper limb spasticity in adult patients
- For lower limb spasticity in patients 2 years or older
- for cervical dystonia in adults

We may cover the following indications for **Dysport**[™] (botulinum toxin type a) and **Botox**[™] (onabotulinumtoxin a) in this policy which are FDA approved indications for the Botulinum Toxins and we will cover a **Myobloc**[™] (rimabotulinumtoxin b) or **Xeomin**[®] (incobotulinumtoxin a) when either **Dysport**[™] or **Botox**[™] is tried and failed first:

- Is indicated for the treatment of upper limb spasticity in adult patients, to decrease the severity of increased muscle tone in elbow flexors (biceps), wrist flexors (flexor carpi radialis and flexor carpi ulnaris), finger flexors (flexor digitorum profundus and flexor digitorum sublimis), Focal upper limb dystonia (organic writer's cramp), and thumb flexors (adductor pollicis and flexor pollicis longus) when **ALL** of the following criteria are met:
 - Age 18 years or over
 - AND**
 - **Dysport**[™] (botulinum toxin type a) must be used prior to **Botox**[™] (onabotulinumtoxin a), **Myobloc**[™] (rimabotulinumtoxin b) or **Xeomin**[®] (incobotulinumtoxin a)
- A lower limb spasticity in patients 2 years or older to decrease the severity of increased muscle tone in ankle and toe flexors (gastrocnemius, soleus, tibialis posterior, flexor hallucis longus, and flexor digitorum longus). **ALL** of the following criteria are met:
 - Age 2 years or over
 - AND**
 - **Dysport**[™] (botulinum toxin type a) must be used prior to **Botox**[™] (onabotulinumtoxin a), **Myobloc**[™] (rimabotulinumtoxin b) or **Xeomin**[®] (incobotulinumtoxin a).
- Is indicated for the treatment of adults with cervical dystonia, to reduce the severity of abnormal head position and neck pain associated with cervical dystonia. For this use, cervical dystonia must be associated with sustained head tilt or abnormal posturing with limited range of motion in the neck **AND** a history of recurrent involuntary contraction of one or more of the muscles of the neck, (e.g., sternocleidomastoid, splenius, trapezius, or posterior cervical muscles) and may be covered when **ALL** of the following criteria are met:
 - Age 18 years or over
 - AND**
 - **Dysport**[™] (botulinum toxin type a) must be used prior to **Botox**[™] (onabotulinumtoxin a), **Myobloc**[™] (rimabotulinumtoxin b) or **Xeomin**[®] (incobotulinumtoxin a)
- Is indicated for the treatment of overactive bladder with symptoms of urge urinary incontinence, urgency, and frequency, in adults who have an inadequate response to or are intolerant of an anticholinergic medication
- Is indicated for the treatment of urinary incontinence due to detrusor over activity associated with a neurologic condition (e.g., SCI, MS) in adults who have an inadequate response to or are intolerant of an anticholinergic medication
- Is indicated for the prophylaxis of headaches in adult patients with chronic migraine (≥15 days per month with headache lasting 4 hours a day or longer)
- And will be covered for **Migraine headache** when **ALL** of the following criteria are met:
 - Age 18 years or over
 - Prescribed by a neurologist, ophthalmologist or board certified headache medicine specialist.
 - Episodes of migraine for ≥ 15 days/month with duration ≥ 4 hours/day
 - Previous treatment for at least three months each or contraindication to all of the following therapeutic categories/medications:
 - Beta blockers (e.g. propranolol, timolol)
 - Topiramate
 - Divalproex sodium
 - Non-steroidal anti-inflammatory medications (e.g. ibuprofen, naproxen, diclofenac)
 - Serotonin receptor agonists (e.g. sumatriptan, naratriptan).
- Is indicated for the treatment of strabismus and blepharospasm associated with dystonia, including benign essential blepharospasm or facial (VII) nerve disorders such as hemifacial spasm in patients 12 years of age and above
- Chronic sialorrhea

- Is indicated for the treatment of severe primary axillary hyperhidrosis that is inadequately managed with topical agents and will be covered according to the criteria below.

NOTE: Primary focal hyperhidrosis is defined as excessive sweating induced by sympathetic hyperactivity in selected areas that is not associated with an underlying disease process. The most common locations are underarms (axillary hyperhidrosis), palms (palmar hyperhidrosis), soles (plantar hyperhidrosis) or face (craniofacial hyperhidrosis).

- We may cover **the treatment of primary hyperhidrosis in a small subset of patients** with the following medical conditions:
 - acrocyanosis of the hands;
 - history of recurrent skin maceration with bacterial or fungal infections;
 - history of recurrent secondary infections;
 - history of persistent eczematous dermatitis in spite of medical treatments with topical dermatological or systemic anticholinergic agents; or
 - significant functional impairment:
 - Documentation must be submitted that reports the location of the hyperhidrosis, the frequency and duration of episodes, the specific functions that are impaired (including activities of daily living and/or occupational activities), the severity of impairment, and a description of how the function is impaired.

We cover **the treatment of primary hyperhidrosis based on focal regions** as noted below:

Focal Regions	Covered Treatments
Axillary	Onabotulinumtoxin A (botulinum type A)(intradermal injection) for severe primary axillary hyperhidrosis that is inadequately managed with topical agents, in patients 18 years and older,
Palmar	Onabotulinumtoxin A (botulinum type A) (intradermal injection) for severe primary palmar hyperhidrosis that is inadequately managed with topical agents, in patients 18 years and older;

We also may cover the following Dystonia/Spasticity disorders:

Note: To obtain a Non-Preferred toxin one must try and fail at least one Preferred toxin. In addition, Dysport shall be used prior to Botox, Xeomin & Myobloc for any diagnosis involving spasticity in adults.

Dystonia/spasticity resulting in functional impairment (interference with joint function, mobility, communication, nutritional intake) and/or pain in patients with **any** of the following:

- Focal upper limb dystonia (e.g., organic writer’s cramp)
- Oromandibular dystonia (orofacial dyskinesia, Meige syndrome)
- Laryngeal dystonia (adductor spasmodic dysphonia)
- Idiopathic (primary or genetic) torsion dystonia
- Symptomatic (acquired) torsion dystonia
- Cerebral palsy
- Spasticity related to stroke
- Acquired spinal cord or brain injury
- Hereditary spastic paraparesis
- Spastic hemiplegia
- Neuromyelitis optica
- Multiple sclerosis or Schilder’s disease
- Esophageal achalasia in patients who have not responded to dilation therapy or who are considered poor surgical candidates
- Chronic anal fissure

We do not cover onabotulinumtoxin A (**Botox™**), onabotulinumtoxin A (**Dysport™**) rimabotulinumtoxin B (**Myobloc™**) or incobotulinumtoxin A (**Xeomin®**) injections for conditions, including but not limited to:

- Eye conditions not listed above, including:
 - Chronic paralytic strabismus (except to reduce antagonist contracture in conjuncture with surgical repair)
 - Patients with corneal exposure, persistent epithelial defect, or corneal ulceration
- Headaches including migraine unless criteria met above
- Tourette’s syndrome
- Chronic Motor Tic disorder
- Patients with myasthenia gravis
- Wrinkles, glabellar lines or other cosmetic indications
- Myofascial pain syndrome
- Chronic low back pain
- Tremors such as benign essential tremor
- Lateral epicondylitis
- Benign prostatic hyperplasia
- Detrusor over reactivity not due to spinal cord injury
- Detrusor sphincteric dyssynergia
- Prevention of pain associated with breast reconstruction after mastectomy
- Hirschsprung’s disease
- Gastroparesis.

We do not cover **the following botulinum toxin treatments of primary hyperhidrosis** based on focal region, because they are considered investigational, as they do not meet our Medical Technology Assessment Guidelines, #350:

Focal Region	Non Covered Treatments (Investigational)
Palmar	<ul style="list-style-type: none"> • Rimabotulinumtoxin B (botulinum type B)
Plantar	<ul style="list-style-type: none"> • Onabotulinumtoxin A (botulinum type A) • Rimabotulinumtoxin B (botulinum type B)
Craniofacial	<ul style="list-style-type: none"> • Onabotulinumtoxin A (botulinum type A) • Rimabotulinumtoxin B (botulinum type B)

We do not cover **the following treatments** including, but not limited to, **Onabotulinumtoxin A (botulinum toxin type A) and Rimabotulinumtoxin B (botulinum toxin type B)** as a treatment for **severe gustatory hyperhidrosis¹** because they are considered investigational, as they do not meet our Medical Technology Assessment Guidelines, #350.

For patient safety, we do not cover any type of botulinum injections for:

- Patients who are pregnant or intend to become pregnant
- Patients who are on aminoglycoside therapy, as it may increase the risk of problems between the muscles and the nerves
- Patients with retrobulbar hemorrhages sufficient to compromise retinal circulation
- Patients with severe laryngeal or respiratory weakness
- Patients with sensitivity or allergy to any type of botulinum injections, or known high antibody titers to any type of botulinum injections.

Other Information

Blue Cross Blue Shield of Massachusetts (BCBSMA*) members (other than Medex®; Blue MedicareRx, Medicare Advantage plans that include prescription drug coverage) will be required to fill their prescriptions for the above medications at one of the providers in our retail specialty pharmacy network, see link below:

[Link to Specialty Pharmacy List](#)

Individual Consideration

All our medical policies are written for the majority of people with a given condition. Each policy is based on medical science. For many of our medical policies, each individual's unique clinical circumstances may be considered in light of current scientific literature. Physicians may send relevant clinical information for individual patients for consideration to:

Blue Cross Blue Shield of Massachusetts
Pharmacy Operations Department
25 Technology Place
Hingham, MA 02043
Tel: 1-800-366-7778
Fax: 1-800-583-6289

Managed Care Authorization Instructions

- Prior authorization is required for all out patient sites of service
- For retail pharmacy requests, physicians may call BCBSMA Pharmacy Operations department to request a review for prior authorization for patients.
Pharmacy Operations: (800)366-7778
- For all outpatient sites of service for retail pharmacy exceptions, physicians may also fax or mail the attached form to the address above. The Formulary Exception/Prior Authorization form is included as part of this document for physicians to submit for patients.
- For all outpatient sites of service for retail pharmacy exceptions, physicians may also submit requests for retail pharmacy exceptions via the web using Express PAtH which can be found on the BCBSMA provider website or directly on the web at <https://provider.express-path.com>

PPO and Indemnity Authorization Instructions

- Prior authorization **is** required when these medications are processed under the retail pharmacy benefit and home infusion therapy benefit.
- Prior authorization **is not** required when drugs are not part of the Pharmacy only program and are purchased by the physician and administered in the office in accordance with this medical policy.
- For retail pharmacy requests, physicians may call BCBSMA Pharmacy Operations department to request a review for prior authorization for patients.
Pharmacy Operations: (800)366-7778
- Physicians may also fax or mail the attached form for retail pharmacy exceptions to the address above. The Formulary Exception/Prior Authorization form is included as part of this document for physicians to submit for patients.
- Physicians may also submit requests for retail pharmacy exceptions via the web using Express PAtH which can be found on the BCBSMA provider website or directly on the web at <https://provider.express-path.com>

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member. A draft of future ICD-10 Coding related to this document, as it might look today, is included below for your reference.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

The above medical necessity criteria **MUST be met for the following codes to be covered for Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity:**

HCPCS Codes

HCPCS codes:	Code Description
J0585	Injection, onabotulinumtoxinA, 1 unit (Botox)
J0587	Injection, rimabotulinumtoxinB, 100 units (Myobloc)

The following ICD Diagnosis Codes are considered medically necessary when submitted with the HCPCS codes above if medical necessity criteria are met:

Diagnosis coding

ICD-10 Diagnosis Codes

ICD-10-CM Diagnosis codes:	Code Description
G11.4	Hereditary spastic paraplegia
G24.02	Drug induced acute dystonia
G24.09	Other drug induced dystonia
G24.1	Genetic torsion dystonia
G24.2	Idiopathic nonfamilial dystonia
G24.3	Spasmodic torticollis
G24.4	Idiopathic orofacial dystonia
G24.5	Blepharospasm
G24.8	Other dystonia
G24.9	Dystonia, unspecified
G25.82	Stiff-man syndrome
G25.89	Other specified extrapyramidal and movement disorders
G35	Multiple sclerosis
G36.0	Neuromyelitis optica [Devic]
G36.1	Acute and subacute hemorrhagic leukoencephalitis [Hurst]
G36.8	Other specified acute disseminated demyelination
G36.9	Acute disseminated demyelination, unspecified
G37.0	Diffuse sclerosis of central nervous system
G37.1	Central demyelination of corpus callosum
G37.2	Central pontine myelinolysis
G37.4	Subacute necrotizing myelitis of central nervous system
G37.5	Concentric sclerosis [Balo] of central nervous system
G37.8	Other specified demyelinating diseases of central nervous system
G37.9	Demyelinating disease of central nervous system, unspecified
G43.001	Migraine without aura, not intractable, with status migrainosus
G43.009	Migraine without aura, not intractable, without status migrainosus
G43.011	Migraine without aura, intractable, with status migrainosus
G43.019	Migraine without aura, intractable, without status migrainosus
G43.101	Migraine with aura, not intractable, with status migrainosus
G43.109	Migraine with aura, not intractable, without status migrainosus
G43.111	Migraine with aura, intractable, with status migrainosus
G43.119	Migraine with aura, intractable, without status migrainosus
G43.401	Hemiplegic migraine, not intractable, with status migrainosus
G43.409	Hemiplegic migraine, not intractable, without status migrainosus
G43.411	Hemiplegic migraine, intractable, with status migrainosus
G43.419	Hemiplegic migraine, intractable, without status migrainosus

G43.501	Persistent migraine aura without cerebral infarction, not intractable, with status migrainosus
G43.509	Persistent migraine aura without cerebral infarction, not intractable, without status migrainosus
G43.511	Persistent migraine aura without cerebral infarction, intractable, with status migrainosus
G43.519	Persistent migraine aura without cerebral infarction, intractable, without status migrainosus
G43.601	Persistent migraine aura with cerebral infarction, not intractable, with status migrainosus
G43.609	Persistent migraine aura with cerebral infarction, not intractable, without status migrainosus
G43.611	Persistent migraine aura with cerebral infarction, intractable, with status migrainosus
G43.619	Persistent migraine aura with cerebral infarction, intractable, without status migrainosus
G43.701	Chronic migraine without aura, not intractable, with status migrainosus
G43.709	Chronic migraine without aura, not intractable, without status migrainosus
G43.711	Chronic migraine without aura, intractable, with status migrainosus
G43.719	Chronic migraine without aura, intractable, without status migrainosus
G43.801	Other migraine, not intractable, with status migrainosus
G43.809	Other migraine, not intractable, without status migrainosus
G43.811	Other migraine, intractable, with status migrainosus
G43.819	Other migraine, intractable, without status migrainosus
G43.821	Menstrual migraine, not intractable, with status migrainosus
G43.829	Menstrual migraine, not intractable, without status migrainosus
G43.831	Menstrual migraine, intractable, with status migrainosus
G43.839	Menstrual migraine, intractable, without status migrainosus
G43.901	Migraine, unspecified, not intractable, with status migrainosus
G43.909	Migraine, unspecified, not intractable, without status migrainosus
G43.911	Migraine, unspecified, intractable, with status migrainosus
G43.919	Migraine, unspecified, intractable, without status migrainosus
G43.A0	Cyclical vomiting, not intractable
G43.A1	Cyclical vomiting, intractable
G43.B0	Ophthalmoplegic migraine, not intractable
G43.B1	Ophthalmoplegic migraine, intractable
G43.C0	Periodic headache syndromes in child or adult, not intractable
G43.C1	Periodic headache syndromes in child or adult, intractable
G43.D0	Abdominal migraine, not intractable
G43.D1	Abdominal migraine, intractable
G44.1	Vascular headache, not elsewhere classified
G51.0	Bell's palsy
G51.1	Geniculate ganglionitis
G51.2	Melkersson's syndrome
G51.3	Clonic hemifacial spasm
G51.4	Facial myokymia
G51.8	Other disorders of facial nerve
G51.9	Disorder of facial nerve, unspecified
G80.0	Spastic quadriplegic cerebral palsy
G80.1	Spastic diplegic cerebral palsy
G80.2	Spastic hemiplegic cerebral palsy
G80.4	Ataxic cerebral palsy
G80.8	Other cerebral palsy
G80.9	Cerebral palsy, unspecified
G81.10	Spastic hemiplegia affecting unspecified side
G81.11	Spastic hemiplegia affecting right dominant side
G81.12	Spastic hemiplegia affecting left dominant side
G81.13	Spastic hemiplegia affecting right nondominant side

G81.14	Spastic hemiplegia affecting left nondominant side
H49.00	Third [oculomotor] nerve palsy, unspecified eye
H49.01	Third [oculomotor] nerve palsy, right eye
H49.02	Third [oculomotor] nerve palsy, left eye
H49.03	Third [oculomotor] nerve palsy, bilateral
H49.10	Fourth [trochlear] nerve palsy, unspecified eye
H49.11	Fourth [trochlear] nerve palsy, right eye
H49.12	Fourth [trochlear] nerve palsy, left eye
H49.13	Fourth [trochlear] nerve palsy, bilateral
H49.20	Sixth [abducent] nerve palsy, unspecified eye
H49.21	Sixth [abducent] nerve palsy, right eye
H49.22	Sixth [abducent] nerve palsy, left eye
H49.23	Sixth [abducent] nerve palsy, bilateral
H49.30	Total (external) ophthalmoplegia, unspecified eye
H49.31	Total (external) ophthalmoplegia, right eye
H49.32	Total (external) ophthalmoplegia, left eye
H49.33	Total (external) ophthalmoplegia, bilateral
H49.40	Progressive external ophthalmoplegia, unspecified eye
H49.41	Progressive external ophthalmoplegia, right eye
H49.42	Progressive external ophthalmoplegia, left eye
H49.43	Progressive external ophthalmoplegia, bilateral
H49.881	Other paralytic strabismus, right eye
H49.882	Other paralytic strabismus, left eye
H49.883	Other paralytic strabismus, bilateral
H49.889	Other paralytic strabismus, unspecified eye
H49.9	Unspecified paralytic strabismus
H50.00	Unspecified esotropia
H50.011	Monocular esotropia, right eye
H50.012	Monocular esotropia, left eye
H50.021	Monocular esotropia with A pattern, right eye
H50.022	Monocular esotropia with A pattern, left eye
H50.031	Monocular esotropia with V pattern, right eye
H50.032	Monocular esotropia with V pattern, left eye
H50.041	Monocular esotropia with other noncomitancies, right eye
H50.042	Monocular esotropia with other noncomitancies, left eye
H50.05	Alternating esotropia
H50.06	Alternating esotropia with A pattern
H50.07	Alternating esotropia with V pattern
H50.08	Alternating esotropia with other noncomitancies
H50.10	Unspecified exotropia
H50.111	Monocular exotropia, right eye
H50.112	Monocular exotropia, left eye
H50.121	Monocular exotropia with A pattern, right eye
H50.122	Monocular exotropia with A pattern, left eye
H50.131	Monocular exotropia with V pattern, right eye
H50.132	Monocular exotropia with V pattern, left eye
H50.141	Monocular exotropia with other noncomitancies, right eye
H50.142	Monocular exotropia with other noncomitancies, left eye
H50.15	Alternating exotropia
H50.16	Alternating exotropia with A pattern
H50.17	Alternating exotropia with V pattern

H50.18	Alternating exotropia with other noncomitancies
H50.21	Vertical strabismus, right eye
H50.22	Vertical strabismus, left eye
H50.30	Unspecified intermittent heterotropia
H50.311	Intermittent monocular esotropia, right eye
H50.312	Intermittent monocular esotropia, left eye
H50.32	Intermittent alternating esotropia
H50.331	Intermittent monocular exotropia, right eye
H50.332	Intermittent monocular exotropia, left eye
H50.34	Intermittent alternating exotropia
H50.40	Unspecified heterotropia
H50.411	Cyclotropia, right eye
H50.412	Cyclotropia, left eye
H50.42	Monofixation syndrome
H50.43	Accommodative component in esotropia
H50.50	Unspecified heterophoria
H50.51	Esophoria
H50.52	Exophoria
H50.53	Vertical heterophoria
H50.54	Cyclophoria
H50.55	Alternating heterophoria
H50.60	Mechanical strabismus, unspecified
H50.611	Brown's sheath syndrome, right eye
H50.612	Brown's sheath syndrome, left eye
H50.69	Other mechanical strabismus
H50.811	Duane's syndrome, right eye
H50.812	Duane's syndrome, left eye
H50.89	Other specified strabismus
H50.9	Unspecified strabismus
H51.0	Palsy (spasm) of conjugate gaze
H51.11	Convergence insufficiency
H51.12	Convergence excess
H51.20	Internuclear ophthalmoplegia, unspecified eye
H51.21	Internuclear ophthalmoplegia, right eye
H51.22	Internuclear ophthalmoplegia, left eye
H51.23	Internuclear ophthalmoplegia, bilateral
H51.8	Other specified disorders of binocular movement
H51.9	Unspecified disorder of binocular movement
J38.5	Laryngeal spasm
J38.7	Other diseases of larynx
K22.0	Achalasia of cardia
K59.4	Anal spasm
K60.0	Acute anal fissure
K60.1	Chronic anal fissure
K60.2	Anal fissure, unspecified
M43.6	Torticollis
M62.40	Contracture of muscle, unspecified site
M62.411	Contracture of muscle, right shoulder
M62.412	Contracture of muscle, left shoulder
M62.419	Contracture of muscle, unspecified shoulder
M62.421	Contracture of muscle, right upper arm

M62.422	Contracture of muscle, left upper arm
M62.429	Contracture of muscle, unspecified upper arm
M62.431	Contracture of muscle, right forearm
M62.432	Contracture of muscle, left forearm
M62.439	Contracture of muscle, unspecified forearm
M62.441	Contracture of muscle, right hand
M62.442	Contracture of muscle, left hand
M62.449	Contracture of muscle, unspecified hand
M62.451	Contracture of muscle, right thigh
M62.452	Contracture of muscle, left thigh
M62.459	Contracture of muscle, unspecified thigh
M62.461	Contracture of muscle, right lower leg
M62.462	Contracture of muscle, left lower leg
M62.469	Contracture of muscle, unspecified lower leg
M62.471	Contracture of muscle, right ankle and foot
M62.472	Contracture of muscle, left ankle and foot
M62.479	Contracture of muscle, unspecified ankle and foot
M62.48	Contracture of muscle, other site
M62.49	Contracture of muscle, multiple sites
M62.831	Muscle spasm of calf
M62.838	Other muscle spasm
N31.9	Neuromuscular dysfunction of bladder, unspecified
N32.81	Overactive bladder
N39.3	Stress incontinence (female) (male)
N39.41	Urge incontinence
N39.42	Incontinence without sensory awareness
N39.43	Post-void dribbling
N39.44	Nocturnal enuresis
N39.45	Continuous leakage
N39.46	Mixed incontinence
N39.490	Overflow incontinence
N39.498	Other specified urinary incontinence
Q68.0	Congenital deformity of sternocleidomastoid muscle
R29.898	Other symptoms and signs involving the musculoskeletal system
R32	Unspecified urinary incontinence
R49.8	Other voice and resonance disorders
R51	Headache
S13.4xxA	Sprain of ligaments of cervical spine, initial encounter
S13.4xxD	Sprain of ligaments of cervical spine, subsequent encounter
S13.4xxS	Sprain of ligaments of cervical spine, sequela
S13.8xxA	Sprain of joints and ligaments of other parts of neck, initial encounter
S13.8xxD	Sprain of joints and ligaments of other parts of neck, subsequent encounter
S13.8xxS	Sprain of joints and ligaments of other parts of neck, sequela
S16.1xxA	Strain of muscle, fascia and tendon at neck level, initial encounter
S16.1xxD	Strain of muscle, fascia and tendon at neck level, subsequent encounter
S16.1xxS	Strain of muscle, fascia and tendon at neck level, sequela

The above **medical necessity criteria MUST** be met for the following codes to be covered for Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity:

HCPCS Codes

HCPCS codes:	Code Description
J0586	Injection, abobotulinumtoxinA, 5 units (Dysport)

The following ICD Diagnosis Codes are considered medically necessary when submitted with the HCPCS code above if medical necessity criteria are met:

ICD-10 Diagnosis Codes

ICD-10-CM diagnosis codes:	Code Description
G11.4	Hereditary spastic paraplegia
G24.3	Spasmodic Torticollis
G35	Multiple sclerosis
G80.0	Spastic quadriplegic cerebral palsy
G80.1	Spastic diplegic cerebral palsy
G80.2	Spastic hemiplegic cerebral palsy
G80.3	Athetoid cerebral palsy
G80.4	Ataxic cerebral palsy
G80.8	Other cerebral palsy
G80.9	Cerebral palsy, unspecified
G81.10	Spastic hemiplegia affecting unspecified side
G81.11	Spastic hemiplegia affecting right dominant side
G81.12	Spastic hemiplegia affecting left dominant side
G81.13	Spastic hemiplegia affecting right nondominant side
G81.14	Spastic hemiplegia affecting left nondominant side

The above medical necessity criteria **MUST** be met for the following codes to be covered for Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity:

HCPCS Codes

HCPCS codes:	Code Description
J0588	Injection, incobotulinumtoxinA, 1 unit (Xeomin)

The following ICD Diagnosis Codes are considered medically necessary when submitted with the CPT codes above if medical necessity criteria are met:

ICD-10 Diagnosis Codes:

ICD-10-CM diagnosis codes:	Code Description
G24.3	Spasmodic Torticollis
G24.5	Blepharospasm

Policy History

Date	Action
11/2018	BCBSA National medical policy review. No changes to policy statements. New references added.
11/2018	Updated new FDA indication for chronic sialorrhea.
6/2018	Updated to clarify coverage and to add Specialty Pharmacy link.
1/2018	Updated to add Dysport's updated spasticity FDA indication.

07/2017	Updated to Prefer Dysport & Botox and to include hyperhidrosis to this policy and retired policy 405. Clarified coding information.
11/2015	Clarified coding information.
7/2014	Updated Coding section with ICD10 procedure and diagnosis codes, effective 10/2015.
3/2014	Updated to include adding the sub specialty of board certified headache medicine.
1/2014	Updated to remove Blue Value.
12/2012	Updated to add new CPT code 64615 effective 1/1/2013.
10/2012	Updated to reclassify as a pharmacy medical policy.
11/2011-4/2012	Medical policy ICD 10 remediation: Formatting, editing and coding updates. No changes to policy statements.
7/2012	Updated to clarify coverage criteria and coding for Dysport™ (abobotulinumtoxinA), add diagnosis codes for cervical dystonia, clarify the patient safety section, and add ophthalmologist under migraine criteria.
1/2012	Reviewed - Medical Policy Group - Neurology and Neurosurgery. No changes to policy statements.
11/2011	Reviewed - Medical Policy Group - Plastic Surgery and Dermatology. No changes to policy statements.
5/2011	Updated to include coverage criteria for new FDA approved indication of migraine for Botox
2/2011	Reviewed - Medical Policy Group - Psychiatry and Ophthalmology. No changes to policy statements.
1/2011	Reviewed - Medical Policy Group - Neurology and Neurosurgery. No changes to policy statements.
12/2010	Reviewed - Medical Policy Group - Plastic Surgery and Dermatology. No changes to policy statements.
12/2010	Updated to include coverage criteria for new FDA-approved product Xeomin® (incobotulinumtoxinA).
6/2010	Updated to include coverage criteria for new FDA-approved product Dysport™ (abobotulinumtoxinA).
6/2010	BCBSA National medical policy review. Changes to policy statements.
2/2010	Reviewed - Medical Policy Group - Psychiatry and Ophthalmology. No changes to policy statements.
1/2010	Reviewed - Medical Policy Group - Neurology and Neurosurgery. No changes to policy statements.
1/2010	Updated to include 10/1 UM requirements.
12/2009	Reviewed - Medical Policy Group - Plastic Surgery and Dermatology. No changes to policy statements.
12/2009	Updated to remove coverage of Botulinum Type B, Myobloc™ for all types of hyperhidrosis.
2/2009	Reviewed - Medical Policy Group - Psychiatry and Ophthalmology. No changes to policy statements.
1/2009	Reviewed - Medical Policy Group - Neurology and Neurosurgery. No changes to policy statements.
12/2008	Reviewed - Medical Policy Group - Plastic Surgery and Dermatology. No changes to policy statements.
1/2008	Reviewed - Medical Policy Group - Neurology and Neurosurgery. No changes to policy statements.
12/2007	Reviewed - Medical Policy Group - Plastic Surgery and Dermatology. No changes to policy statements.
1/2007	Reviewed - Medical Policy Group - Neurology and Neurosurgery. No changes to policy statements.
1/2007	BCBSA National medical policy review. Changes to policy statements.

1/1/2001	New policy, effective 1/1/2001, describing covered and non-covered indications.
----------	---

References

#5.01.05

1. Hyman N. Botulinum toxin for focal dystonia. *Pract Neurol* 2004;4(1):30-35.
2. Hsiung GY, Das SK, Ranawaya R, et al. Long-term efficacy of botulinum toxin A in treatment of various movement disorders over a 10-year period. *Mov Disord*. Nov 2002;17(6):1288-1293. PMID 12465070
3. Mejia NI, Vuong KD, Jankovic J. Long-term botulinum toxin efficacy, safety, and immunogenicity. *Mov Disord*. May 2005;20(5):592-597. PMID 15645481
4. Naumann M, Toyka KV, Mansouri Taleghani B, et al. Depletion of neutralising antibodies resensitises a secondary non-responder to botulinum A neurotoxin. *J Neurol Neurosurg Psychiatry*. Dec 1998;65(6):924-927. PMID 9854974
5. Mahant N, Clouston PD, Lorentz IT. The current use of botulinum toxin. *J Clin Neurosci*. Sep 2000;7(5):389-394. PMID 10942658
6. Herrmann J, Geth K, Mall V, et al. Clinical impact of antibody formation to botulinum toxin A in children. *Ann Neurol*. May 2004;55(5):732-735. PMID 15122715
7. Koman LA, Brashers A, Rosenfeld S, et al. Botulinum toxin type a neuromuscular blockade in the treatment of equinus foot deformity in cerebral palsy: a multicenter, open-label clinical trial. *Pediatrics*. Nov 2001;108(5):1062-1071. PMID 11694682
8. Sankhla C, Jankovic J, Duane D. Variability of the immunologic and clinical response in dystonic patients immunoresistant to botulinum toxin injections. *Mov Disord*. Jan 1998;13(1):150-154. PMID 9452341
9. Dutton JJ, White JJ, Richard MJ. Myobloc for the treatment of benign essential blepharospasm in patients refractory to botox. *Ophthal Plast Reconstr Surg*. May-Jun 2006;22(3):173-177. PMID 16714924
10. Pearce LB, Borodic GE, First ER, et al. Measurement of botulinum toxin activity: evaluation of the lethality assay. *Toxicol Appl Pharmacol*. Sep 1994;128(1):69-77. PMID 8079356
11. Goschel H, Wohlfarth K, Frevert J, et al. Botulinum A toxin therapy: neutralizing and nonneutralizing antibodies--therapeutic consequences. *Exp Neurol*. Sep 1997;147(1):96-102. PMID 9294406
12. Cordivari C, Misra VP, Vincent A, et al. Secondary nonresponsiveness to botulinum toxin A in cervical dystonia: the role of electromyogram-guided injections, botulinum toxin A antibody assay, and the extensor digitorum brevis test. *Mov Disord*. Oct 2006;21(10):1737-1741. PMID 16874756
13. Hanna PA, Jankovic J. Mouse bioassay versus Western blot assay for botulinum toxin antibodies: correlation with clinical response. *Neurology*. Jun 1998;50(6):1624-1629. PMID 9633703
14. Food and Drug Administration (FDA). Information for Healthcare Professionals: OnabotulinumtoxinA (marketed as Botox/Botox Cosmetic), AbobotulinumtoxinA (marketed as Dysport) and RimabotulinumtoxinB (marketed as Myobloc). 2009; <https://wayback.archive-it.org/7993/20170406044653/https://www.fda.gov/Drugs/DrugSafety/PostmarketDrugSafetyInformationforPatientsandProviders/DrugSafetyInformationforHealthcareProfessionals/ucm174949.htm>. Accessed October 1, 2018.
15. Castela M, Marques RE, Duarte GS, et al. Botulinum toxin type A therapy for cervical dystonia. *Cochrane Database Syst Rev*. Dec 12 2017;12:CD003633. PMID 29230798
16. Dong Y, Wu T, Hu X, et al. Efficacy and safety of botulinum toxin type A for upper limb spasticity after stroke or traumatic brain injury: a systematic review with meta-analysis and trial sequential analysis. *Eur J Phys Rehabil Med*. Apr 2017;53(2):256-267. PMID 27834471
17. Baker JA, Pereira G. The efficacy of Botulinum Toxin A for limb spasticity on improving activity restriction and quality of life: a systematic review and meta-analysis using the GRADE approach. *Clin Rehabil*. Jun 2016;30(6):549-558. PMID 26150020
18. Marques RE, Duarte GS, Rodrigues FB, et al. Botulinum toxin type B for cervical dystonia. *Cochrane Database Syst Rev*. May 13 2016(5):CD004315. PMID 27176573
19. Duarte GS, Castela M, Rodrigues FB, et al. Botulinum toxin type A versus botulinum toxin type B for cervical dystonia. *Cochrane Database Syst Rev*. Oct 26 2016;10:CD004314. PMID 27782297

20. Dashtipour K, Chen JJ, Walker HW, et al. Systematic literature review of abobotulinumtoxinA in clinical trials for adult upper limb spasticity. *Am J Phys Med Rehabil.* Mar 2015;94(3):229-238. PMID 25299523
21. Marsh WA, Monroe DM, Brin MF, et al. Systematic review and meta-analysis of the duration of clinical effect of onabotulinumtoxinA in cervical dystonia. *BMC Neurol.* Apr 27 2014;14:91. PMID 24767576
22. Foley N, Pereira S, Salter K, et al. Treatment with botulinum toxin improves upper-extremity function post stroke: a systematic review and meta-analysis. *Arch Phys Med Rehabil.* May 2013;94(5):977-989. PMID 23262381
23. Baker JA, Pereira G. The efficacy of botulinum toxin A for spasticity and pain in adults: a systematic review and meta-analysis using the Grades of Recommendation, Assessment, Development and Evaluation approach. *Clin Rehabil.* Dec 2013;27(12):1084-1096. PMID 23864518
24. Blue Cross Blue Shield Association Technology Evaluation Center (TEC). Botulinum-A toxin for treatment of chronic spasticity. *TEC Assessments* 1996;Volume 11, Tab 6.
25. Blue Cross Blue Shield Association Technology Evaluation Center (TEC). Botulinum toxin for treatment of primary chronic headache disorders. *TEC Assessments.* 2004;Volume 19, Tab 10.
26. Naumann M, So Y, Argoff CE, et al. Assessment: Botulinum neurotoxin in the treatment of autonomic disorders and pain (an evidence-based review): report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology.* May 6 2008;70(19):1707-1714. PMID 18458231
27. Simpson DM, Gracies JM, Graham HK, et al. Assessment: Botulinum neurotoxin for the treatment of spasticity (an evidence-based review): report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology.* May 06 2008;70(19):1691-1698. PMID 18458229
28. Wein T, Esquenazi A, Jost WH, et al. OnabotulinumtoxinA for the treatment of poststroke distal lower limb spasticity: a randomized trial. *PM R.* Jul 2018;10(7):693-703. PMID 29330071
29. Wissel J, Ganapathy V, Ward AB, et al. OnabotulinumtoxinA improves pain in patients with post-stroke spasticity: findings from a randomized, double-blind, placebo-controlled trial. *J Pain Symptom Manage.* Jul 2016;52(1):17-26. PMID 27037050
30. Gracies JM, Brashear A, Jech R, et al. Safety and efficacy of abobotulinumtoxinA for hemiparesis in adults with upper limb spasticity after stroke or traumatic brain injury: a double-blind randomised controlled trial. *Lancet Neurol.* Oct 2015;14(10):992-1001. PMID 26318836
31. Shaw LC, Price CI, van Wijck FM, et al. Botulinum Toxin for the Upper Limb after Stroke (BoTULS) Trial: effect on impairment, activity limitation, and pain. *Stroke.* May 2011;42(5):1371-1379. PMID 21415398
32. Kanovsky P, Slawek J, Denes Z, et al. Efficacy and safety of botulinum neurotoxin NT 201 in poststroke upper limb spasticity. *Clin Neuropharmacol.* Sep-Oct 2009;32(5):259-265. PMID 19644361
33. Kanovsky P, Slawek J, Denes Z, et al. Efficacy and safety of treatment with incobotulinum toxin A (botulinum neurotoxin type A free from complexing proteins; NT 201) in post-stroke upper limb spasticity. *J Rehabil Med.* May 2011;43(6):486-492. PMID 21533328
34. Delgado MR, Tilton A, Russman B, et al. AbobotulinumtoxinA for equinus foot deformity in cerebral palsy: a randomized controlled trial. *Pediatrics.* Feb 2016;137(2):e20152830. PMID 26812925
35. Rowe FJ, Noonan CP. Botulinum toxin for the treatment of strabismus. *Cochrane Database Syst Rev.* Feb 15 2012;2(2):CD006499. PMID 22336817
36. Tejedor J, Rodriguez JM. Early retreatment of infantile esotropia: comparison of reoperation and botulinum toxin. *Br J Ophthalmol.* Jul 1999;83(7):783-787. PMID 10381663
37. Lee J, Harris S, Cohen J, et al. Results of a prospective randomized trial of botulinum toxin therapy in acute unilateral sixth nerve palsy. *J Pediatr Ophthalmol Strabismus.* Sep-Oct 1994;31(5):283-286. PMID 7837013
38. Dashtipour K, Chen JJ, Frei K, et al. Systematic literature review of AbobotulinumtoxinA in clinical trials for blepharospasm and hemifacial spasm. *Tremor Other Hyperkinet Mov (N Y).* Nov 2015;5:338. PMID 26566457
39. Jankovic J. Clinical efficacy and tolerability of Xeomin in the treatment of blepharospasm. *Eur J Neurol.* Dec 2009;16(Suppl 2):14-18. PMID 20002742

40. Nussgens Z, Roggenkamper P. Comparison of two botulinum-toxin preparations in the treatment of essential blepharospasm. *Graefes Arch Clin Exp Ophthalmol*. Apr 1997;235(4):197-199. PMID 9143885
41. Roggenkamper P, Jost WH, Bihari K, et al. Efficacy and safety of a new botulinum toxin type A free of complexing proteins in the treatment of blepharospasm. *J Neural Transm*. Mar 2006;113(3):303-312. PMID 15959841
42. Headache Classification Committee of the International Headache Society. The International Classification of Headache Disorders, 3rd edition (beta version). *Cephalalgia*. Jul 2013;33(9):629-808. PMID 23771276
43. Shamliyan TA, Kane RL, Taylor FR. *Migraine in Adults: Preventive Pharmacologic Treatments (Comparative Effectiveness Review No. 103)*. Rockville MD: Agency for Healthcare Research and Quality; 2013.
44. Jackson JL, Kuriyama A, Hayashino Y. Botulinum toxin A for prophylactic treatment of migraine and tension headaches in adults: a meta-analysis. *JAMA*. Apr 25 2012;307(16):1736-1745. PMID 22535858
45. Shuhendler AJ, Lee S, Siu M, et al. Efficacy of botulinum toxin type A for the prophylaxis of episodic migraine headaches: a meta-analysis of randomized, double-blind, placebo-controlled trials. *Pharmacotherapy*. Jul 2009;29(7):784-791. PMID 19558252
46. Aurora SK, Dodick DW, Turkel CC, et al. OnabotulinumtoxinA for treatment of chronic migraine: results from the double-blind, randomized, placebo-controlled phase of the PREEMPT 1 trial. *Cephalalgia*. Jul 2010;30(7):793-803. PMID 20647170
47. Diener HC, Dodick DW, Aurora SK, et al. OnabotulinumtoxinA for treatment of chronic migraine: results from the double-blind, randomized, placebo-controlled phase of the PREEMPT 2 trial. *Cephalalgia*. Jul 2010;30(7):804-814. PMID 20647171
48. Dodick DW, Turkel CC, DeGryse RE, et al. OnabotulinumtoxinA for treatment of chronic migraine: pooled results from the double-blind, randomized, placebo-controlled phases of the PREEMPT clinical program. *Headache*. Jun 2010;50(6):921-936. PMID 20487038
49. Aurora SK, Winner P, Freeman MC, et al. OnabotulinumtoxinA for treatment of chronic migraine: pooled analyses of the 56-week PREEMPT clinical program. *Headache*. Oct 2011;51(9):1358-1373. PMID 21883197
50. Matharu M, Halker R, Pozo-Rosich P, et al. The impact of onabotulinumtoxinA on severe headache days: PREEMPT 56-week pooled analysis. *J Headache Pain*. Dec 2017;18(1):78. PMID 28766236
51. Cady RK, Schreiber CP, Porter JA, et al. A multi-center double-blind pilot comparison of onabotulinumtoxinA and topiramate for the prophylactic treatment of chronic migraine. *Headache*. Jan 2011;51(1):21-32. PMID 21070228
52. Silberstein SD, Olesen J, Bousser MG, et al. The International Classification of Headache Disorders, 2nd Edition (ICHD-II)--revision of criteria for 8.2 Medication-overuse headache. *Cephalalgia*. Jun 2005;25(6):460-465. PMID 15910572
53. Silberstein SD, Blumenfeld AM, Cady RK, et al. OnabotulinumtoxinA for treatment of chronic migraine: PREEMPT 24-week pooled subgroup analysis of patients who had acute headache medication overuse at baseline. *J Neurol Sci*. Aug 15 2013;331(1-2):48-56. PMID 23790235
54. Silberstein SD, Gobel H, Jensen R, et al. Botulinum toxin type A in the prophylactic treatment of chronic tension-type headache: a multicentre, double-blind, randomized, placebo-controlled, parallel-group study. *Cephalalgia*. Jul 2006;26(7):790-800. PMID 16776693
55. Ondo WG, Vuong KD, Derman HS. Botulinum toxin A for chronic daily headache: a randomized, placebo-controlled, parallel design study. *Cephalalgia*. Jan 2004;24(1):60-65. PMID 14687015
56. Mathew NT, Frishberg BM, Gawel M, et al. Botulinum toxin type A (BOTOX) for the prophylactic treatment of chronic daily headache: a randomized, double-blind, placebo-controlled trial. *Headache*. Apr 2005;45(4):293-307. PMID 15836565
57. Silberstein SD, Stark SR, Lucas SM, et al. Botulinum toxin type A for the prophylactic treatment of chronic daily headache: a randomized, double-blind, placebo-controlled trial. *Mayo Clin Proc*. Sep 2005;80(9):1126-1137. PMID 16178492
58. Peloso P, Gross A, Haines T, et al. Medicinal and injection therapies for mechanical neck disorders. *Cochrane Database Syst Rev*. Jul 18 2007(3):CD000319. PMID 17636629

59. Linde M, Hagen K, Salvesen O, et al. Onabotulinum toxin A treatment of cervicogenic headache: a randomised, double-blind, placebo-controlled crossover study. *Cephalalgia*. May 2011;31(7):797-807. PMID 21300635
60. Braker C, Yariv S, Adler R, et al. The analgesic effect of botulinum-toxin A on postwhiplash neck pain. *Clin J Pain*. Jan 2008;24(1):5-10. PMID 18180629
61. Freund BJ, Schwartz M. Treatment of chronic cervical-associated headache with botulinum toxin A: a pilot study. *Headache*. Mar 2000;40(3):231-236. PMID 10759926
62. Padberg M, de Bruijn SF, Tavy DL. Neck pain in chronic whiplash syndrome treated with botulinum toxin. A double-blind, placebo-controlled clinical trial. *J Neurol*. Mar 2007;254(3):290-295. PMID 17345052
63. Leyden JE, Moss AC, MacMathuna P. Endoscopic pneumatic dilation versus botulinum toxin injection in the management of primary achalasia. *Cochrane Database Syst Rev*. Dec 2014;12(12):CD005046. PMID 25485740
64. Lagalla G, Millevolte M, Capecchi M, et al. Botulinum toxin type A for drooling in Parkinson's disease: a double-blind, randomized, placebo-controlled study. *Mov Disord*. May 2006;21(5):704-707. PMID 16440332
65. Ondo WG, Hunter C, Moore W. A double-blind placebo-controlled trial of botulinum toxin B for sialorrhea in Parkinson's disease. *Neurology*. Jan 13 2004;62(1):37-40. PMID 14718694
66. Mancini F, Zangaglia R, Cristina S, et al. Double-blind, placebo-controlled study to evaluate the efficacy and safety of botulinum toxin type A in the treatment of drooling in parkinsonism. *Mov Disord*. Jun 2003;18(6):685-688. PMID 12784273
67. Squires N, Humberstone M, Wills A, et al. The use of botulinum toxin injections to manage drooling in amyotrophic lateral sclerosis/motor neurone disease: a systematic review. *Dysphagia*. Aug 2014;29(4):500-508. PMID 24847964
68. Rodwell K, Edwards P, Ware RS, et al. Salivary gland botulinum toxin injections for drooling in children with cerebral palsy and neurodevelopmental disability: a systematic review. *Dev Med Child Neurol*. Nov 2012;54(11):977-987. PMID 22946706
69. Gonzalez LM, Martinez C, Bori YFI, et al. Factors in the efficacy, safety, and impact on quality of life for treatment of drooling with botulinum toxin type a in patients with cerebral palsy. *Am J Phys Med Rehabil*. Feb 2017;96(2):68-76. PMID 28099276
70. Reid SM, Johnstone BR, Westbury C, et al. Randomized trial of botulinum toxin injections into the salivary glands to reduce drooling in children with neurological disorders. *Dev Med Child Neurol*. Feb 2008;50(2):123-128. PMID 18201301
71. Chan KH, Liang C, Wilson P, et al. Long-term safety and efficacy data on botulinum toxin type A: an injection for sialorrhea. *JAMA Otolaryngol Head Neck Surg*. Feb 2013;139(2):134-138. PMID 23429943
72. Friedmacher F, Puri P. Comparison of posterior internal anal sphincter myectomy and intrasphincteric botulinum toxin injection for treatment of internal anal sphincter achalasia: a meta-analysis. *Pediatr Surg Int*. Aug 2012;28(8):765-771. PMID 22806601
73. Emile SH, Elfeki HA, Elbanna HG, et al. Efficacy and safety of botulinum toxin in treatment of anismus: A systematic review. *World J Gastrointest Pharmacol Ther*. Aug 06 2016;7(3):453-462. PMID 27602248
74. Farid M, El Monem HA, Omar W, et al. Comparative study between biofeedback retraining and botulinum neurotoxin in the treatment of anismus patients. *Int J Colorectal Dis*. Jan 2009;24(1):115-120. PMID 18719924
75. Farid M, Youssef T, Mahdy T, et al. Comparative study between botulinum toxin injection and partial division of puborectalis for treating anismus. *Int J Colorectal Dis*. Mar 2009;24(3):327-334. PMID 19039596
76. Yiannakopoulou E. Botulinum toxin and anal fissure: efficacy and safety systematic review. *Int J Colorectal Dis*. Jan 2012;27(1):1-9. PMID 21822595
77. Brisinda G, Cadeddu F, Brandara F, et al. Randomized clinical trial comparing botulinum toxin injections with 0.2 per cent nitroglycerin ointment for chronic anal fissure. *Br J Surg*. Feb 2007;94(2):162-167. PMID 17256809
78. Brisinda G, Albanese A, Cadeddu F, et al. Botulinum neurotoxin to treat chronic anal fissure: results of a randomized "Botox vs. Dysport" controlled trial. *Aliment Pharmacol Ther*. Mar 15 2004;19(6):695-701. PMID 15023172

79. Brisinda G, Maria G, Bentivoglio AR, et al. A comparison of injections of botulinum toxin and topical nitroglycerin ointment for the treatment of chronic anal fissure. *N Engl J Med*. Jul 8 1999;341(2):65-69. PMID 10395629
80. Maria G, Cassetta E, Gui D, et al. A comparison of botulinum toxin and saline for the treatment of chronic anal fissure. *N Engl J Med*. Jan 22 1998;338(4):217-220. PMID 9435326
81. De Nardi P, Ortolano E, Radaelli G, et al. Comparison of glycerine trinitrate and botulinum toxin-a for the treatment of chronic anal fissure: long-term results. *Dis Colon Rectum*. Apr 2006;49(4):427-432. PMID 16456637
82. Fruehauf H, Fried M, Wegmueller B, et al. Efficacy and safety of botulinum toxin a injection compared with topical nitroglycerin ointment for the treatment of chronic anal fissure: a prospective randomized study. *Am J Gastroenterol*. Sep 2006;101(9):2107-2112. PMID 16848808
83. Iswariah H, Stephens J, Rieger N, et al. Randomized prospective controlled trial of lateral internal sphincterotomy versus injection of botulinum toxin for the treatment of idiopathic fissure in ano. *ANZ J Surg*. Jul 2005;75(7):553-555. PMID 15972045
84. Nasr M, Ezzat H, Elsebae M. Botulinum toxin injection versus lateral internal sphincterotomy in the treatment of chronic anal fissure: a randomized controlled trial. *World J Surg*. Nov 2010;34(11):2730-2734. PMID 20703472
85. Chen HL, Woo XB, Wang HS, et al. Botulinum toxin injection versus lateral internal sphincterotomy for chronic anal fissure: a meta-analysis of randomized control trials. *Tech Coloproctol*. Aug 2014;18(8):693-698. PMID 24500725
86. Drake MJ, Nitti VW, Ginsberg DA, et al. Comparative assessment of the efficacy of onabotulinumtoxinA and oral therapies (anticholinergics and mirabegron) for overactive bladder: a systematic review and network meta-analysis. *BJU Int*. Nov 2017;120(5):611-622. PMID 28670786
87. Freemantle N, Ginsberg DA, McCool R, et al. Comparative assessment of onabotulinumtoxinA and mirabegron for overactive bladder: an indirect treatment comparison. *BMJ Open*. Feb 23 2016;6(2):e009122. PMID 26908514
88. Cheng T, Shuang WB, Jia DD, et al. Efficacy and safety of onabotulinumtoxinA in patients with neurogenic detrusor overactivity: a systematic review and meta-analysis of randomized controlled trials. *PLoS One*. Jul 2016;11(7):e0159307. PMID 27463810
89. Cui Y, Zhou X, Zong H, et al. The efficacy and safety of onabotulinumtoxinA in treating idiopathic OAB: A systematic review and meta-analysis. *NeuroUrol Urodyn*. Jun 2015;34(5):413-419. PMID 24676791
90. Cui Y, Wang L, Liu L, et al. Botulinum toxin-A injections for idiopathic overactive bladder: a systematic review and meta-analysis. *Urol Int*. Aug 2013;91(4):429-438. PMID 23970316
91. Duthie JB, Vincent M, Herbison GP, et al. Botulinum toxin injections for adults with overactive bladder syndrome. *Cochrane Database Syst Rev*. Dec 07 2011(12):CD005493. PMID 22161392
92. Soljanik I. Efficacy and safety of botulinum toxin a intradetrusor injections in adults with neurogenic detrusor overactivity/neurogenic overactive bladder: a systematic review. *Drugs*. Jul 2013;73(10):1055-1066. PMID 23775527
93. Mehta S, Hill D, McIntyre A, et al. Meta-analysis of botulinum toxin A detrusor injections in the treatment of neurogenic detrusor overactivity after spinal cord injury. *Arch Phys Med Rehabil*. Aug 2013;94(8):1473-1481. PMID 23632286
94. Karsenty G, Denys P, Amarenco G, et al. Botulinum toxin A (Botox) intradetrusor injections in adults with neurogenic detrusor overactivity/neurogenic overactive bladder: a systematic literature review. *Eur Urol*. Feb 2008;53(2):275-287. PMID 17988791
95. Herschorn S, Kohan A, Aliotta P, et al. The efficacy and safety of onabotulinumtoxinA or solifenacin compared with placebo in solifenacin naive patients with refractory overactive bladder: results from a multicenter, randomized, double-blind phase 3b trial. *J Urol*. Jul 2017;198(1):167-175. PMID 28161352
96. Nitti VW, Dmochowski R, Herschorn S, et al. OnabotulinumtoxinA for the treatment of patients with overactive bladder and urinary incontinence: results of a phase 3, randomized, placebo controlled trial. *J Urol*. Feb 2017;197(2S):S216-S223. PMID 28012773
97. Amundsen CL, Richter HE, Menefee SA, et al. OnabotulinumtoxinA vs sacral neuromodulation on refractory urgency urinary incontinence in women: a randomized clinical trial. *JAMA*. Oct 04 2016;316(13):1366-1374. PMID 27701661

98. Amundsen CL, Komesu YM, Chermansky C, et al. Two-year outcomes of sacral neuromodulation versus onabotulinumtoxinA for refractory urgency urinary incontinence: a randomized trial. *Eur Urol.* Jul 2018;74(1):66-73. PMID 29482936
99. Nitti VW, Dmochowski R, Herschorn S, et al. OnabotulinumtoxinA for the treatment of patients with overactive bladder and urinary incontinence: results of a phase 3, randomized, placebo controlled trial. *J Urol.* Jun 2013;189(6):2186-2193. PMID 23246476
100. Chapple C, Sievert KD, Macdiarmid S, et al. OnabotulinumtoxinA 100 U significantly improves all idiopathic overactive bladder symptoms and quality of life in patients with overactive bladder and urinary incontinence: a randomised, double-blind, placebo-controlled trial. *Eur Urol.* Aug 2013;64(2):249-256. PMID 23608668
101. Sievert KD, Chapple C, Herschorn S, et al. OnabotulinumtoxinA 100U provides significant improvements in overactive bladder symptoms in patients with urinary incontinence regardless of the number of anticholinergic therapies used or reason for inadequate management of overactive bladder. *Int J Clin Pract.* Oct 2014;68(10):1246-1256. PMID 24754838
102. Nitti VW, Ginsberg D, Sievert KD, et al. Durable efficacy and safety of long-term onabotulinumtoxinA treatment in patients with overactive bladder syndrome: final results of a 3.5-year study. *J Urol.* Sep 2016;196(3):791-800. PMID 27038769
103. Ginsberg D, Gousse A, Keppenne V, et al. Phase 3 efficacy and tolerability study of onabotulinumtoxinA for urinary incontinence from neurogenic detrusor overactivity. *J Urol.* Jun 2012;187(6):2131-2139. PMID 22503020
104. Mehta S, Hill D, Foley N, et al. A meta-analysis of botulinum toxin sphincteric injections in the treatment of incomplete voiding after spinal cord injury. *Arch Phys Med Rehabil.* Apr 2012;93(4):597-603. PMID 22365478
105. Karsenty G, Baazeem A, Elzayat E, et al. Injection of botulinum toxin type A in the urethral sphincter to treat lower urinary tract dysfunction: a review of indications, techniques and results. *Can J Urol.* Apr 2006;13(2):3027-3033. PMID 16672114
106. de Seze M, Petit H, Gallien P, et al. Botulinum toxin and detrusor sphincter dyssynergia: a double-blind lidocaine-controlled study in 13 patients with spinal cord disease. *Eur Urol.* Jul 2002;42(1):56-62. PMID 12121731
107. Marchal C, Perez JE, Herrera B, et al. The use of botulinum toxin in benign prostatic hyperplasia. *Neurourol Urodyn.* Jan 2012;31(1):86-92. PMID 21905088
108. Maria G, Brisinda G, Civello IM, et al. Relief by botulinum toxin of voiding dysfunction due to benign prostatic hyperplasia: results of a randomized, placebo-controlled study. *Urology.* Aug 2003;62(2):259-264; discussion 264-255. PMID 12893330
109. Wang J, Wang Q, Wu Q, et al. Intravesical botulinum toxin A injections for bladder pain syndrome/interstitial cystitis: a systematic review and meta-analysis of controlled studies. *Med Sci Monit.* Sep 14 2016;22:3257-3267. PMID 27624897
110. Tirumuru S, Al-Kurdi D, Latthe P. Intravesical botulinum toxin A injections in the treatment of painful bladder syndrome/interstitial cystitis: a systematic review. *Int Urogynecol J.* Oct 2010;21(10):1285-1300. PMID 20449567
111. Akiyama Y, Nomiya A, Niimi A, et al. Botulinum toxin type A injection for refractory interstitial cystitis: A randomized comparative study and predictors of treatment response. *Int J Urol.* Sep 2015;22(9):835-841. PMID 26041274
112. Kuo HC, Jiang YH, Tsai YC, et al. Intravesical botulinum toxin-A injections reduce bladder pain of interstitial cystitis/bladder pain syndrome refractory to conventional treatment - A prospective, multicenter, randomized, double-blind, placebo-controlled clinical trial. *Neurourol Urodyn.* Jun 2016;35(5):609-614. PMID 25914337
113. Manning J, Dwyer P, Rosamilia A, et al. A multicentre, prospective, randomised, double-blind study to measure the treatment effectiveness of abobotulinum A (AboBTXA) among women with refractory interstitial cystitis/bladder pain syndrome. *Int Urogynecol J.* May 2014;25(5):593-599. PMID 24276074
114. Brin MF, Lyons KE, Doucette J, et al. A randomized, double masked, controlled trial of botulinum toxin type A in essential hand tremor. *Neurology.* Jun 12 2001;56(11):1523-1528. PMID 11402109
115. Jankovic J, Schwartz K, Clemence W, et al. A randomized, double-blind, placebo-controlled study to evaluate botulinum toxin type A in essential hand tremor. *Mov Disord.* May 1996;11(3):250-256. PMID 8723140

116. Mittal SO, Machado D, Richardson D, et al. Botulinum toxin in Parkinson disease tremor: a randomized, double-blind, placebo-controlled study with a customized injection approach. *Mayo Clin Proc.* Sep 2017;92(9):1359-1367. PMID 28789780
117. Foster L, Clapp L, Erickson M, et al. Botulinum toxin A and chronic low back pain: a randomized, double-blind study. *Neurology.* May 22 2001;56(10):1290-1293. PMID 11376175
118. Lin YC, Wu WT, Hsu YC, et al. Comparative effectiveness of botulinum toxin versus non-surgical treatments for treating lateral epicondylitis: a systematic review and meta-analysis. *Clin Rehabil.* Mar 01 2017;269215517702517. PMID 28349703
119. Krogh TP, Bartels EM, Ellingsen T, et al. Comparative effectiveness of injection therapies in lateral epicondylitis: a systematic review and network meta-analysis of randomized controlled trials. *Am J Sports Med.* Jun 2013;41(6):1435-1446. PMID 22972856
120. Sims SE, Miller K, Elfar JC, et al. Non-surgical treatment of lateral epicondylitis: a systematic review of randomized controlled trials. *Hand (N Y).* Dec 2014;9(4):419-446. PMID 25414603
121. Mahowald ML, Singh JA, Dykstra D. Long term effects of intra-articular botulinum toxin A for refractory joint pain. *Neurotox Res.* Apr 2006;9(2-3):179-188. PMID 16785116
122. Singer BJ, Silbert PL, Dunne JW, et al. An open label pilot investigation of the efficacy of Botulinum toxin type A [Dysport] injection in the rehabilitation of chronic anterior knee pain. *Disabil Rehabil.* Jun 15 2006;28(11):707-713. PMID 16809213
123. Soares A, Andriolo RB, Atallah AN, et al. Botulinum toxin for myofascial pain syndromes in adults. *Cochrane Database Syst Rev.* Jul 25 2014;7(7):CD007533. PMID 25062018
124. Desai MJ, Shkolnikova T, Nava A, et al. A critical appraisal of the evidence for botulinum toxin type A in the treatment for cervico-thoracic myofascial pain syndrome. *Pain Pract.* Feb 2014;14(2):185-195. PMID 23692187
125. Langevin P, Lowcock J, Weber J, et al. Botulinum toxin intramuscular injections for neck pain: a systematic review and metaanalysis. *J Rheumatol.* Feb 2011;38(2):203-214. PMID 21123322
126. Nicol AL, Wu, II, Ferrante FM. Botulinum toxin type a injections for cervical and shoulder girdle myofascial pain using an enriched protocol design. *Anesth Analg.* Jun 2014;118(6):1326-1335. PMID 24842179
127. Chen YW, Chiu YW, Chen CY, et al. Botulinum toxin therapy for temporomandibular joint disorders: a systematic review of randomized controlled trials. *Int J Oral Maxillofac Surg.* Aug 2015;44(8):1018-1026. PMID 25920597
128. Morra ME, Elgebaly A, Elmaraezy A, et al. Therapeutic efficacy and safety of Botulinum Toxin A Therapy in Trigeminal Neuralgia: a systematic review and meta-analysis of randomized controlled trials. *J Headache Pain.* Dec 2016;17(1):63. PMID 27377706
129. Zhang H, Lian Y, Ma Y, et al. Two doses of botulinum toxin type A for the treatment of trigeminal neuralgia: observation of therapeutic effect from a randomized, double-blind, placebo-controlled trial. *J Headache Pain.* Sep 27 2014;15:65. PMID 25263254
130. Shehata HS, El-Tamawy MS, Shalaby NM, et al. Botulinum toxin-type A: could it be an effective treatment option in intractable trigeminal neuralgia? *J Headache Pain.* Nov 19 2013;14:92. PMID 24251833
131. Wu CJ, Lian YJ, Zheng YK, et al. Botulinum toxin type A for the treatment of trigeminal neuralgia: results from a randomized, double-blind, placebo-controlled trial. *Cephalalgia.* Apr 2012;32(6):443-450. PMID 22492424
132. Patti R, Almasio PL, Muggeo VM, et al. Improvement of wound healing after hemorrhoidectomy: a double-blind, randomized study of botulinum toxin injection. *Dis Colon Rectum.* Dec 2005;48(12):2173-2179. PMID 16400513
133. Patti R, Almasio PL, Arcara M, et al. Botulinum toxin vs. topical glyceryl trinitrate ointment for pain control in patients undergoing hemorrhoidectomy: a randomized trial. *Dis Colon Rectum.* Nov 2006;49(11):1741-1748. PMID 16990976
134. Ziade M, Domergue S, Batifol D, et al. Use of botulinum toxin type A to improve treatment of facial wounds: a prospective randomised study. *J Plast Reconstr Aesthet Surg.* Feb 2013;66(2):209-214. PMID 23102873
135. Gassner HG, Brissett AE, Otley CC, et al. Botulinum toxin to improve facial wound healing: A prospective, blinded, placebo-controlled study. *Mayo Clin Proc.* Aug 2006;81(8):1023-1028. PMID 16901024

136. Abbott JA, Jarvis SK, Lyons SD, et al. Botulinum toxin type A for chronic pain and pelvic floor spasm in women: a randomized controlled trial. *Obstet Gynecol.* Oct 2006;108(4):915-923. PMID 17012454
137. Dykstra DD, Presthus J. Botulinum toxin type A for the treatment of provoked vestibulodynia: an open-label, pilot study. *J Reprod Med.* Jun 2006;51(6):467-470. PMID 16846084
138. Jarvis SK, Abbott JA, Lenart MB, et al. Pilot study of botulinum toxin type A in the treatment of chronic pelvic pain associated with spasm of the levator ani muscles. *Aust N Z J Obstet Gynaecol.* Feb 2004;44(1):46-50. PMID 15089868
139. Vasan CW, Liu WC, Klusmann JP, et al. Botulinum toxin type A for the treatment of chronic neck pain after neck dissection. *Head Neck.* Jan 2004;26(1):39-45. PMID 14724905
140. Wittekindt C, Liu WC, Preuss SF, et al. Botulinum toxin A for neuropathic pain after neck dissection: a dose-finding study. *Laryngoscope.* Jul 2006;116(7):1168-1171. PMID 16826054
141. Slengerik-Hansen J, Ovesen T. Botulinum toxin treatment of objective tinnitus because of essential palatal tremor: a systematic review. *Otol Neurotol.* Aug 2016;37(7):820-828. PMID 27273401
142. Stidham KR, Solomon PH, Roberson JB. Evaluation of botulinum toxin A in treatment of tinnitus. *Otolaryngol Head Neck Surg.* Jun 2005;132(6):883-889. PMID 15944559
143. Winocour S, Murad MH, Bidgoli-Moghaddam M, et al. A systematic review of the use of Botulinum toxin type A with subpectoral breast implants. *J Plast Reconstr Aesthet Surg.* Jan 2014;67(1):34-41. PMID 24094619
144. Koivusalo AI, Pakarinen MP, Rintala RJ. Botox injection treatment for anal outlet obstruction in patients with internal anal sphincter achalasia and Hirschsprung's disease. *Pediatr Surg Int.* Oct 2009;25(10):873-876. PMID 19662428
145. Minkes RK, Langer JC. A prospective study of botulinum toxin for internal anal sphincter hypertonicity in children with Hirschsprung's disease. *J Pediatr Surg.* Dec 2000;35(12):1733-1736. PMID 11101725
146. Patrus B, Nasr A, Langer JC, et al. Intrasphincteric botulinum toxin decreases the rate of hospitalization for postoperative obstructive symptoms in children with Hirschsprung disease. *J Pediatr Surg.* Jan 2011;46(1):184-187. PMID 21238663
147. Bai Y, Xu MJ, Yang X, et al. A systematic review on intrapyloric botulinum toxin injection for gastroparesis. *Digestion.* Dec 2010;81(1):27-34. PMID 20029206
148. Arts J, Holvoet L, Caenepeel P, et al. Clinical trial: a randomized-controlled crossover study of intrapyloric injection of botulinum toxin in gastroparesis. *Aliment Pharmacol Ther.* Nov 1 2007;26(9):1251-1258. PMID 17944739
149. FriedenberG FK, Palit A, Parkman HP, et al. Botulinum toxin A for the treatment of delayed gastric emptying. *Am J Gastroenterol.* Feb 2008;103(2):416-423. PMID 18070232
150. Magid M, Finzi E, Kruger TH, et al. Treating depression with botulinum toxin: a pooled analysis of randomized controlled trials. *Pharmacopsychiatry.* Sep 2015;48(6):205-210. PMID 26252721
151. American Urological Association (AUA). Diagnosis and treatment of non-neurogenic overactive bladder (OAB) in adults: AUA/SUFU guideline. 2012, amended in 2014; [https://www.auanet.org/guidelines/incontinence-non-neurogenic-overactive-bladder-\(2012-amended-2014\)](https://www.auanet.org/guidelines/incontinence-non-neurogenic-overactive-bladder-(2012-amended-2014)). Accessed October 1, 2018.
152. American Urological Association (AUA), Hanno PH, Burks DA, et al. Diagnosis and treatment of interstitial cystitis/bladder pain syndrome. 2011; amended 2014; <https://www.auanet.org/education/guidelines/ic-bladder-pain-syndrome.cfm>. Accessed October 1, 2018.
153. Simpson DM, Hallett M, Ashman EJ, et al. Practice guideline update summary: Botulinum neurotoxin for the treatment of blepharospasm, cervical dystonia, adult spasticity, and headache: Report of the Guideline Development Subcommittee of the American Academy of Neurology. *Neurology.* May 10 2016;86(19):1818-1826. PMID 27164716
154. Zesiewicz TA, Elble RJ, Louis ED, et al. Evidence-based guideline update: treatment of essential tremor: report of the Quality Standards subcommittee of the American Academy of Neurology. *Neurology.* Nov 8 2011;77(19):1752-1755. PMID 22013182
155. American Academy of Neurology. Update: Treatment of Essential Tremor. 2011, reaffirmed 2014; <https://www.aan.com/Guidelines/>. Accessed October 1, 2018.
156. Quality Standards Subcommittee of the American Academy of N, the Practice Committee of the Child Neurology S, Delgado MR, et al. Practice parameter: pharmacologic treatment of spasticity in children and adolescents with cerebral palsy (an evidence-based review): report of the Quality

- Standards Subcommittee of the American Academy of Neurology and the Practice Committee of the Child Neurology Society. *Neurology*. Jan 26 2010;74(4):336-343. PMID 20101040
157. American Academy of Neurology. 2010 Pharmacologic Treatment of Spasticity in Children and Adolescents with Cerebral Palsy. 2010, reaffirmed 2013; <https://www.aan.com/Guidelines/>. Accessed October 1, 2018.
158. Naumann M, So Y, Argoff CE, et al. Assessment: Botulinum neurotoxin in the treatment of autonomic disorders and pain (an evidence-based review): report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology*. May 6 2008;70(19):1707-1714. PMID 18458231
159. Stewart DB, Sr., Gaertner W, Glasgow S, et al. Clinical practice guideline for the management of anal fissures. *Dis Colon Rectum*. Jan 2017;60(1):7-14. PMID 27926552

#8.01.19

1. Wade R, Rice S, Llewellyn A, et al. Interventions for hyperhidrosis in secondary care: a systematic review and value-of-information analysis. *Health Technol Assess*. Dec 2017;21(80):1-280. PMID 29271741
2. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Iontophoresis for Medical Indications. *TEC Assessments* 2003;Volume 18, Tab 3.
3. Rajagopal R, Mallya NB. Comparative evaluation of botulinum toxin versus iontophoresis with topical aluminium chloride hexahydrate in treatment of palmar hyperhidrosis. *Med J Armed Forces India*. Jul 2014;70(3):247-252. PMID 25378778
4. Solish N, Bertucci V, Dansereau A, et al. A comprehensive approach to the recognition, diagnosis, and severity-based treatment of focal hyperhidrosis: recommendations of the Canadian Hyperhidrosis Advisory Committee. *Dermatol Surg*. Aug 2007;33(8):908-923. PMID 17661933
5. Dogruk Kacar S, Ozuguz P, Eroglu S, et al. Treatment of primary hyperhidrosis with tap water iontophoresis in paediatric patients: a retrospective analysis. *Cutan Ocul Toxicol*. Dec 2014;33(4):313-316. PMID 24405389
6. McAleer MA, Collins P. A study investigating patients' experience of hospital and home iontophoresis for hyperhidrosis. *J Dermatolog Treat*. Aug 2014;25(4):342-344. PMID 23356798
7. Mirkovic SE, Rystedt A, Balling M, et al. Hyperhidrosis substantially reduces quality of life in children: a retrospective study describing symptoms, consequences and treatment with botulinum toxin. *Acta Derm Venereol*. Jan 12 2018;98(1):103-107. PMID 28761964
8. Lowe NJ, Glaser DA, Eadie N, et al. Botulinum toxin type A in the treatment of primary axillary hyperhidrosis: a 52-week multicenter double-blind, randomized, placebo-controlled study of efficacy and safety. *J Am Acad Dermatol*. Apr 2007;56(4):604-611. PMID 17306417
9. Baumann L, Slezinger A, Halem M, et al. Double-blind, randomized, placebo-controlled pilot study of the safety and efficacy of Myobloc (botulinum toxin type B) for the treatment of palmar hyperhidrosis. *Dermatol Surg*. Mar 2005;31(3):263-270. PMID 15841624
10. Baumann L, Slezinger A, Halem M, et al. Pilot study of the safety and efficacy of Myobloc (botulinum toxin type B) for treatment of axillary hyperhidrosis. *Int J Dermatol*. May 2005;44(5):418-424. PMID 15869543
11. Naumann MK, Hamm H, Lowe NJ, et al. Effect of botulinum toxin type A on quality of life measures in patients with excessive axillary sweating: a randomized controlled trial. *Br J Dermatol*. Dec 2002;147(6):1218-1226. PMID 12452874
12. Heckmann M, Ceballos-Baumann AO, Plewig G, et al. Botulinum toxin A for axillary hyperhidrosis (excessive sweating). *N Engl J Med*. Feb 15 2001;344(7):488-493. PMID 11172190
13. Dressler D. Comparing Botox and Xeomin for axillar hyperhidrosis. *J Neural Transm (Vienna)*. Mar 2010;117(3):317-319. PMID 20143241
14. Talarico-Filho S, Mendonca DO, Nascimento M, et al. A double-blind, randomized, comparative study of two type A botulinum toxins in the treatment of primary axillary hyperhidrosis. *Dermatol Surg*. Jan 2007;33(1 Spec No.):S44-50. PMID 17241414
15. Frasson E, Brigo F, Acler M, et al. Botulinum toxin type A vs type B for axillary hyperhidrosis in a case series of patients observed for 6 months. *Arch Dermatol*. Jan 2011;147(1):122-123. PMID 21242408
16. An JS, Hyun Won C, Si Han J, et al. Comparison of onabotulinumtoxinA and rimabotulinumtoxinB for the treatment of axillary hyperhidrosis. *Dermatol Surg*. Aug 2015;41(8):960-967. PMID 26218729

17. Lowe NJ, Yamauchi PS, Lask GP, et al. Efficacy and safety of botulinum toxin type a in the treatment of palmar hyperhidrosis: a double-blind, randomized, placebo-controlled study. *Dermatol Surg.* Sep 2002;28(9):822-827. PMID 12269876
18. Saadia D, Voustianiouk A, Wang AK, et al. Botulinum toxin type A in primary palmar hyperhidrosis: randomized, single-blind, two-dose study. *Neurology.* Dec 11 2001;57(11):2095-2099. PMID 11739832
19. Campanati A, Giuliadori K, Martina E, et al. Onabotulinumtoxin type A (Botox((R))) versus Incobotulinumtoxin type A (Xeomin((R))) in the treatment of focal idiopathic palmar hyperhidrosis: results of a comparative double-blind clinical trial. *J Neural Transm.* Jan 2014;121(1):21-26. PMID 24052109
20. Hsu TH, Chen YT, Tu YK, et al. A systematic review of microwave-based therapy for axillary hyperhidrosis. *J Cosmet Laser Ther.* Oct 2017;19(5):275-282. PMID 28281850
21. Glaser DA, Coleman WP, 3rd, Fan LK, et al. A randomized, blinded clinical evaluation of a novel microwave device for treating axillary hyperhidrosis: the dermatologic reduction in underarm perspiration study. *Dermatol Surg.* Feb 2012;38(2):185-191. PMID 22289389
22. Hong HC, Lupin M, O'Shaughnessy KF. Clinical evaluation of a microwave device for treating axillary hyperhidrosis. *Dermatol Surg.* May 2012;38(5):728-735. PMID 22452511
23. Purtuloglu T, Atim A, Deniz S, et al. Effect of radiofrequency ablation and comparison with surgical sympathectomy in palmar hyperhidrosis. *Eur J Cardiothorac Surg.* Jun 2013;43(6):e151-154. PMID 23428574
24. Hafner J, Beer GM. Axillary sweat gland excision. *Curr Probl Dermatol.* Dec 2002;30:57-63. PMID 12471699
25. Deng B, Tan QY, Jiang YG, et al. Optimization of sympathectomy to treat palmar hyperhidrosis: the systematic review and meta-analysis of studies published during the past decade. *Surg Endosc.* Jun 2011;25(6):1893-1901. PMID 21136103
26. Baumgartner FJ, Reyes M, Sarkisyan GG, et al. Thoracoscopic sympathectomy for disabling palmar hyperhidrosis: a prospective randomized comparison between two levels. *Ann Thorac Surg.* Dec 2011;92(6):2015-2019. PMID 22115211
27. Yuncu G, Turk F, Ozturk G, et al. Comparison of only T3 and T3-T4 sympathectomy for axillary hyperhidrosis regarding treatment effect and compensatory sweating. *Interact Cardiovasc Thorac Surg.* Aug 2013;17(2):263-267. PMID 23644731
28. de Andrade Filho LO, Kuzniec S, Wolosker N, et al. Technical difficulties and complications of sympathectomy in the treatment of hyperhidrosis: an analysis of 1731 cases. *Ann Vasc Surg.* May 2013;27(4):447-453. PMID 23406790
29. Karamustafaoglu YA, Kuzucuoglu M, Yanik F, et al. 3-year follow-up after uniportal thoracoscopic sympathectomy for hyperhidrosis: undesirable side effects. *J Laparoendosc Adv Surg Tech A.* Nov 2014;24(11):782-785. PMID 25376004
30. Smidfelt K, Drott C. Late results of endoscopic thoracic sympathectomy for hyperhidrosis and facial blushing. *Br J Surg.* Dec 2011;98(12):1719-1724. PMID 21928403
31. Wait SD, Killory BD, Lekovic GP, et al. Thoracoscopic sympathectomy for hyperhidrosis: analysis of 642 procedures with special attention to Horner's syndrome and compensatory hyperhidrosis. *Neurosurgery.* Sep 2010;67(3):652-656; discussion 656-657. PMID 20647968
32. Lembranca L, Wolosker N, de Campos JRM, et al. Videothoracoscopic sympathectomy results after oxybutynin chloride treatment failure. *Ann Vasc Surg.* Aug 2017;43:283-287. PMID 28478174
33. de Campos JRM, Lembranca L, Fukuda JM, et al. Evaluation of patients who underwent resympathectomy for treatment of primary hyperhidrosis. *Interact Cardiovasc Thorac Surg.* Nov 1 2017;25(5):716-719. PMID 29049566
34. Fukuda JM, Varella AYM, Teivelis MP, et al. Video-Assisted thoracoscopic sympathectomy for facial hyperhidrosis: the influence of the main site of complaint. *Ann Vasc Surg.* Jan 2018;46:337-344. PMID 28689957
35. Rieger R, Pedevilla S, Pochlauer S. Endoscopic lumbar sympathectomy for plantar hyperhidrosis. *Br J Surg.* Dec 2009;96(12):1422-1428. PMID 19918855
36. Reisfeld R. Endoscopic lumbar sympathectomy for focal plantar hyperhidrosis using the clamping method. *Surg Laparosc Endosc Percutan Tech.* Aug 2010;20(4):231-236. PMID 20729691
37. Hornberger J, Grimes K, Naumann M, et al. Recognition, diagnosis, and treatment of primary focal hyperhidrosis. *J Am Acad Dermatol.* Aug 2004;51(2):274-286. PMID 15280848

38. Li C, Wu F, Zhang Q, et al. Interventions for the treatment of Frey's syndrome. *Cochrane Database Syst Rev*. Mar 17 2015;3(3):CD009959. PMID 25781421
39. Clayman MA, Clayman SM, Seagle MB. A review of the surgical and medical treatment of Frey syndrome. *Ann Plast Surg*. Nov 2006;57(5):581-584. PMID 17060744
40. de Bree R, van der Waal I, Leemans CR. Management of Frey syndrome. *Head Neck*. Aug 2007;29(8):773-778. PMID 17230557
41. Cerfolio RJ, De Campos JR, Bryant AS, et al. The Society of Thoracic Surgeons expert consensus for the surgical treatment of hyperhidrosis. *Ann Thorac Surg*. May 2011;91(5):1642-1648. PMID 21524489
42. Naumann M, So Y, Argoff CE, et al. Assessment: Botulinum neurotoxin in the treatment of autonomic disorders and pain (an evidence-based review): report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology*. May 6 2008;70(19):1707-1714. PMID 18458231
43. Naumann M, Dressler D, Hallett M, et al. Evidence-based review and assessment of botulinum neurotoxin for the treatment of secretory disorders. *Toxicon*. Jun 1 2013;67:141-152. PMID 23178324
44. National Institute of Health and Care Excellence (NICE). Endoscopic thoracic sympathectomy for primary facial blushing [IPG480]. 2014; <https://www.nice.org.uk/guidance/ipg480>. Accessed March 1, 2018.
45. National Institute of Health and Care Excellence (NICE). Endoscopic throacic sympathectomy for primary hyperhidrosis of the upper limb [IPG487]. 2014; <https://www.nice.org.uk/guidance/ipg487>. Accessed March 1, 2018.

Endnotes

1. FDA-approved indications
2. From National Blue Cross Blue Shield Association policy 5.01.05
3. Local Medicare policy <http://www.medicarenhic.com/> and CMS guidelines http://www.hcfa.gov/pubforms/14%5Fcar/3b2049.htm#_1_7.
4. Intrasphincteric botulinum toxin for the treatment of achalasia. *NEJM* 1995 March 23;322:744-8, by Paricha et al.
5. A comparison of Botulinum toxin an saline for the treatment of chronic anal fissure, *NEJM* 1998 Jan 22;338:217-20 by Maria et al. Based upon advice from representative physician experts of the Massachusetts Neurologic Association, 2000.
6. Recommendations from Joel Stein, MD; Massachusetts General Hospital, EBR 1/01.
7. 12/00 FDA-approved indication
8. In accordance with local Medicare policy published in the September 2001 Medicare B Resource. See also: www.medicarenhic.gov.
9. Based upon the 2002 and 2004 Blue Cross Blue Shield Association national policy 5.01.05.
10. Pediatrics volume 108 number 5 November 2001. Botulinum toxin type A neuromuscular blockade in the treatment of Equinus deformity in cerebral palsy. Based upon the 2002 and 2004 Blue Cross Blue Shield Association National policy 5.01.05. Foster L, Clapp L, Erickson M et al. Botulinum toxin A and chronic low back pain: a randomized, double-blind study. *Neurology* 2001;56(10):1290-3.
11. Based upon the 2002 and 2004 Blue Cross Blue Shield Association National policy 5.01.05. The National policy offered the following rationale:
12. Individual consideration guideline for cervicogenic migraine headache is based upon local expert opinion.
13. Based upon the 2004 Blue Cross Blue Shield Association National policy 5.01.05. Botulinum toxin for Sialorrhea (drooling).
Based upon the 2005 Blue Cross Blue Shield Association National policy 5.01.05 issued 12/05.
14. Based upon the 2005 Blue Cross Blue Shield Association National policy 8.01.19 Treatment of Hyperhidrosis issued 12/05 with a coding update only. Based upon Blue Cross Blue Shield National Policy 5.01.05 Botulinum Toxin issued 4/06.
15. Dysport™ [package insert]. Brisbane, CA: Tercica, Inc., May 2009.
16. Xeomin® [package insert]. Greensboro, NC: Merz Pharmaceuticals, LLC.; August 2010.

17. "FDA Approves Botox to Treat Chronic Migraines." FDA.gov. Food and Drug Administration, 15 Oct. 2010. Web. 23 Mar. 2011.
<http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm229782.htm>.

To request prior authorization using the Massachusetts Standard Form for Medication Prior Authorization Requests (eForm), click the link below:

http://www.bluecrossma.com/common/en_US/medical_policies/023%20E%20Form%20medication%20prior%20auth%20instruction%20prn.pdf