



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an independent
Licensee of the Blue Cross and Blue Shield Association

Medical Policy

Lung and Lobar Lung Transplant

Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

Policy Number: 015

BCBSA Reference Number: 7.03.07

NCD/LCD: NA

Related Policies

- Heart/Lung Transplant, #[269](#)
- Outpatient Pulmonary Rehabilitation, #[136](#)

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO BlueSM and Medicare PPO BlueSM Members

Lung transplantation may be considered [MEDICALLY NECESSARY](#) for carefully selected patients with irreversible, progressively disabling, end-stage pulmonary disease unresponsive to maximum medical therapy.

Lobar lung transplant from a living or deceased donor may be considered [MEDICALLY NECESSARY](#) for carefully selected patients with end-stage pulmonary disease.

Etiologies of End-Stage Lung Disease

- Bilateral bronchiectasis
- Alpha-1 antitrypsin deficiency
- Primary pulmonary hypertension
- Cystic fibrosis (both lungs to be transplanted)
- Bronchopulmonary dysplasia
- Postinflammatory pulmonary fibrosis
- Idiopathic/interstitial pulmonary fibrosis
- Sarcoidosis
- Scleroderma
- Lymphangiomyomatosis
- Emphysema
- Eosinophilic granuloma
- Bronchiolitis obliterans

- Recurrent pulmonary embolism
- Pulmonary hypertension due to cardiac disease
- Chronic obstructive pulmonary disease
- Eisenmenger's syndrome.

Lung or lobar lung transplants in patients with any of the following conditions are **NOT MEDICALLY NECESSARY**:

1. Known active malignancy, including metastatic cancer
2. Recently treated malignancy with a high risk of recurrence
Note: *the assessment of risk of recurrence of a recently treated malignancy is made by the transplant team; providers must submit a statement with an explanation of why the patient with a recently treated malignancy is an appropriate candidate for a transplant.*
3. Untreated systemic infection making immunosuppression unsafe, including chronic infection
4. Other irreversible end-stage disease not attributed to lung disease
5. History of cancer with a moderate risk of recurrence
6. Systemic disease that could be exacerbated by immunosuppression
7. Psychosocial conditions or chemical dependence affecting the ability to adhere to therapy
8. Coronary artery disease not amenable to percutaneous intervention or bypass grafting, or associated with significant impairment of left ventricular function, or
9. Colonization with highly resistant or highly virulent bacteria, fungi, or mycobacteria.

Lung or lobar lung retransplantation after a failed lung or lobar lung transplant may be considered **MEDICALLY NECESSARY** in patients who meet criteria for lung transplantation.

Lung or lobar lung transplantation is considered **INVESTIGATIONAL** in all other situations.

Prior Authorization Information

Pre-service approval is required for all inpatient services for all products.

See below for situations where prior authorization may be required or may not be required for outpatient services.

Yes indicates that prior authorization is required.

No indicates that prior authorization is not required.

N/A indicates that this service is primarily performed in an inpatient setting.

	Outpatient
Commercial Managed Care (HMO and POS)	N/A
Commercial PPO and Indemnity	N/A
Medicare HMO BlueSM	N/A
Medicare PPO BlueSM	N/A

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

The above medical necessity criteria MUST be met for the following codes to be covered for Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:

CPT Codes

CPT codes:	Code Description
32851	Lung transplant, single; without cardiopulmonary bypass
32852	Lung transplant, single; with cardiopulmonary bypass
32853	Lung transplant, double (bilateral sequential or en bloc); without cardiopulmonary bypass
32854	Lung transplant, double (bilateral sequential or en bloc); with cardiopulmonary bypass

HCPCS Codes

HCPCS codes:	Code Description
S2060	Lobar lung transplantation

ICD-10 Procedure Codes

ICD-10-CM Diagnosis codes:	Code Description
0BYC0Z0	Transplantation of Right Upper Lung Lobe, Allogeneic, Open Approach
0BYC0Z1	Transplantation of Right Upper Lung Lobe, Syngeneic, Open Approach
0BYD0Z0	Transplantation of Right Middle Lung Lobe, Allogeneic, Open Approach
0BYD0Z1	Transplantation of Right Middle Lung Lobe, Syngeneic, Open Approach
0BYF0Z0	Transplantation of Right Lower Lung Lobe, Allogeneic, Open Approach
0BYF0Z1	Transplantation of Right Lower Lung Lobe, Syngeneic, Open Approach
0BYG0Z0	Transplantation of Left Upper Lung Lobe, Allogeneic, Open Approach
0BYG0Z1	Transplantation of Left Upper Lung Lobe, Syngeneic, Open Approach
0BYH0Z0	Transplantation of Lung Lingula, Allogeneic, Open Approach
0BYH0Z1	Transplantation of Lung Lingula, Syngeneic, Open Approach
0BYJ0Z0	Transplantation of Left Lower Lung Lobe, Allogeneic, Open Approach
0BYJ0Z1	Transplantation of Left Lower Lung Lobe, Syngeneic, Open Approach
0BYK0Z0	Transplantation of Right Lung, Allogeneic, Open Approach
0BYK0Z1	Transplantation of Right Lung, Syngeneic, Open Approach
0BYL0Z0	Transplantation of Left Lung, Allogeneic, Open Approach
0BYL0Z1	Transplantation of Left Lung, Syngeneic, Open Approach
0BYM0Z0	Transplantation of Bilateral Lungs, Allogeneic, Open Approach
0BYM0Z1	Transplantation of Bilateral Lungs, Syngeneic, Open Approach

DESCRIPTION

END-STAGE LUNG DISEASE

End-stage lung disease may derive from different etiologies. The most common indications for lung transplantation are chronic obstructive pulmonary disease, idiopathic pulmonary fibrosis, cystic fibrosis, α_1 -antitrypsin deficiency, and idiopathic pulmonary arterial hypertension.

Treatment

Before consideration for transplant, patients should be receiving maximal medical therapy, including oxygen supplementation, or surgical options, such as lung-volume reduction surgery for chronic obstructive pulmonary disease. Lung or lobar lung transplantation is an option for patients with end-stage lung disease despite these measures.

A lung transplant refers to single-lung or double-lung replacement. In a single-lung transplant, only 1 lung from a deceased donor is provided to the recipient. In a double-lung transplant, both the recipient's lungs are removed and replaced by the donor's lungs. In a lobar transplant, a lobe of the donor's lung is excised, sized appropriately for the recipient's thoracic dimensions, and transplanted. Donors for lobar transplant have primarily been living-related donors, with 1 lobe obtained from each of 2 donors

(generally friends or family members) in cases for which bilateral transplantation is required. There are also cases of cadaver lobe transplants.

Since 2005, potential recipients have been ranked according to the Lung Allocation Score.¹ Patients 12 years of age and older receive a score between 1 and 100 based on predicted survival after transplantation reduced by predicted survival on the waiting list; the Lung Allocation Score takes into consideration the patient’s disease and clinical parameters. In 2010, a simple priority system was implemented for children younger than age 12 years. Under this system, children younger than 12 years with respiratory lung failure and/or pulmonary hypertension who meet criteria are considered “priority 1” and all other candidates in the age group are considered “priority 2”. A lung review board has the authority to adjust scores on appeal for adults and children.

Summary

A lung transplant consists of replacing all or part of diseased lungs with healthy lung(s) or lobes. Transplantation is an option for patients with end-stage lung disease.

For individuals who have end-stage pulmonary disease who receive a lung transplant, the evidence includes case series and registry studies. Relevant outcomes are overall survival, change in disease status, and treatment-related mortality and morbidity. International registry data on a large number of patients receiving lung transplantation (>50,000) found relatively high patient survival rates, especially among those who survived the first year posttransplant. After adjusting for potential confounding factors, survival did not differ significantly after single- or double-lung transplant. Lung transplantation may be the only option for some patients with end-stage lung disease. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have end-stage pulmonary disease who receive a lobar lung transplant, the evidence includes case series and systematic reviews. Relevant outcomes are overall survival, change in disease status, and treatment-related mortality and morbidity. There are less data on lung lobar transplants than on whole-lung transplants, but several case series have reported reasonably similar survival outcomes between the procedures, and lung lobar transplants may be the only option for patients unable to wait for a whole-lung transplant. A 2017 systematic review found 1-year survival rates in available published studies ranging from 50% to 100%. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have a prior lung or lobar transplant who meet criteria for a lung transplant who receive a lung or lobar lung retransplant, the evidence includes case series and registry studies. Relevant outcomes are overall survival, change in disease status, treatment-related mortality and morbidity. Data from registries and case series have found favorable outcomes with lung retransplantation in patients who meet criteria for initial lung transplantation. Given the exceedingly poor survival prognosis without retransplantation of patients who have exhausted other treatments, the evidence of a moderate level of posttransplant survival may be considered sufficient in this patient population. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

Policy History

Date	Action
10/2018	BCBSA National medical policy review. Description, summary and references updated. Policy statement(s) unchanged.
10/2018	BCBSA National medical policy review. No changes to policy statements. New references added. Summary clarified.
1/2018	Clarified coding information.
10/2017	BCBSA National medical policy review. Conditions for covered indications moved to Policy Guidelines. 10/1/2017
8/2015	Added coding language.
3/2015	New references added from BCBSA National medical policy.

10/2014	Coding information clarified.
6/2014	BCBSA National medical policy review. New medically necessary and investigational indications described. Effective 6/1/2014.
5/2014	Updated Coding section with ICD10 procedure and diagnosis codes, effective 10/2015.
12/2013	Removed ICD-9 diagnosis codes as the policy requires prior authorization
6/2013	BCBSA National medical policy review. In lobar lung statement, “children and adolescents” replaced with “carefully selected patients.” Effective 6/1/2013.
11/2011-4/2012	Medical policy ICD 10 remediation: Formatting, editing and coding updates. No changes to policy statements.
10/2011	Reviewed - Medical Policy Group – GI, Nutrition and Organ Transplantation. No changes to policy statements.
3/2011	National Policy Review. Changes to policy statement.
11/2010	Reviewed - Medical Policy Group – Gastroenterology, Nutrition and Organ Transplantation. No changes to policy statements.
11/2009	Reviewed - Medical Policy Group – Gastroenterology, Nutrition and Organ Transplantation. No changes to policy statements.
9/2009	National Policy Review. No changes to policy statements.
6/2009	National Policy Review. No changes to policy statements.
11/2008	Reviewed - Medical Policy Group – Gastroenterology, Nutrition & Organ Transplants, No changes to policy statements.
5/2008	National Policy Review. Policy guidelines updated. No changes to policy statements.
11/2007	Reviewed - Medical Policy Group – Gastroenterology, Nutrition and Organ Transplants. No changes to policy statements.
11/2006	Reviewed - Medical Policy Group – Gastroenterology, Nutrition and Organ Transplants. No changes to policy statements.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

References

1. Organ Procurement and Transplantation Network (OPTN). Policy 10: Allocation of Lungs. 2018; https://optn.transplant.hrsa.gov/media/1200/optn_policies.pdf. Accessed July 20, 2018.
2. Paraskeva MA, Edwards LB, Levvey B, et al. Outcomes of adolescent recipients after lung transplantation: An analysis of the International Society for Heart and Lung Transplantation Registry. *J Heart Lung Transplant*. Feb 17 2018;37(3):323-331. PMID 28320631
3. Yusen RD, Edwards LB, Dipchand AI, et al. The Registry of the International Society for Heart and Lung Transplantation: Thirty-third Adult Lung and Heart-Lung Transplant Report-2016; Focus Theme: Primary Diagnostic Indications for Transplant. *J Heart Lung Transplant*. Oct 2016;35(10):1170-1184. PMID 27772669
4. Goldfarb SB, Levvey BJ, Edwards LB, et al. The Registry of the International Society for Heart and Lung Transplantation: Nineteenth Pediatric Lung and Heart-Lung Transplantation Report-2016; Focus Theme: Primary Diagnostic Indications for Transplant. *J Heart Lung Transplant*. Oct 2016;35(10):1196-1205. PMID 27772671
5. Thabut G, Christie JD, Kremers WK, et al. Survival differences following lung transplantation among US transplant centers. *JAMA*. Jul 07 2010;304(1):53-60. PMID 20606149
6. Black MC, Trivedi J, Schumer EM, et al. Double lung transplants have significantly improved survival compared with single lung transplants in high lung allocation score patients. *Ann Thorac Surg*. Nov 2014;98(5):1737-1741. PMID 25110334

7. Yusen RD, Shearon TH, Qian Y, et al. Lung transplantation in the United States, 1999-2008. *Am J Transplant.* Apr 2010;10(4 Pt 2):1047-1068. PMID 20420652
8. Shafii AE, Mason DP, Brown CR, et al. Too high for transplantation? Single-center analysis of the lung allocation score. *Ann Thorac Surg.* Nov 2014;98(5):1730-1736. PMID 25218678
9. Date H. Update on living-donor lobar lung transplantation. *Curr Opin Organ Transplant.* Oct 2011;16(5):453-457. PMID 21836512
10. Eberlein M, Reed RM, Chahla M, et al. Lobar lung transplantation from deceased donors: A systematic review. *World J Transplant.* Feb 24 2017;7(1):70-80. PMID 28280698
11. Barr ML, Schenkel FA, Bowdish ME, et al. Living donor lobar lung transplantation: current status and future directions. *Transplant Proc.* Nov 2005;37(9):3983-3986. PMID 16386604
12. Date H, Sato M, Aoyama A, et al. Living-donor lobar lung transplantation provides similar survival to cadaveric lung transplantation even for very ill patients. *Eur J Cardiothorac Surg.* Jun 2015;47(6):967-972; discussion 972-963. PMID 25228745
13. Date H, Shiraishi T, Sugimoto S, et al. Outcome of living-donor lobar lung transplantation using a single donor. *J Thorac Cardiovasc Surg.* Sep 2012;144(3):710-715. PMID 22717276
14. Slama A, Ghanim B, Kliekovits T, et al. Lobar lung transplantation--is it comparable with standard lung transplantation? *Transpl Int.* Sep 2014;27(9):909-916. PMID 24810771
15. Yusen RD, Christie JD, Edwards LB, et al. The Registry of the International Society for Heart and Lung Transplantation: thirtieth adult lung and heart-lung transplant report--2013; focus theme: age. *J Heart Lung Transplant.* Oct 2013;32(10):965-978. PMID 24054805
16. Kilic A, Beaty CA, Merlo CA, et al. Functional status is highly predictive of outcomes after redo lung transplantation: an analysis of 390 cases in the modern era. *Ann Thorac Surg.* Nov 2013;96(5):1804-1811. PMID 23968759
17. Kawut SM. Lung retransplantation. *Clin Chest Med.* Jun 2011;32(2):367-377. PMID 21511096
18. Biswas Roy S, Panchanathan R, Walia R, et al. Lung retransplantation for chronic rejection: a single-center experience. *Ann Thorac Surg.* Nov 1 2018;105(1):221-227. PMID 29100649
19. Organ Procurement and Transplantation Network (OPTN). National Data. n.d.; <https://optn.transplant.hrsa.gov/data/view-data-reports/national-data/>. Accessed July 20, 2018.
20. Organ Procurement and Transplantation Network (OPTN). Organ Procurement and Transplantation Network Policies. 2018; https://optn.transplant.hrsa.gov/media/1200/optn_policies.pdf. Accessed July 20, 2018.
21. Working Party of the British Transplantation Society. *Kidney and Pancreas Transplantation in Patients with HIV. Second Edition (Revised). British Transplantation Society Guidelines.* Macclesfield, UK: British Transplantation Society; 2017.
22. Alexander BD, Petzold EW, Reller LB, et al. Survival after lung transplantation of cystic fibrosis patients infected with Burkholderia cepacia complex. *Am J Transplant.* May 2008;8(5):1025-1030. PMID 18318775
23. Murray S, Charbeneau J, Marshall BC, et al. Impact of burkholderia infection on lung transplantation in cystic fibrosis. *Am J Respir Crit Care Med.* Aug 15 2008;178(4):363-371. PMID 18535253
24. Boussaud V, Guillemain R, Grenet D, et al. Clinical outcome following lung transplantation in patients with cystic fibrosis colonised with Burkholderia cepacia complex: results from two French centres. *Thorax.* Aug 2008;63(8):732-737. PMID 18408050
25. Orens JB, Estenne M, Arcasoy S, et al. International guidelines for the selection of lung transplant candidates: 2006 update--a consensus report from the Pulmonary Scientific Council of the International Society for Heart and Lung Transplantation. *J Heart Lung Transplant.* Jul 2006;25(7):745-755. PMID 16818116
26. Weill D, Benden C, Corris PA, et al. A consensus document for the selection of lung transplant candidates: 2014--an update from the Pulmonary Transplantation Council of the International Society for Heart and Lung Transplantation. *J Heart Lung Transplant* Jun 26 2015 34(1):1-15. PMID 25085497
27. Raghu G, Collard HR, Egan JJ, et al. An official ATS/ERS/JRS/ALAT statement: idiopathic pulmonary fibrosis: evidence-based guidelines for diagnosis and management. *Am J Respir Crit Care Med.* Mar 15 2011;183(6):788-824. PMID 21471066
28. Centers for Medicare & Medicaid. Medicare-approved transplant programs: Lung-only. 2018; <https://www.cms.gov/Medicare/Provider-Enrollment-and->

Certification/CertificationandCompliance/downloads/ApprovedTransplantPrograms.pdf. Accessed July 20, 2018.