



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

## Medical Policy

# Surgical Treatment of Femoroacetabular Impingement

### Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

### Policy Number: 145

BCBSA Reference Number: 7.01.118

NCD/LCD: N/A

### Related Policies

- Hip Resurfacing, #[046](#)
- Surgery for Groin Pain in Athletes, #[695](#)

### Policy

#### Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO Blue<sup>SM</sup> and Medicare PPO Blue<sup>SM</sup> Members

Open or arthroscopic treatment of femoroacetabular impingement (FAI) may be **MEDICALLY NECESSARY** when **ALL** the age, symptom and imaging conditions below have been met:

#### Age

- Candidates should be skeletally mature with documented closure of growth plates (e.g., 15 years or older).

#### Symptoms

- Moderate-to-severe hip pain that is worsened by flexion activities (e.g., squatting or prolonged sitting) that significantly limits activities, AND
- Unresponsive to conservative therapy for at least 3 months (including activity modifications, restriction of athletic pursuits and avoidance of symptomatic motion), AND
- Positive impingement sign on clinical examination (pain elicited with 90 degrees of flexion and internal rotation and adduction of the femur).

#### Imaging

- Morphology indicative of cam or pincer-type FAI, e.g., pistol-grip deformity, femoral head-neck offset with an alpha angle greater than 50 degrees, a positive wall sign, acetabular retroversion (overcoverage with crossover sign), coxa profunda or protrusion, or damage of the acetabular rim, AND

- High probability of a causal association between the FAI morphology and damage, e.g., a pistol-grip deformity with a tear of the acetabular labrum and articular cartilage damage in the anterosuperior quadrant, AND
- No evidence of advanced osteoarthritis, defined as Tonnis grade II or III, or joint space of less than 2 mm, AND
- No evidence of severe (Outerbridge grade IV) chondral damage.

Treatment of femoroacetabular impingement in all other situations is **INVESTIGATIONAL**.

## Prior Authorization Information

### Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

### Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

|  | Outpatient  |
|--|---|
| <b>Commercial Managed Care (HMO and POS)</b> | Prior authorization is <b><u>not required</u></b> . |
| <b>Commercial PPO and Indemnity</b>          | Prior authorization is <b><u>not required</u></b> . |
| <b>Medicare HMO Blue<sup>SM</sup></b>        | Prior authorization is <b><u>not required</u></b> . |
| <b>Medicare PPO Blue<sup>SM</sup></b>        | Prior authorization is <b><u>not required</u></b> . |

## CPT Codes / HCPCS Codes / ICD Codes

*Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.*

*Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.*

### CPT Codes

| CPT codes: | Code Description  |
|------------|---|
| 29914      | Arthroscopy, hip, surgical; with femoroplasty (ie, treatment of cam lesion)       |
| 29915      | Arthroscopy, hip, surgical; with acetabuloplasty (ie, treatment of pincer lesion) |
| 29916      | Arthroscopy, hip, surgical; with labral repair                                    |

## Description

### Femoroacetabular Impingement

FAI arises from an anatomic mismatch between the head of the femur and the acetabulum, causing compression of the labrum or articular cartilage during flexion. The mismatch can arise from subtle morphologic alterations in the anatomy or orientation of the ball-and-socket components (eg, a bony prominence at the head-neck junction or acetabular overcoverage), with articular cartilage damage initially occurring from abutment of the femoral neck against the acetabular rim, typically at the anterosuperior aspect of the acetabulum. Although hip joints can possess the morphologic features of FAI without symptoms, FAI may become pathologic with repetitive movement and/or increased force on the hip joint. High-demand activities may also result in pathologic impingement in hips with normal morphology.

Two types of impingement, cam, and pincer may occur alone or, more frequently, together. Cam impingement is associated with an asymmetric or nonspherical contour of the head or neck of the femur jamming against the acetabulum, resulting in cartilage damage and delamination (detachment from the subchondral bone). Deformity of the head/neck junction that looks like a pistol-grip on radiographs is associated with damage to the anterosuperior area of the acetabulum. Symptomatic cam impingement is

found most frequently in young male athletes. Pincer impingement is associated with overcoverage of the acetabulum and pinching of the labrum, with pain more typically beginning in women of middle age. In cases of isolated pincer impingement, the damage may be limited to a narrow strip of the acetabular cartilage.

Epidemiologic and radiographic studies have found correlations between hip osteoarthritis (OA) and FAI lesions, supporting the theory that prolonged contact between the anatomically mismatched acetabulum and femur may lead not only to cam and pincer lesions but also to further cartilage damage and subsequent joint deterioration. It is believed that osteoplasty of the impinging bone is needed to protect the cartilage from further damage and to preserve the natural joint. Therefore, if FAI morphology is shown to be an etiology of OA, a strategy to reduce the occurrence of idiopathic hip OA could be early recognition and treatment of FAI before cartilage damage and joint deterioration occurs.

An association between FAI and athletic pubalgia, sometimes called sports hernia, has been proposed. Athletic pubalgia is an umbrella term for a large variety of musculoskeletal injuries involving attachments and/or soft tissue support structures of the pubis (see medical policy #695, Surgery for Groin Pain in Athletes).

### **Treatment**

A technique for hip dislocation with open osteochondroplasty that preserved the femoral blood supply was reported by Ganz. Visualization of the entire joint with this procedure led to the identification and acceptance of FAI as an etiology of cartilage damage and the possibility of correcting the abnormal femoroacetabular morphology. Open osteochondroplasty of bony abnormalities and treatment of the symptomatic cartilage defect is considered the criterion standard for complex bony abnormalities. However, open osteochondroplasty is invasive, requiring transection of the greater trochanter (separation of the femoral head from the femoral shaft) and dislocation of the hip joint to provide full access to the femoral head and acetabulum. In addition to the general adverse events of open surgical procedures, open osteochondroplasty with dislocation has been associated with nonunion and neurologic and soft tissue lesions.

Less invasive hip arthroscopy and an arthroscopy-assisted mini-approach were developed by 2004. Arthroscopy requires specially designed instruments and is considered technically more difficult due to reduced visibility and limited access to the joint space. Advanced imaging techniques, including computed tomography and fluoroscopy, have been used to improve visualization of the 3-dimensional head/neck morphology during arthroscopy.

FAI can also be a source of hip pain and decreased hip internal rotation in the pediatric population. When nonoperative management of FAI in children and adolescents is ineffective, surgical procedures may be indicated. Surgical techniques include arthroscopy, open hip dislocation, limited open with arthroscopy, and osteotomy.

### **Slipped Capital Femoral Epiphysis**

Patients with slipped capital femoral epiphysis (SCFE) have a displaced femoral head in relation to the femoral neck within the confines of the acetabulum, which can result in hip pain, thigh pain, knee pain, and the onset of a limp. SCFE occurs most frequently in children between the ages of 10 to 16. Upon reaching skeletal maturity patients diagnosed with SCFE, 32% were found to have clinical signs of impingement. It is not uncommon for patients with SCFE to develop premature OA and require total hip arthroplasty within 20 years.

### **Treatment**

The standard treatment for SCFE is stabilization across the physis by in situ pinning. Alternative treatments proposed for pediatric patients with SCFE-related FAI include osteoplasty without dislocation, or with the open dislocation technique described by Ganz. The Ganz technique (capital realignment with open dislocation) is technically demanding, with a steep learning curve and a high-risk of complications, including avascular necrosis. Therefore, early treatment to decrease impingement must be weighed against the increased risk of adverse events.

## Summary

Femoroacetabular impingement (FAI) results from localized compression within the joint as a result of an anatomic mismatch between the head of the femur and the acetabulum. Symptoms of impingement typically occur in young to middle-aged adults before the onset of osteoarthritis (OA) but may be present in younger patients with developmental hip disorders. The objective of surgical treatment of FAI is to provide symptom relief and reduce further joint damage.

For individuals who are adults with asymptomatic FAI who receive FAI surgery, there is no direct evidence that the surgical treatment will prevent the development of OA. The relevant outcomes are symptoms, functional outcomes, health status measures, quality of life (QOL), and change in disease status. Indirect evidence consists of observational studies. In retrospective studies of patients with OA, the relevant outcomes were radiographic evidence of hip joint malformations. In prospective studies of patients with FAI, the relevant outcome is progression to OA. Several large observational studies (>1000 patients), as well as smaller studies, have shown radiographic evidence of relationships between abnormal hip morphology and the development of OA. There have been no studies in which FAI surgery was performed on patients with FAI morphology but no symptoms. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who are adults with symptomatic FAI who receive FAI surgery, the evidence includes systematic reviews of large and small observational studies and a small randomized controlled trial. The relevant outcomes are symptoms, functional outcomes, health status measures, QOL, and change in disease status. Open hip dislocation surgery and arthroscopic surgery are the most common surgical techniques performed on patients with FAI. Systematic reviews have evaluated open hip dislocation surgery and arthroscopic surgery, compared with no comparator, nonsurgical management, and other surgical techniques. Compared with nonsurgical management, all types of surgical techniques have resulted in significant improvements in functional outcomes, pain, and radiographic measurements. The reviews were limited when comparing surgical techniques with each other because patient characteristics and outcome measurements were heterogeneous among studies. The evidence is sufficient to determine the technology results in a meaningful improvement in the net health outcome.

For individuals who are children 15 years of age or younger with symptomatic FAI who receive FAI surgery, the evidence includes systematic reviews evaluating small observational studies and case series. The relevant outcomes are symptoms, functional outcomes, health status measures, QOL, and change in disease status. While the studies reported reductions in pain and improvements in functional outcomes, the sample sizes were relatively small, with an average of 54 patients per study. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who are children 15 years of age or younger with slipped capital femoral epiphysis-associated FAI who receive FAI surgery, the evidence includes small observational studies (range, 19-51 patients). The relevant outcomes are symptoms, functional outcomes, health status measures, QOL, and change in disease status. While most patients experienced symptom relief following FAI surgery, the surgery is invasive and complications (eg, nonunions) were reported. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have residual FAI symptoms following a primary surgery who receive revision arthroscopic surgery, the evidence includes systematic reviews of observational studies (>400 patients). The relevant outcomes are symptoms, functional outcomes, health status measures, QOL, and change in disease status. Though the studies were of low-quality, consistent improvements in functional outcomes, pain relief, and patient satisfaction were reported, in some cases beyond three years. The evidence is sufficient to determine the technology results in a meaningful improvement in the net health outcome.

## Policy History

| Date | Action |
|------|--------|
|------|--------|

|                |  |
|----------------|--|
| 5/2019         | BCBSA National medical policy review. Description, summary and references updated. Policy statements unchanged.  |
| 5/2018         | New references added from BCBSA National medical policy. Background and summary clarified.   |
| 5/2017         | New references added from BCBSA National medical policy.   |
| 7/2015         | New references added from BCBSA National medical policy.   |
| 9/2014         | New references added from BCBSA National medical policy.   |
| 11/1/2013      | BCBSA National medical policy review.<br>Age restriction on older adults removed; age restriction on pediatric patients clarified.<br>Effective 11/1/2013. |
| 11/2011-4/2012 | Medical policy ICD 10 remediation: Formatting, editing and coding updates.<br>No changes to policy statements.   |
| 6/2011         | Medical Policy Group - Orthopedics, Rehabilitation and Rheumatology.<br>No changes to policy statements.   |
| 7/2010         | Medical Policy Group - Orthopedics, Rehabilitation Medicine and Rheumatology.<br>No changes to policy statements.  |
| 12/1/09        | Medical Policy 145 effective 12/1/09.  |

## Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

## References

1. Oner A, Koksal A, Sofu H, et al. The prevalence of femoroacetabular impingement as an aetiologic factor for end-stage degenerative osteoarthritis of the hip joint: analysis of 1,000 cases. *Hip Int.* Mar-Apr 2016;26(2):164- 168. PMID 26916653.
2. Beck M, Kalhor M, Leunig M, et al. Hip morphology influences the pattern of damage to the acetabular cartilage: femoroacetabular impingement as a cause of early osteoarthritis of the hip. *J Bone Joint Surg Br.* Jul 2005;87(7):1012-1018. PMID 15972923.
3. Reichenbach S, Leunig M, Werlen S, et al. Association between cam-type deformities and magnetic resonance imaging-detected structural hip damage: a cross-sectional study in young men. *Arthritis Rheum.* Dec 2011;63(12):4023-4030. PMID 21904996.
4. Thomas GE, Palmer AJ, Batra RN, et al. Subclinical deformities of the hip are significant predictors of radiographic osteoarthritis and joint replacement in women. A 20 year longitudinal cohort study. *Osteoarthritis Cartilage.* Oct 2014;22(10):1504-1510. PMID 25047637.
5. Gosvig KK, Jacobsen S, Sonne-Holm S, et al. Prevalence of malformations of the hip joint and their relationship to sex, groin pain, and risk of osteoarthritis: a population-based survey. *J Bone Joint Surg Am.* May 2010;92(5):1162-1169. PMID 20439662.
6. Takeyama A, Naito M, Shiramizu K, et al. Prevalence of femoroacetabular impingement in Asian patients with osteoarthritis of the hip. *Int Orthop.* Oct 2009;33(5):1229-1232. PMID 19277653.
7. Tanzer M, Noiseux N. Osseous abnormalities and early osteoarthritis: the role of hip impingement. *Clin Orthop Relat Res.* Dec 2004(429):170-177. PMID 15577483.
8. Wall PD, Brown JS, Parsons N, et al. Surgery for treating hip impingement (femoroacetabular impingement). *Cochrane Database Syst Rev.* Sep 8 2014;9(9):CD010796. PMID 25198064.
9. Harris JD, Erickson BJ, Bush-Joseph CA, et al. Treatment of femoroacetabular impingement: a systematic review. *Curr Rev Musculoskelet Med.* Sep 2013;6(3):207-218. PMID 23743861.
10. Bedi A, Chen N, Robertson W, et al. The management of labral tears and femoroacetabular impingement of the hip in the young, active patient. *Arthroscopy.* Oct 2008;24(10):1135-1145. PMID 19028166.
11. Beck M, Leunig M, Parvizi J, et al. Anterior femoroacetabular impingement: part II. Midterm results of surgical treatment. *Clin Orthop Relat Res.* Jan 2004(418):67-73. PMID 15043095.

12. Espinosa N, Rothenfluh DA, Beck M, et al. Treatment of femoro-acetabular impingement: preliminary results of labral refixation. *J Bone Joint Surg Am*. May 2006;88(5):925-935. PMID 16651565.
13. Peters CL, Erickson JA. Treatment of femoro-acetabular impingement with surgical dislocation and debridement in young adults. *J Bone Joint Surg Am*. Aug 2006;88(8):1735-1741. PMID 16882895.
14. Kierkegaard S, Langeskov-Christensen M, Lund B, et al. Pain, activities of daily living and sport function at different time points after hip arthroscopy in patients with femoroacetabular impingement: a systematic review with meta-analysis. *Br J Sports Med*. Apr 2017;51(7):572-579. PMID 27845683.
15. Krych AJ, Thompson M, Knutson Z, et al. Arthroscopic labral repair versus selective labral debridement in female patients with femoroacetabular impingement: a prospective randomized study. *Arthroscopy*. Jan 2013;29(1):46-53. PMID 23276413.
16. Minkara AA, Westermann RW, Rosneck J, et al. Systematic review and meta-analysis of outcomes after hip arthroscopy in femoroacetabular impingement. *Am J Sports Med*. Jan 1 2018;363546517749475. PMID 29373805.
17. Griffin DR, Dickenson EJ, Wall PDH, et al. Hip arthroscopy versus best conservative care for the treatment of femoroacetabular impingement syndrome (UK FASHIoN): a multicenter randomised controlled trial. *Lancet*. 2018 Jun 2;391(10136):2225-2235. PMID: 29893223.
18. Palmer AJR, Ayyar Gupta V, Fernquest S, et al. *BMJ*. 2019 Feb 7;364:l185. PMID: 30733197.
19. Malviya A, Stafford GH, Villar RN. Impact of arthroscopy of the hip for femoroacetabular impingement on quality of life at a mean follow-up of 3.2 years. *J Bone Joint Surg Br*. Apr 2012;94(4):466-470. PMID 22434460.
20. Philippon MJ, Schroder ESBG, Briggs KK. Hip arthroscopy for femoroacetabular impingement in patients aged 50 years or older. *Arthroscopy*. Jan 2012;28(1):59-65. PMID 21982390.
21. Palmer DH, Ganesh V, Comfort T, et al. Midterm outcomes in patients with cam femoroacetabular impingement treated arthroscopically. *Arthroscopy*. Nov 2012;28(11):1671-1681. PMID 22959221.
22. Javed A, O'Donnell JM. Arthroscopic femoral osteochondroplasty for cam femoroacetabular impingement in patients over 60 years of age. *J Bone Joint Surg Br*. Mar 2011;93(3):326-331. PMID 21357953.
23. Larson CM, Giveans MR, Taylor M. Does arthroscopic FAI correction improve function with radiographic arthritis? *Clin Orthop Relat Res*. Jun 2011;469(6):1667-1676. PMID 21181460.
24. Horisberger M, Brunner A, Herzog RF. Arthroscopic treatment of femoral acetabular impingement in patients with preoperative generalized degenerative changes. *Arthroscopy*. May 2010;26(5):623-629. PMID 20434659.
25. Byrd JW, Jones KS. Arthroscopic femoroplasty in the management of cam-type femoroacetabular impingement. *Clin Orthop Relat Res*. Mar 2009;467(3):739-746. PMID 19096902.
26. Larson CM, Giveans MR. Arthroscopic management of femoroacetabular impingement: early outcomes measures. *Arthroscopy*. May 2008;24(5):540-546. PMID 18442686.
27. Zhang D, Chen L, Wang G. Hip arthroscopy versus open surgical dislocation for femoroacetabular impingement: A systematic review and meta-analysis. *Medicine (Baltimore)*. Oct 2016;95(41):e5122. PMID 27741133.
28. Nwachukwu BU, Rebolledo BJ, McCormick F, et al. Arthroscopic versus open treatment of femoroacetabular impingement: a systematic review of medium- to long-term outcomes. *Am J Sports Med*. Apr 2016;44(4):1062-1068. PMID 26059179.
29. Matsuda DK, Carlisle JC, Arthurs SC, et al. Comparative systematic review of the open dislocation, mini-open, and arthroscopic surgeries for femoroacetabular impingement. *Arthroscopy*. Feb 2011;27(2):252-269. PMID 21266276.
30. Botser IB, Smith TW, Jr., Nasser R, et al. Open surgical dislocation versus arthroscopy for femoroacetabular impingement: a comparison of clinical outcomes. *Arthroscopy*. Feb 2011;27(2):270-278. PMID 21266277.
31. Papalia R, Del Buono A, Franceschi F, et al. Femoroacetabular impingement syndrome management: arthroscopy or open surgery? *Int Orthop*. May 2012;36(5):903-914. PMID 22190060.
32. Zingg PO, Ulbrich EJ, Buehler TC, et al. Surgical hip dislocation versus hip arthroscopy for femoroacetabular impingement: clinical and morphological short-term results. *Arch Orthop Trauma Surg*. Jan 2013;133(1):69-79. PMID 23064993.
33. Domb BG, Stake CE, Botser IB, et al. Surgical dislocation of the hip versus arthroscopic treatment of femoroacetabular impingement: a prospective matched-pair study with average 2-year follow-up. *Arthroscopy*. Sep 2013;29(9):1506-1513. PMID 23992988.

34. Chiron P, Espie A, Reina N, et al. Surgery for femoroacetabular impingement using a minimally invasive anterolateral approach: analysis of 118 cases at 2.2-year follow-up. *Orthop Traumatol Surg Res.* Feb 2012;98(1):30-38. PMID 22257764.
35. Laude F, Stimesi E, Nogier A. Femoroacetabular impingement treatment using arthroscopy and anterior approach. *Clin Orthop Relat Res.* Mar 2009;467(3):747-752. PMID 19089524.
36. de Sa D, Cargnelli S, Catapano M, et al. Femoroacetabular impingement in skeletally immature patients: a systematic review examining indications, outcomes, and complications of open and arthroscopic treatment. *Arthroscopy.* Feb 2015;31(2):373-384. PMID 25262968.
37. Tran P, Pritchard M, O'Donnell J. Outcome of arthroscopic treatment for cam type femoroacetabular impingement in adolescents. *ANZ J Surg.* Sep 3 2013;83(5):382-386. PMID 22943465.
38. Guindani N, Eberhardt O, Wirth T, et al. Surgical dislocation for pediatric and adolescent hip deformity: clinical and radiographical results at 3 years follow-up. *Arch Orthop Trauma Surg.* Apr 2017;137(4):471-479. PMID 28197752.
39. Nwachukwu BU, Chang B, Kahlenberg CA, et al. Arthroscopic treatment of femoroacetabular impingement in adolescents provides clinically significant outcome improvement. *Arthroscopy.* Oct 2017;33(10):1812-1818. PMID 28623078.
40. Sink EL, Zaltz I, Heare T, et al. Acetabular cartilage and labral damage observed during surgical hip dislocation for stable slipped capital femoral epiphysis. *J Pediatr Orthop.* Jan-Feb 2010;30(1):26-30. PMID 20032738.
41. Ziebarth K, Zilkens C, Spencer S, et al. Capital realignment for moderate and severe SCFE using a modified Dunn procedure. *Clin Orthop Relat Res.* Mar 2009;467(3):704-716. PMID 19142692.
42. Spencer S, Millis MB, Kim YJ. Early results of treatment of hip impingement syndrome in slipped capital femoral epiphysis and pistol grip deformity of the femoral head-neck junction using the surgical dislocation technique. *J Pediatr Orthop.* May-Jun 2006;26(3):281-285. PMID 16670535.
43. Sardana V, Philippon MJ, de Sa D, et al. Revision hip arthroscopy indications and outcomes: a systematic review. *Arthroscopy.* Oct 2015;31(10):2047-2055. PMID 26033461.
44. Cvetanovich GL, Harris JD, Erickson BJ, et al. Revision hip arthroscopy: a systematic review of diagnoses, operative findings, and outcomes. *Arthroscopy.* Jul 2015;31(7):1382-1390. PMID 25703289.
45. Newman JT, Briggs KK, McNamara SC, et al. Outcomes after revision hip arthroscopic surgery in adolescent patients compared with a matched cohort undergoing primary arthroscopic surgery. *Am J Sports Med.* Dec 2016;44(12):3063-3069. PMID 27514736.
46. Gwathmey FW, Jones KS, Thomas Byrd JW. Revision hip arthroscopy: findings and outcomes. *J Hip Preserv Surg.* Dec 2017;4(4):318-323. PMID 29250340.
47. National Institute for Health and Clinical Excellence (NICE). Arthroscopic femoro-acetabular surgery for hip impingement syndrome [IPG408]. 2011; <https://www.nice.org.uk/guidance/ipg408>. Accessed March 22, 2018.
48. National Institute for Health and Care Excellence (NICE). Open femoro-acetabular surgery for hip impingement syndrome [IPG403]. 2011; <https://www.nice.org.uk/guidance/ipg403>. Accessed March 22, 2018.