

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

Medical Policy

Electrical Stimulation Devices for Psychiatric and Neurological Conditions

Table of Contents

- Policy: Commercial
- Policy: Medicare
- Authorization Information
- Coding Information
- Description
- Policy History
- Information Pertaining to All Policies
- References
- Endnotes

Policy Number: 157

BCBSA Reference Number: N/A

NCD/LCD: NA

Related Policies

- Cranial Electrotherapy Stimulation CES and Auricular Electrostimulation, #362
- Outpatient Electroconvulsive Therapy, #319
- Percutaneous Electrical nerve Stimulation (PENS) and Percutaneous Neuromodulation Therapy (PNT), #172
- Transcranial Magnetic Stimulation as a Treatment of Depression and Other Psychiatric/Neurologic Disorders, #297
- Transcutaneous Electrical Nerve Stimulation (TENS), #003

Policy¹

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO BlueSM and Medicare PPO BlueSM Members

Transcranial direct current stimulation devices (TDCS) are considered INVESTIGATIONAL for the treatment of all conditions including but not limited to brain and spinal cord injury, Parkinson's disease and behavioral health conditions such as major depressive disorder, anxiety, substance use disorders, and post-traumatic stress disorder (PTSD).

Transcranial alternating current stimulation (TACS) devices are considered <u>INVESTIGATIONAL</u> for the treatment of all conditions including Parkinson's disease, Alzheimer's, Seizure disorders, and behavioral health conditions such as major depressive disorder, anxiety, substance use disorders, and post-traumatic stress disorder (PTSD).

INVESTIGATIONAL devices include but are not limited to:

- The Brain Stimulator v3.0
- The Brain Driver v2.1
- APEX Type A
- OMNI Stimulator

- TCT TDCS Device
- Soterix 1x1
- Focus V2 (TES)
- Focus V3
- NeuroConn DC Stimulator PLUS
- Activa Dose II Caputron
- Foc.us V2.

Note: BCBSMA does not approve any medical devices that have not received FDA approval.

Prior Authorization Information

Inpatient

• For services described in this policy, precertification/preauthorization <u>IS REQUIRED</u> for all products if the procedure is performed **inpatient**.

Outpatient

For services described in this policy, see below for products where prior authorization <u>might be</u> <u>required</u> if the procedure is performed <u>outpatient</u>.

Outpatient	
Commercial Managed Care (HMO and POS)	This is not a covered service.
Commercial PPO and Indemnity	This is not a covered service.
ledicare HMO BlueSM This is not a covered service.	
Medicare PPO Blue SM	This is not a covered service.

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

CPT Codes

There is no specific CPT code for this service.

Description

Transcranial direct current stimulation (TDCS) devices neurostimulation to specifically targeted areas of the brain through low levels of constant electric currents. TDCS devices deliver cathodal (negative) stimulation or anodal (positive) stimulation via small electrodes placed on different regions of the scalp and are thought to influence the rate of neuronal activity.

Transcranial alternating current stimulation (TACS) devices deliver neurostimulation to specifically targeted areas of the brain with alternating electoral currents. The alternating currents are thought to change the rate of abnormal neuronal activity that may be related to a variety of behavioral and cognitive conditions.

TDCS and TACS are non-invasive, painless treatments that are being evaluated for a variety of conditions including, chronic pain, brain and spinal cord injury, Parkinson's disease, and psychiatric conditions such as major depressive disorder, substance use disorders and anxiety. TDCS and TACS devices can be used in the home setting or in a therapeutic setting under the direction of a physician. TDCS and TACS devices can be manually set to the desired electrical current and length of time. Recommended time for treatment is 20-30 minutes of neurostimulation per session. Results from the treatments depend on how long the neurostimulation is delivered.

Some of the devices used for TDCS include The Brain Stimulator v3.0tm, the APEX type Atm, and the Omni Stimulator. Other devices include the Activa Dose II, The Brain Driver v2.1, and the NeuroConn. Some of the devices come with software that help improve reproducible neurostimulation levels and headbands that hold the electrodes in place during treatment. TACS devices include Soterix 1x1, Starstim, and the Brain Stimulator tm.

Summary

Transcranial Direct Current Stimulation (TDCS) and Transcranial Alternating Current Stimulation devices are being investigated for a variety of conditions including chronic pain, brain and spinal cord injury, Parkinson's disease and psychiatric conditions including major depressive disorder, anxiety, and post-traumatic stress disorder.

Small randomized controlled trials and multiple case studies have evaluated the use of TDCS and TACS devices in Parkinson's and Alzheimer's disease, major depressive disorder, anxiety and post traumatic stress disorder (PTSD). While some of the studies showed small improvements in symptom reduction and increased mood, outcomes did not demonstrate clinically significant improvements in symptoms. The study designs and treatment protocols differed significantly and were not reproducible. The evidence is insufficient to determine the effects of this technology on health outcomes.

Policy History

· • · · · · · · · · · · · · · · · · · ·	oj meterj					
Date	Action					
4/2019	New medical policy describing investigational indications. Effective 4/1/2019.					

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

Medical Policy Terms of Use

Managed Care Guidelines

Indemnity/PPO Guidelines

Clinical Exception Process

Medical Technology Assessment Guidelines

References

- 1. Fregni, F., Nitsche, M.A., Loo, C. K. Regulatory Considerations for the Clinical and Research Use of Transcranial Direct Current Stimulation (tDCS): review and recommendations from an expert panel. Clin Res Regul Aff. 2015 Mar 1; 32(1): 22–35.
- 2. Alonzo, A., Aaronson, S., Bikson, M. Study design and methodology for a multicentre, randomised controlled trial of transcranial direct current stimulation as a treatment for unipolar and bipolar depression.
- 3. Dell'Osso, B., Dobrea, C., Arici, C. Augmentative transcranial direct current stimulation (tDCS) in poor responder depressed patients: a follow-up study. CNS Spectr. 2014 Aug;19(4):347-54. doi: 10.1017/S1092852913000497. Epub 2013 Aug 20.
- 4. Dondé, C., Amad, A., Nieto, I. Transcranial direct-current stimulation (tDCS) for bipolar depression: A systematic review and meta-analysis. Prog Neuropsychopharmacol Biol Psychiatry. 2017 Aug 1;78:123-131. doi: 10.1016/j.pnpbp.2017.05.021. Epub 2017 May 25.
- 5. Benninger, D., Lomarev, M., Lopez, G. Transcranial Direct Current Stimulation for the Treatment of Parkinson's Disease. Journal of Neurosurgical Psychiatry, 2010 October, 81(10):1105-1111.
- 6. Loo, C.K., Husain, M.M., McDonald, W.M. International randomized-controlled trial of transcranial Direct Current Stimulation in depression. Brain Stimul. 2018 Jan Feb;11(1):125-133. doi: 10.1016/j.brs.2017.10.011. Epub 2017 Oct 27.
- 7. Sampaio-Junior, B., Tortella, G., Borrione, L. Efficacy and Safety of Transcranial Direct Current Stimulation as an Add-on Treatment for Bipolar Depression. A Randomized Clinical Trial. *JAMA Psychiatry*. February 2018. 75(2) 158-166. doi:10.1001/jamapsychiatry.2017.4040
- 8. Brunoni, A.R., Moffa, A.H., Fregni, F. Transcranial direct current stimulation for acute major depressive episodes: meta-analysis of individual patient data. <u>Br J Psychiatry.</u> 2016 Jun;208(6):522-31. doi: 10.1192/bjp.bp.115.164715. Epub 2016 Apr 7.

- 9. Shiozawa, P., Fregni, F., Benseñor, I.M. Transcranial direct current stimulation for major depression: an updated systematic review and meta-analysis. Int J Neuropsychopharmacol. 2014 Sep;17(9):1539.
- 10. Sheng, J., Li, B. Recent Advances in Non-invasive Brain Stimulation for Major Depressive Disorder. Front Hum Neurosci. 2017 Nov 6;11:526. doi: 10.3389/fnhum.2017.00526. eCollection 2017.
- 11. Meron, D., Hedger, N., Garner, M. Transcranial direct current stimulation (tDCS) in the treatment of depression: Systematic review and meta-analysis of efficacy and tolerability. Neurosci Biobehav Rev. 2015 Oct;57:46-62. doi: 10.1016/j.neubiorev.2015.07.012. Epub 2015 Jul 29.
- 12. Van den Eynde, F., Daskalakis, Z.J. Clinical utility of transcranial direct current stimulation (tDCS) for treating major depression: a systematic review and meta-analysis of randomized, double-blind and sham-controlled trials. <u>J Psychiatr Res.</u> 2013 Jan;47(1):1-7. doi: 10.1016/j.jpsychires.2012.09.025. Epub 2012 Oct 22.
- 13. Al-Kaysi, A.M., Al-Ani, A., Loo, C.K. Predicting brain stimulation treatment outcomes of depressed patients through the classification of EEG oscillations. <u>Conf Proc IEEE Eng Med Biol Soc.</u> 2016 Aug;2016:5266-5269. doi: 10.1109/EMBC.2016.7591915.
- Mondino M¹, Bennabi D, Poulet E. Can transcranial direct current stimulation (tDCS) alleviate symptoms and improve cognition in psychiatric disorders? World J Biol Psychiatry. 2014 May;15(4):261-75. doi: 10.3109/15622975.2013.876514. Epub 2014 Jan 21.
- 15. Kekic, M., Boysen, E., Campbell, I.C. A systematic review of the clinical efficacy of transcranial direct current stimulation (tDCS) in psychiatric disorders. J Psychiatr Res. 2016 Mar;74:70-86. doi: 10.1016/j.jpsychires.2015.12.018. Epub 2015 Dec 22.
- Brunoni, A.R., Moffa, A.H., Sampaio-Junior, B. Trial of Electrical Direct-Current Therapy versus Escitalopram for Depression. N Engl J Med. 2017 Jun 29;376(26):2523-2533. doi: 10.1056/NEJMoa1612999
- 17. Brunoni, A.R., Valiengo, L., Baccaro, A. The sertraline vs. electrical current therapy for treating depression clinical study: results from a factorial, randomized, controlled trial. JAMA Psychiatry. 2013 Apr;70(4):383-91. doi: 10.1001/2013.jamapsychiatry.32.
- 18. Martin, D.M., Alonzo, A., Mitchell, P.B. Fronto-extracephalic transcranial direct current stimulation as a treatment for major depression: an open-label pilot study. J Affect Disord. 2011 Nov;134(1-3):459-63. doi: 10.1016/j.jad.2011.05.018.
- 19. Philip, P., Nelson, B.G., Frohlich, F. Low-Intensity Transcranial Current Stimulation in Psychiatry. American Journal of Psychiatry July 2017. 174(7). 628-639.
- 20. Szymkowicz, S., McLaren, M., Suryadevara, U. Transcranial Direct Current Stimulation Use in the Treatment of Neuropsychiatric Disorders: A Brief Review. <u>Psychiatr Ann. 2016 Nov; 46(11): 642–646.</u>
- 21. Murphy, D.N., Boggio, P., Fregni, F. Transcranial direct current stimulation as a therapeutic tool for the treatment of major depression: insights from past and recent clinical studies. Curr Opin Psychiatry. 2009 May;22(3):306-11. doi: 10.1097/YCO.0b013e32832a133f.
- 22. Vigod, S., Dennis, C.L., Daskalakis, Z. Transcranial direct current stimulation (tDCS) for treatment of major depression during pregnancy: study protocol for a pilot randomized controlled trial. Trials. 2014 Sep 18;15:366. doi: 10.1186/1745-6215-15-366.
- 23. Lefaucheur, J.P., Antal, A., Ayache, S.S. Evidence-based guidelines on the therapeutic use of transcranial direct current stimulation (tDCS). <u>Clin Neurophysiol.</u> 2017 Jan;128(1):56-92. doi: 10.1016/j.clinph.2016.10.087. Epub 2016 Oct 29.
- 24. Tavakoli, A. V., Yun, K. Transcranial Alternating Current Stimulation (tACS) Mechanisms and Protocols. Front Cell Neurosci. 2017. 11:214. doi: 10.3389/fncel.2017.00214
- 25. Zaehle, T., Rach, S., Herrmann, C.S. Transcranial alternating current stimulation enhances individual alpha activity in human EEG. PloS One. 2010 Nov. 5(11) doi: 10.1371/journal.pone.0013766.

M		

¹ Based on expert opinion