



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

Medical Policy

Absorbable Nasal Implant for Treatment of Nasal Valve Collapse

Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

Policy Number: 186

BCBSA Reference Number: 7.01.163

NCD/LCD: N/A

Related Policies

None

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO BlueSM and Medicare PPO BlueSM Members

The insertion of an absorbable lateral nasal implant for the treatment of symptomatic nasal valve collapse is considered INVESTIGATIONAL.

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed inpatient.

Outpatient

- For services described in this policy, see below for products where prior authorization might be required if the procedure is performed outpatient.

	Outpatient
Commercial Managed Care (HMO and POS)	This is not a covered service.
Commercial PPO and Indemnity	This is not a covered service.
Medicare HMO BlueSM	This is not a covered service.
Medicare PPO BlueSM	This is not a covered service.

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

The following HCPCS code is considered investigational for the conditions listed for Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:

HCPCS Codes

HCPCS codes:	Code Description
C9749	Repair of nasal vestibular lateral wall stenosis with implant(s)

Description

NASAL OBSTRUCTION

Nasal obstruction is defined clinically as a patient symptom that presents as a sensation of reduced or insufficient airflow through the nose. Commonly, patients will feel that they have nasal congestion or stuffiness. In adults, clinicians focus the evaluation of important features of the history provided by the patient such as whether symptoms are unilateral or bilateral. Unilateral symptoms are more suggestive of structural causes of nasal obstruction. A history of trauma or previous nasal surgery, especially septoplasty or rhinoplasty, is also important. Diurnal or seasonal variation in symptoms is associated with allergic conditions.

Etiology

Nasal obstruction associated with the external nasal valve is commonly associated with post-rhinoplasty or traumatic sequelae and may require functional rhinoplasty procedures. A common cause of internal nasal valve collapse is septal deviation. Prior nasal surgery, nasal trauma, and congenital anomaly are additional causes.

Pathophysiology

The internal nasal valve, bordered by the collapsible soft tissue between the upper and lower lateral cartilages, anterior end of the inferior turbinate, and the nasal septum, forms the narrowest part of the nasal airway. During inspiration, the lateral wall cartilage is dynamic and draws inward toward the septum and the internal nasal valve narrows providing protection to the upper airways. The angle at the junction between the septum and upper lateral cartilage is normally 10° to 15° in white populations. Given that the internal nasal valve accounts for at least half of the nasal airway resistance; even minor further narrowing of this area can lead to symptomatic obstruction for a patient. Damaged or weakened lateral nasal cartilage will further decrease airway capacity of the internal nasal valve area, increasing airflow resistance and symptoms of congestion.¹

Physical Examination

A thorough physical examination of the nose, nasal cavity, and the nasopharynx is generally sufficient to identify the most likely etiology for the nasal obstruction. Both the external and internal nasal valve areas should be examined. The external nasal valve is at the level of the internal nostril. It is formed by the caudal portion of the lower lateral cartilage, surrounding soft tissue and the membranous septum. The Cottle maneuver is an examination in which the cheek on the symptomatic side is gently pulled laterally with 1 to 2 fingers. If the patient is less symptomatic with inspiration during the maneuver, the assumption is that the nasal valve has been widened from a collapsed state or dynamic nasal valve collapse. An individual can perform the maneuver on oneself and it is subjective. A clinician performs the modified Cottle maneuver. A cotton swab or curette is inserted into the nasal cavity to support the nasal cartilage and the patient reports whether there is an improvement in the symptoms with inspiration. In both instances, a change in the external contour of the lateral nose may be apparent to both the patient and the examiner.

Measuring Nasal Obstruction

Stewart et al (2004) proposed the Nasal Obstruction Symptom Evaluation as a validated sinonasal-specific health status instrument that is used to assess the impact of nasal obstruction on the quality of life of affected persons.² It is a 5-item questionnaire on breathing problems: nasal congestion or stuffiness, nasal blockage or obstruction, trouble breathing through the nose, trouble sleeping, and inability to get enough air through the nose during exercise or exertion. The responses are made on a Likert-type scale ranging from 0 (not a problem) to 4 (severe problem). The range of raw scores is 0 to 20. The score is then scaled to a potential total score of 0 to 100 by multiplying the raw score by 5. A score of 100 means the worst possible problem with nasal obstruction.

Lipan and Most (2013) developed a Nasal Obstruction Symptom Evaluation scale-based nasal obstruction severity classification system.³ The system is proposed as a means to classify patients for clinical management as well as to better define study populations and describe treatment or intervention responses (see Table 1).

Table 1. NOSE Severity Classification

Severity Class	NOSE Score Range
Mild	5-25
Moderate	30-50
Severe	55-75
Extreme	80-100

NOSE: Nasal Obstruction Symptom Evaluation.

Treatment

Treatment of symptomatic nasal valve collapse includes the use of nonsurgical interventions such as the adhesive strips applied externally across the nose (applying the principle of the Cottle maneuver) or use of nasal dilators, cones, or other devices that support the lateral nasal wall internally (applying the principle of the modified Cottle maneuver).

Severe cases of obstruction result from nasal valve deformities are treated with surgical grafting to widen and/or strengthen the valve. Common materials include cartilaginous autografts and allografts, as well as permanent synthetic grafts. Cartilage grafts are most commonly harvested from the patient's nasal septum or ear.

Nasal Implants

The placement of an absorbable implant to support the lateral nasal cartilages has been proposed as an alternative to more invasive grafting procedures in patients with severe nasal obstruction.

Summary

Nasal valve collapse is a readily identifiable cause of nasal obstruction. Specifically, the internal nasal valve represents the narrowest portion of the nasal airway with the upper lateral nasal cartilages present as supporting structures. The external nasal valve is an area of potential dynamic collapse that is supported by the lower lateral cartilages. Damaged or weakened cartilage will further decrease airway capacity and increase airflow resistance and may be associated with symptoms of obstruction. Patients with nasal valve collapse may be treated with nonsurgical interventions in an attempt to increase the airway capacity but severe symptoms and anatomic distortion are treated with surgical cartilage graft procedures. The placement of an absorbable implant to support the lateral nasal cartilages has been proposed as an alternative to more invasive grafting procedures in patients with severe nasal obstruction. The concept is that the implant may provide support to the lateral nasal wall prior to resorption and then stiffen the wall with scarring as it is resorbed.

For individuals with symptomatic nasal obstruction due to internal nasal valve collapse who receive an absorbable lateral nasal valve implant, the evidence includes 2 nonrandomized prospective, single-cohort industry-sponsored studies. Relevant outcomes are symptoms, change in disease status, treatment-related morbidity, functional outcomes, and quality of life. Both studies are limited by the heterogeneity of

the populations evaluated. Specifically, the types and rates of prior nasal procedures were not well described, nor was the clinical rationale for alternative or adjunctive procedural interventions. Overall, improvements in the Nasal Obstruction Symptom Evaluation score have been demonstrated in the study reports. However, a clinically significant difference may not be consistently apparent in small study populations. Some patients meeting the positive responder criteria still reported severe symptoms, and many patients reported some loss of improvement at 1 year. Data elements are missing or difficult to determine for important outcomes. As reported, adverse events appeared to be mild in severity and self-limiting, but still appeared common. Device retrievals are incompletely characterized. They occurred in 10% of patients in the primary cohort study, and it is not known, eg, whether a device retrieval occurred in a patient who had only a unilateral nasal implant. The need for device retrievals appears to occur early in the course of follow-up (1 month); suggesting technical experience limitations on the part of the operator or inappropriate patient selection. The duration of outcomes reporting is less than the duration of absorption of the device (18 months) and the purported completion of tissue remodeling phase (24 months). Randomized controlled trials with a sham control are feasible and should be performed. The evidence is insufficient to determine the effects of the technology on health outcomes.

Policy History

Date	Action
3/2019	New medical policy describing investigational indications. Effective 3/1/2019.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

References

1. Howard BK, Rohrich RJ. Understanding the nasal airway: principles and practice. *Plast Reconstr Surg*. Mar 2002;109(3):1128-1146; quiz 1145-1146. PMID 11884847
2. Stewart MG, Witsell DL, Smith TL, et al. Development and validation of the Nasal Obstruction Symptom Evaluation (NOSE) scale. *Otolaryngol Head Neck Surg*. Feb 2004;130(2):157-163. PMID 14990910
3. Lipan MJ, Most SP. Development of a severity classification system for subjective nasal obstruction. *JAMA Facial Plast Surg*. Sep-Oct 2013;15(5):358-361. PMID 23846399
4. Rhee JS, Weaver EM, Park SS, et al. Clinical consensus statement: Diagnosis and management of nasal valve compromise. *Otolaryngol Head Neck Surg*. Jul 2010;143(1):48-59. PMID 20620619
5. Fraser L, Kelly G. An evidence-based approach to the management of the adult with nasal obstruction. *Clin Otolaryngol*. Apr 2009;34(2):151-155. PMID 19413614
6. Spirox. Latera. 2017; <http://www.spiroxmed.com/latera>. Accessed September 10, 2018.
7. Stolovitzky P, Sidle DM, Ow RA, et al. A prospective study for treatment of nasal valve collapse due to lateral wall insufficiency: Outcomes using a bioabsorbable implant. *Laryngoscope*. May 14 2018. PMID 29756407
8. San Nicoló M, Stelter K, Sadick H, et al. Absorbable implant to treat nasal valve collapse. *Facial Plast Surg*. Apr 2017;33(2):233-240. PMID 28388804