



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an independent
Licensee of the Blue Cross and Blue Shield Association

Medical Policy

Magnetic Resonance–Guided Focused Ultrasound

Table of Contents

- [Policy: Commercial](#)
- [Coding Information](#)
- [Information Pertaining to All Policies](#)
- [Policy: Medicare](#)
- [Description](#)
- [References](#)
- [Authorization Information](#)
- [Policy History](#)

Policy Number: 243

BCBSA Reference Number: 7.01.109

NCD/LCD: Magnetic Resonance Image Guided High Intensity Focused Ultrasound (MRgFUS) for Essential Tremor (L37421)

Related Policies

- Occlusion of Uterine Arteries Using Transcatheter Embolization #[242](#)
- Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors #[259](#)

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity

Magnetic resonance guided high-intensity ultrasound ablation may be considered **MEDICALLY NECESSARY** for pain palliation in adult patients with metastatic bone cancer who failed or are not candidates for radiotherapy.

Magnetic resonance-guided high-intensity ultrasound ablation may be considered **MEDICALLY NECESSARY** for the treatment of medicine-refractory essential tremors.

Magnetic resonance-guided high-intensity ultrasound ablation is considered **INVESTIGATIONAL** in all other situations including but not limited to:

- Treatment of uterine fibroids; and
- Treatment of other tumors (eg, brain cancer, prostate cancer, and breast cancer, desmoid).

Medicare HMO BlueSM and Medicare PPO BlueSM Members

Medical necessity criteria and coding guidance for **Medicare Advantage members living in Massachusetts** can be found through the link below.

[Local Coverage Determinations \(LCDs\) for National Government Services, Inc.](#)

Local Coverage Determination (LCD): Magnetic Resonance Image Guided High Intensity Focused Ultrasound (MRgFUS) for Essential Tremor (L37421)

Note: To review the specific LCD, please remember to click “accept” on the CMS licensing agreement at the bottom of the CMS webpage.

For medical necessity criteria and coding guidance for **Medicare Advantage members living outside of Massachusetts**, please see the Centers for Medicare and Medicaid Services website at <https://www.cms.gov> for information regarding your specific jurisdiction.

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	Prior authorization is not required .
Commercial PPO and Indemnity	Prior authorization is not required .
Medicare HMO Blue SM	Prior authorization is not required .
Medicare PPO Blue SM	Prior authorization is not required .

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member’s contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

The above **medical necessity criteria MUST** be met for the following codes to be covered for Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:

CPT Codes

CPT codes:	Code Description
0398T	Magnetic resonance image guided high intensity focused ultrasound (mrgfus), stereotactic ablation lesion, intracranial for movement disorder including stereotactic navigation and frame placement when performed

The following ICD Diagnosis Code is considered medically necessary when submitted with the CPT code above if **medical necessity criteria** are met:

ICD-10 diagnosis coding

ICD-10 Diagnosis code	Description
G25.0	Essential tremor

The following CPT codes are considered investigational for **Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:**

CPT Codes

CPT codes:	Code Description
0071T	Focused ultrasound ablation of uterine leiomyomata, including MR guidance; total leiomyomata volume less than 200 cc of tissue
0072T	Focused ultrasound ablation of uterine leiomyomata, including MR guidance; total leiomyomata volume greater or equal to 200 cc of tissue

Description

UTERINE FIBROIDS

Uterine fibroids are one of the most common conditions affecting women in the reproductive years. Symptoms of uterine fibroids include menorrhagia, pelvic pressure, or pain.

Treatment

Several approaches currently available to treat symptomatic uterine fibroids include hysterectomy, abdominal myomectomy, laparoscopic and hysteroscopic myomectomy, hormone therapy, uterine artery embolization, and watchful waiting. Hysterectomy and various myomectomy procedures are considered the criterion standard treatments.

METASTATIC BONE DISEASE

Metastatic bone disease is one of the most common causes of cancer pain.

Treatment

Existing treatments include conservative measures (eg, massage, exercise) and pharmacologic agents (eg, analgesics, bisphosphonates, corticosteroids). For patients who do not respond to these treatments, standard care is external-beam radiotherapy. However, a substantial proportion of patients have residual pain after radiotherapy, and there is a need for alternative treatments for these patients. (One option, radiofrequency ablation, is addressed in related policy #259).

ESSENTIAL TREMORS

Essential tremor (ET) is the most common movement disorder, with an estimated prevalence of 5% worldwide. ET most often affects the hands and arms, may affect head and voice, and rarely includes the face, legs, and trunk. ET is heterogeneous among patients, varying in frequency, amplitude, causes of exacerbation, and association with other neurologic deficits.

Treatment

The neuropathology of ET is uncertain, with some evidence suggesting that ET is localized in the brainstem and cerebellum. If patients with ET experience intermittent or persistent disability due to the tremors, initial therapy is with drugs (β -blockers or anticonvulsants). For medicine-refractory patients, surgery (deep brain stimulation or thalamotomy) may be offered, though high rates of adverse events have been observed.

MAGNETIC RESONANCE-GUIDED FOCUSED ULTRASOUND

Magnetic resonance-guided focused ultrasound (MRgFUS) is a noninvasive treatment that combines 2 technologies: focused ultrasound and magnetic resonance imaging (MRI). The ultrasound beam penetrates through the soft tissues and, using MRI for guidance and monitoring, the beam can be focused on targeted sites. Ultrasound causes a local increase in temperature in the target tissue, resulting in coagulation necrosis while sparing the surrounding normal structures. Ultrasound waves from each sonication are directed at a focal point that has a maximum focal volume of 20 nm in diameter and 15 nm in height/length. This causes a rapid rise in temperature (ie, to 65°C-85°C), which is sufficient to ablate tissue at the focal point. In addition to providing guidance, the associated MRI can provide online thermometric imaging, a temperature “map”, to confirm the therapeutic effect of the ablation treatment and allow for real-time adjustment of the treatment parameters.

The U.S. Food and Drug Administration (FDA) approved the ExAblate MRgFUS system (InSightec) for 2 indications: treatment of uterine fibroids (leiomyomata) and palliation of pain associated with tumors metastatic to bone. The ultrasound equipment is specifically designed to be compatible with magnetic resonance magnets, and it is integrated into standard clinical MRI units; it also includes a patient table, which has a cradle that houses the focused ultrasound transducer in water or a light oil bath. Some models have a detachable cradle; only certain cradle types can be used for palliation of pain associated with metastatic bone cancer. For treating pain associated with bone metastases, the aim of MRgFUS is to destroy nerves in the bone surface surrounding the tumor.

MRgFUS is also being investigated for the treatment of other tumors, including breast, prostate, brain, and desmoid tumors as well as nonspinal osteoid osteoma.

Summary

An integrated system providing magnetic resonance–guided focused ultrasound (MRgFUS) treatment is proposed as a noninvasive therapy for uterine fibroids and pain palliation of bone metastases. MRgFUS is also being investigated as a treatment of other benign and malignant tumors as well as essential tremors.

For individuals who have uterine fibroids who receive MRgFUS, the evidence includes 2 small randomized controlled trials (RCTs), nonrandomized comparative studies, and case series. Relevant outcomes are symptoms, quality of life, resource utilization, and treatment-related morbidity. One RCT (N=20) has reported some health outcomes, but its primary purpose was to determine the feasibility of a larger trial. It did not find statistically significant differences in quality of life outcomes between active and sham treatment groups, but it did find lower fibroid volumes after active treatment. This trial did not have an active comparator, the clinical significance of the primary outcome was unclear, and there were no follow-up data beyond 1 year. The second RCT (N=49) is ongoing; preliminary results at 6 weeks posttreatment, comparing MRgFUS with uterine artery embolization have shown that the 2 groups are comparable in medication use and symptom improvement following treatments. Patients in the MRgFUS group reported recovering significantly faster than patients in the uterine artery embolization group, as measured by time to return to work and time to normal activities. In a separate 2013 comparative study, outcomes appeared to be better with uterine artery embolization than with MRgFUS. Long-term data on the treatment effects, recurrence rates, and impact on future fertility and pregnancy are lacking. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals with metastatic bone cancer who have failed or are not candidates for radiotherapy who receive MRgFUS, the evidence includes a sham-controlled randomized trial and several case series. Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, and treatment-related morbidity. The RCT found statistically significant improvements after MRgFUS in a composite outcome comprised of a reduction in pain and morphine use, and in pain reduction as a stand-alone outcome. A substantial proportion of patients in the treatment group experienced adverse events, but most events were transient and not severe. The case series reported reductions in pain following MRgFUS treatment, consistent with the RCT. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals with other tumors (eg, breast cancer, brain cancer, prostate cancer, desmoid, nonspinal osteoid osteoma) who receive MRgFUS, the evidence includes small case series. Relevant outcomes are symptoms, health status measures, and treatment-related morbidity. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals with medicine-refractory essential tremors who receive MRgFUS, the evidence includes 2 systematic reviews that identified an RCT and several observational studies. Relevant outcomes include symptoms, functional outcomes, quality of life, and treatment-related morbidity. The assessment did not pool study results but concluded that, overall, MRgFUS decreased tremor severity and improved quality of life. The sham-controlled randomized trial found significant improvements in the treatment group in tremor severity, functional improvement, and quality of life after 3 months of follow-up. The improvements

in hand tremor score, function, and quality of life were maintained at the 2-year follow-up. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

Policy History

Date	Action
2/2019	BCBSA National medical policy review. New medically necessary statement added for MRgFUS ablation for treatment of medicine-refractory essential tremors. Clarified coding information. Effective 2/1/2019.
4/2018	Local Coverage Determination (LCD): Magnetic Resonance Image Guided High Intensity Focused Ultrasound (MRgFUS) for Essential Tremor (L37421) added. Effective 4/1/2018.
8/2017	New references added from BCBSA National medical policy.
4/2016	BCBSA National medical policy review. Policy statements unchanged. Global change to policy to remove “imaging” (eg, title, policy statement) to standardize terminology to magnetic resonance–guided focused ultrasound (MRgFUS).
1/2016	Clarified coding information.
7/2015	BCBSA National medical policy review. New medically necessary indications described. Effective 7/1/2015.
5/2014	New references from BCBSA National medical policy.
8/2013	BCBSA National medical policy review. Policy changed to single investigational statement; no change to intent of policy. Policy title changed to MRI-Guided Focused Ultrasound (MRgFUS). Effective 8/1/2013.
11/2011-4/2012	Medical policy ICD 10 remediation: Formatting, editing and coding updates. No changes to policy statements.
9/2011	Reviewed - Medical Policy Group - Obstetrics and Gynecology. No changes to policy statements.
10/2010	Reviewed - Medical Policy Group - Obstetrics and Gynecology. No changes to policy statements.
7/2010	New policy describing on-going non-coverage. Effective 7/1/2010.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

References

1. Blue Cross Blue Shield Association Technology Evaluation Center (TEC). Magnetic resonance-guided focused ultrasound therapy for symptomatic uterine fibroids. *TEC Assessments*. 2005;Volume 20:Tab 10.
2. Barnard EP, AbdElmagied AM, Vaughan LE, et al. Periprocedural outcomes comparing fibroid embolization and focused ultrasound: a randomized controlled trial and comprehensive cohort analysis. *Am J Obstet Gynecol*. May 2017;216(5):500 e501-500 e511. PMID 28063909
3. Jacoby VL, Kohi MP, Poder L, et al. PROMISE trial: a pilot, randomized, placebo-controlled trial of magnetic resonance guided focused ultrasound for uterine fibroids. *Fertil Steril*. Mar 2016;105(3):773-780. PMID 26658133
4. Gizzo S, Saccardi C, Patrelli TS, et al. Magnetic resonance-guided focused ultrasound myomectomy: safety, efficacy, subsequent fertility and quality-of-life improvements, a systematic review. *Reprod Sci*. Apr 2014;21(4):465-476. PMID 23868442
5. Chen R, Keserci B, Bi H, et al. The safety and effectiveness of volumetric magnetic resonance-guided high-intensity focused ultrasound treatment of symptomatic uterine fibroids: early clinical experience in China. *J Ther Ultrasound*. Nov 2016;4:27. PMID 27822376

6. Froeling V, Meckelburg K, Schreiter NF, et al. Outcome of uterine artery embolization versus MR-guided high-intensity focused ultrasound treatment for uterine fibroids: long-term results. *Eur J Radiol.* Dec 2013;82(12):2265-2269. PMID 24075785
7. Rabinovici J, David M, Fukunishi H, et al. Pregnancy outcome after magnetic resonance-guided focused ultrasound surgery (MRgFUS) for conservative treatment of uterine fibroids. *Fertil Steril.* Jan 2010;93(1):199-209. PMID 19013566
8. Hurwitz MD, Ghanouni P, Kanaev SV, et al. Magnetic resonance-guided focused ultrasound for patients with painful bone metastases: phase III trial results. *J Natl Cancer Inst.* May 2014;106(5). PMID 24760791
9. Liberman B, Gianfelice D, Inbar Y, et al. Pain palliation in patients with bone metastases using MR-guided focused ultrasound surgery: a multicenter study. *Ann Surg Oncol.* Jan 2009;16(1):140-146. PMID 19002530
10. Arrigoni F, Barile A, Zugaro L, et al. Intra-articular benign bone lesions treated with magnetic resonance-guided focused ultrasound (MRgFUS): imaging follow-up and clinical results. *Med Oncol.* Apr 2017;34(4):55. PMID 28244018
11. Zippel DB, Papa MZ. The use of MR imaging guided focused ultrasound in breast cancer patients; a preliminary phase one study and review. *Breast Cancer.* 2005;12(1):32-38. PMID 15657521
12. Hynynen K, Pomeroy O, Smith DN, et al. MR imaging-guided focused ultrasound surgery of fibroadenomas in the breast: a feasibility study. *Radiology.* Apr 2001;219(1):176-185. PMID 11274554
13. Gianfelice D, Khiat A, Amara M, et al. MR imaging-guided focused US ablation of breast cancer: histopathologic assessment of effectiveness-- initial experience. *Radiology.* Jun 2003;227(3):849-855. PMID 12714680
14. Gianfelice D, Khiat A, Amara M, et al. MR imaging-guided focused ultrasound surgery of breast cancer: correlation of dynamic contrast-enhanced MRI with histopathologic findings. *Breast Cancer Res Treat.* Nov 2003;82(2):93-101. PMID 14692653
15. Merckel LG, Knuttel FM, Deckers R, et al. First clinical experience with a dedicated MRI-guided high-intensity focused ultrasound system for breast cancer ablation. *Eur Radiol.* Nov 2016;26(11):4037-4046. PMID 26852219
16. McDannold N, Clement GT, Black P, et al. Transcranial magnetic resonance imaging- guided focused ultrasound surgery of brain tumors: initial findings in 3 patients. *Neurosurgery.* Feb 2010;66(2):323-332; discussion 332. PMID 20087132
17. Napoli A, Anzidei M, De Nunzio C, et al. Real-time magnetic resonance-guided high-intensity focused ultrasound focal therapy for localised prostate cancer: preliminary experience. *Eur Urol.* Feb 2013;63(2):395-398. PMID 23159454
18. Geiger D, Napoli A, Conchiglia A, et al. MR-guided focused ultrasound (MRgFUS) ablation for the treatment of nonspinal osteoid osteoma: a prospective multicenter evaluation. *J Bone Joint Surg Am.* May 7 2014;96(9):743-751. PMID 24806011
19. Avedian RS, Bitton R, Gold G, et al. Is MR-guided high-intensity focused ultrasound a feasible treatment modality for desmoid tumors? *Clin Orthop Relat Res.* Mar 2016;474(3):697-704. PMID 26040967
20. Bucknor MD, Rieke V. MRgFUS for desmoid tumors within the thigh: early clinical experiences. *J Ther Ultrasound.* Feb 2017;5:4. PMID 28174660
21. Ghanouni P, Dobrotwir A, Bazzocchi A, et al. Magnetic resonance-guided focused ultrasound treatment of extra-abdominal desmoid tumors: a retrospective multicenter study. *Eur Radiol.* Feb 2017;27(2):732-740. PMID 27147222
22. Health Quality Ontario (HQO). Magnetic Resonance-Guided Focused Ultrasound Neurosurgery for Essential Tremor: A Health Technology Assessment. *Ont Health Technol Assess Ser.* May 2018;18(4):1-141. PMID 29805721
23. Mohammed N, Patra D, Nanda A. A meta-analysis of outcomes and complications of magnetic resonance-guided focused ultrasound in the treatment of essential tremor. *Neurosurg Focus.* Feb 2018;44(2):E4. PMID 29385917
24. Elias WJ, Lipsman N, Ondo WG, et al. A randomized trial of focused ultrasound thalamotomy for essential tremor. *N Engl J Med.* Aug 25 2016;375(8):730-739. PMID 27557301

25. Chang JW, Park CK, Lipsman N, et al. A prospective trial of magnetic resonance-guided focused ultrasound thalamotomy for essential tremor: Results at the 2-year follow-up. *Ann Neurol*. Jan 2018;83(1):107-114. PMID 29265546
26. Carranza-Mamane B, Havelock J, Hemmings R, et al. The management of uterine fibroids in women with otherwise unexplained infertility. *J Obstet Gynaecol Can*. Mar 2015;37(3):277-288. PMID 26001875
27. Lutz S, Balboni T, Jones J, et al. Palliative radiation therapy for bone metastases: Update of an ASTRO Evidence-Based Guideline. *Pract Radiat Oncol*. Jan - Feb 2017;7(1):4-12. PMID 27663933
28. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Bone Cancer. Version 2.2018. https://www.nccn.org/professionals/physician_gls/pdf/bone.pdf. Accessed June 4, 2018.
29. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Breast Cancer. Version 1.2018. https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf. Accessed June 4, 2018.
30. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Central Nervous System Cancers. Version 1.2018. https://www.nccn.org/professionals/physician_gls/pdf/cns.pdf. Accessed June 4, 2018.
31. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Prostate Cancer. Version 2.2018. https://www.nccn.org/professionals/physician_gls/pdf/prostate.pdf. Accessed June 4, 2018.