



# MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

## Medical Policy

### Diagnosis and Treatment of Chronic Cerebrospinal Venous Insufficiency in Multiple Sclerosis

#### Table of Contents

- [Policy: Commercial](#)
- [Coding Information](#)
- [Information Pertaining to All Policies](#)
- [Policy: Medicare](#)
- [Description](#)
- [References](#)
- [Authorization Information](#)
- [Policy History](#)

#### Policy Number: 352

BCBSA Reference Number: 8.01.56

NCD/LCD: N/A

#### Related Policies

None

#### Policy

#### Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO Blue<sup>SM</sup> and Medicare PPO Blue<sup>SM</sup> Members

The identification and subsequent treatment of chronic cerebrospinal venous insufficiency in patients with multiple sclerosis is considered **NOT MEDICALLY NECESSARY**.

#### Prior Authorization Information

Pre-service approval is required for all inpatient services for all products.

See below for situations where prior authorization may be required or may not be required.

Yes indicates that prior authorization is required.

No indicates that prior authorization is not required.

N/A indicates that this service is primarily performed in an inpatient setting.

#### Outpatient

<b>Commercial Managed Care (HMO and POS)</b>	This is not a covered service.
<b>Commercial PPO and Indemnity</b>	This is not a covered service.
<b>Medicare HMO Blue<sup>SM</sup></b>	This is not a covered service.
<b>Medicare PPO Blue<sup>SM</sup></b>	This is not a covered service.

## CPT Codes / HCPCS Codes / ICD Codes

The following codes are included below for informational purposes. Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

### CPT Codes

There is no specific code CPT for this test.

### Description

MS is generally considered a chronic inflammatory demyelinating disease of the central nervous system (brain, spinal cord, optic nerve) believed to be triggered by an autoimmune response to myelin. However, in part due to the periventricular predilection of the lesions of MS, vascular etiologies (CCSVI) have also been considered. The core foundation of this vascular theory is that venous drainage from the brain is abnormal due to outflow obstruction in the draining jugular vein and/or azygos veins. This abnormal venous drainage, which is characterized by special ultrasound criteria, is said to cause intracerebral flow disturbance or outflow problems that lead to periventricular deposits. In the CCSVI theory, these deposits have a similarity to the iron deposits seen around the veins in the legs of patients with chronic deep vein thrombosis. Balloon dilatation, with or without stenting, has been proposed as a means to treat the outflow problems, thereby alleviating CCSVI and MS complaints.

The following 5 criteria were defined by Zamboni et al as features of CCSVI.1 To make the diagnosis of CCSVI, at least 2 of the 5 criteria need to be present:

1. Reflux constantly present (for a duration >0.8 s) in the supine and upright positions at the level of an internal jugular or vertebral vein. This parameter was evaluated during a short breath-hold following normal breathing and not under Valsalva maneuver.
2. Reflux at the level of veins of the deep cerebral system (for a duration >0.5 s). This was evaluated with the patient in the sitting and supine positions, and venous flow was enhanced by inviting the patient to breath in.
3. Stenosis (<0.3 cm), valve abnormalities and septa on B-mode imaging.
4. Absence of flow at the level of the internal jugular or vertebral vein, despite numerous deep inspirations.
5. No increase in the diameter of the internal jugular vein when changing from an upright to a supine position (lack of  $\Delta$ -).

### Summary

Chronic cerebrospinal venous insufficiency (CCSVI) may be associated with multiple sclerosis (MS), although this is controversial and an active area of research. Correction of CCSVI has been attempted via percutaneous venoplasty. The intent of this procedure is to relieve MS symptoms by improving venous drainage of the central nervous system. Correction of CCSVI by this method may be referred to as the "Liberation Procedure."

The association of CCSVI with MS is uncertain. The rate of CCSVI in MS patients varies widely in the literature for unclear reasons, from 0% to 100%. Some studies report higher rates of CCSVI in patients with MS compared with non-MS patients, but others do not. If there is an association between MS and CCSVI, it is not known whether this is a causative factor for MS or a secondary result of the disease. It also appears that CCSVI can occur in other disorders and is not specific for MS.

Treatment of CCSVI with endovascular interventions has been attempted. Some currently available studies report improvement in patient-reported symptoms following treatment, but this evidence is not sufficient to establish efficacy. A prospective, double-blind, sham-controlled randomized controlled trial

(RCT) of venous angioplasty in MS patients (N=20) with CCSVI published in 2014 showed no significant differences in venous outflow characteristics between the treated and control groups, nor any significant improvements in clinical disease scores among treated patients compared with controls. The results of this RCT are limited by the small number of patients. However, the failure to show a beneficial effect of venous angioplasty on blood flow or symptoms supports a lack of efficacy for this treatment.

Adverse events occur at a low overall rate, but serious adverse events can occur, and the U.S. Food and Drug Administration (FDA) issued an alert in 2012 concerning the potential for serious adverse events with treatment of CCSVI.

## Policy History

Date	Action
10/2015	BCBSA National medical policy review. Policy statement changed from investigational to not medically necessary. Effective 10/1/2015.
9/2014	New references added from BCBSA National medical policy.
6/2013	New references from BCBSA National medical policy.
5/1/12	New policy describing ongoing non-coverage.

## Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

## References

1. Zamboni P, Galeotti R, Menegatti E et al. Chronic cerebrospinal venous insufficiency in patients with multiple sclerosis. *J Neurol Neurosurg Psychiatry* 2009; 80(4):392-9.
2. Zamboni P, Galeotti R, Menegatti E et al. A prospective open-label study of endovascular treatment of chronic cerebrospinal venous insufficiency. *J Vasc Surg* 2009; 50(6):1348-58 e1-3.
3. Tsvigoulis G, Sergentanis TN, Chan A et al. Chronic cerebrospinal venous insufficiency and multiple sclerosis: a comprehensive meta-analysis of case-control studies. *Ther Adv Neurol Disord* 2014; 7(2):114-36.
4. Zwischenberger BA, Beasley MM, Davenport DL et al. Meta-analysis of the correlation between chronic cerebrospinal venous insufficiency and multiple sclerosis. *Vasc Endovascular Surg* 2013; 47(8):620-4.
5. Laupacis A, Lillie E, Dueck A et al. Association between chronic cerebrospinal venous insufficiency and multiple sclerosis: a meta-analysis. *CMAJ* 2011; 183(16):E1203-12.
6. Thapar A, Lane T, Nicholas R et al. Systematic review of sonographic chronic cerebrospinal venous insufficiency findings in multiple sclerosis. *Phlebology* 2011; 26(8):319-25.
7. Zivadinov R, Marr K, Cutter G et al. Prevalence, sensitivity, and specificity of chronic cerebrospinal venous insufficiency in MS. *Neurology* 2011; 77(2):138-44.
8. Zivadinov R, Cutter G, Marr K et al. No association between conventional brain MR imaging and chronic cerebrospinal venous insufficiency in multiple sclerosis. *AJNR Am J Neuroradiol* 2012; 33(10):1913-7.
9. Weinstock-Guttman B, Ramanathan M, Marr K et al. Clinical correlates of chronic cerebrospinal venous insufficiency in multiple sclerosis. *BMC Neurol* 2012; 12:26.
10. Barreto AD, Brod SA, Bui TT et al. Chronic cerebrospinal venous insufficiency: Case-control neurosonography results. *Ann Neurol* 2012.
11. Floris R, Centonze D, Fabiano S et al. Prevalence study of chronic cerebrospinal venous insufficiency in patients with multiple sclerosis: preliminary data. *Radiol Med* 2012.

12. Centonze D, Floris R, Stefanini M et al. Proposed chronic cerebrospinal venous insufficiency criteria do not predict multiple sclerosis risk or severity. *Ann Neurol* 2011; 70(1):51-8.
13. Doepp F, Paul F, Valdueza JM et al. No cerebrocervical venous congestion in patients with multiple sclerosis. *Ann Neurol* 2010; 68(2):173-83.
14. Siddiqui AH, Zivadinov R, Benedict RH, et al. Prospective randomized trial of venous angioplasty in MS (PREMiSe). *Neurology*. Jul 29 2014;83(5):441-449. PMID 24975855
15. Bourdette DN, Cohen JA. Venous angioplasty for "CCSVI" in multiple sclerosis: ending a therapeutic misadventure. *Neurology*. Jul 29 2014;83(5):388-389. PMID 24975856
16. Tsivgoulis G, Faissner S, Voumvourakis K, et al. "Liberation treatment" for chronic cerebrospinal venous insufficiency in multiple sclerosis: the truth will set you free. *Brain Behav*. Jan 2015;5(1):3-12. PMID 25722945
17. van Zuuren EJ, Fedorowicz Z, Pucci E et al. Percutaneous transluminal angioplasty for treatment of chronic cerebrospinal venous insufficiency (CCSVI) in multiple sclerosis patients. *Cochrane Database Syst Rev* 2012; 12:CD009903.
18. Hubbard D, Ponec D, Gooding J et al. Clinical improvement after extracranial venoplasty in multiple sclerosis. *J Vasc Interv Radiol* 2012; 23(10):1302-8.
19. Zamboni P, Galeotti R, Weinstock-Guttman B et al. Venous angioplasty in patients with multiple sclerosis: results of a pilot study. *Eur J Vasc Endovasc Surg* 2012; 43(1):116-22.
20. Burton JM, Alikhani K, Goyal M et al. Complications in MS patients after CCSVI procedures abroad (Calgary, AB). *Can J Neurol Sci* 2011; 38(5):741-6.
21. Petrov I, Grozdinski L, Kaninski G et al. Safety profile of endovascular treatment for chronic cerebrospinal venous insufficiency in patients with multiple sclerosis. *J Endovasc Ther* 2011; 18(3):314-23.
22. Mandato KD, Hegener PF, Siskin GP et al. Safety of endovascular treatment of chronic cerebrospinal venous insufficiency: a report of 240 patients with multiple sclerosis. *J Vasc Interv Radiol* 2012; 23(1):55-9.
23. FDA News Release: FDA issues alert on potential dangers of unproven treatment for multiple sclerosis. Available online at: <http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm303538.htm?source=govdeli> very. Last accessed April, 2014.
24. Reekers JA, Lee MJ, Belli AM et al. Cardiovascular and Interventional Radiological Society of Europe commentary on the treatment of chronic cerebrospinal venous insufficiency. *Cardiovasc Intervent Radiol* 2011; 34(1):1-2.
25. Vedantham S, Benenati JF, Kundu S et al. Interventional endovascular management of chronic cerebrospinal venous insufficiency in patients with multiple sclerosis: a position statement by the Society of Interventional Radiology, endorsed by the Canadian Interventional Radiology Association. *J Vasc Interv Radiol* 2010; 21(9):1335-7.
26. NICE. Percutaneous venoplasty for chronic cerebrospinal venous insufficiency for multiple sclerosis. March 2012. Available online at: [www.nice.org.uk/ipg420](http://www.nice.org.uk/ipg420). Last accessed April, 2014.
27. Baracchini C, Valdueza JM, Del Sette M et al. CCSVI and MS: a statement from the European Society of neurosonology and cerebral hemodynamics. *J Neurol* 2012; 259(12):2585-9.