



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

Medical Policy

Vertebral Fracture Assessment with Densitometry

Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

Policy Number: 449

BCBSA Reference Number: 6.01.44

NCD/LCD: NA

Related Policies

- Bone Mineral Density Studies, #[450](#)
- Whole Body Dual X-Ray Absorptiometry to Determine Body Composition, #[577](#)

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO BlueSM and Medicare PPO BlueSM Members

Screening for vertebral fractures using dual x-ray absorptiometry (DEXA or DXA) is [INVESTIGATIONAL](#).

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	This is not a covered service.
Commercial PPO and Indemnity	This is not a covered service.
Medicare HMO BlueSM	This is not a covered service.
Medicare PPO BlueSM	This is not a covered service.

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

The following CPT codes are considered investigational for Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:

CPT Codes

CPT codes:	Code Description
77085	Dual-energy X-ray absorptiometry (DXA), bone density study, 1 or more sites; axial skeleton (eg, hips, pelvis, spine), including vertebral fracture assessment
77086	Vertebral fracture assessment via dual-energy X-ray absorptiometry (DXA)

Description

VERTEBRAL FRACTURES

Vertebral fractures are highly prevalent in the elderly population, and epidemiologic studies have found that these fractures are associated with an increased risk of future spine or hip fractures independent of bone mineral density.

Diagnosis

Only 20% to 30% of vertebral fractures are recognized clinically; the rest are discovered incidentally on lateral spine radiographs. Lateral spine radiographs have not been recommended as a component of risk assessment for osteoporosis because of the cost, radiation exposure, and the fact that the radiograph would require a separate procedure in addition to the bone mineral density study using dual-energy x-ray absorptiometry. However, several densitometers with specialized software can perform vertebral fracture assessment (VFA) in conjunction with dual-energy x-ray absorptiometry. The lateral spine scan is performed by using a rotating arm; depending on the densitometer used, the patient can either stay in the supine position after the bone density study or is required to move to the left decubitus position.

VFA differs from radiologic detection of fractures because VFA uses a lower radiation exposure and can detect only fractures, while traditional radiograph images can detect other bone and soft tissue abnormalities in addition to spinal fractures. Manufacturers have also referred to this procedure as instant vertebral assessment, radiographic vertebral assessment, dual-energy vertebral assessment, or lateral vertebral assessment.

For both lateral spine radiographs and images with densitometry, vertebral fractures are assessed visually. A number of grading systems have been proposed, and the Genant semiquantitative method is commonly used. This system grades deformities from I to III, with grade I (mild) representing a 20% to 24% reduction in vertebral height, grade II (moderate) representing a 25% to 39% reduction in height, and grade III (severe) representing a 40% or greater reduction in height. The location of the deformity within the vertebrae may also be noted. For example, if only the mid-height of the vertebrae is affected, the deformity is defined as an endplate deformity; if both the anterior and mid-heights are deformed, it is a wedge deformity; and if the entire vertebrae is deformed, it is classed as a crush deformity. A vertebral deformity of at least 20% loss in height is typically considered a fracture. Accurate interpretation of both lateral spine radiographs and VFA imaging depends on radiologic training. Thus, device location and availability of appropriately trained personnel may influence diagnostic accuracy.

Summary

Vertebral fracture assessment (VFA) with densitometry is a technique to assess vertebral fractures at the same time as bone mineral density, using additional software with dual-energy x-ray absorptiometry. The addition of VFA to bone mineral density may augment diagnostic information on fracture risk.

For individuals who are at risk of having vertebral fractures but are not known to have them who receive VFA with densitometry by dual-energy x-ray absorptiometry, the evidence includes diagnostic accuracy

studies and subgroup reanalyses of treatment studies. Relevant outcomes are test accuracy, test validity, and morbid events. There is a lack of direct evidence from screening trials that use densitometry with and without VFA improves health outcomes. Because direct evidence was not available, a chain of evidence was sought. Evidence was examined on the diagnostic accuracy of VFA in nonosteoporotic patients (ie, those not already eligible for treatment), the ability of VFA to identify patients for treatment who would not otherwise be identified, and the effectiveness of treatment in this population. Diagnostic accuracy studies have reported variable findings; recent studies have suggested higher diagnostic accuracy of VFA overall compared with standard radiographs than older studies. Studies have found that VFA can identify patients without osteoporosis who may be appropriate candidates for treatment according to recommendations from the National Osteoporosis Foundation. However, there is limited evidence on the effectiveness of treatment in this population. No treatment data have been published on patients whose vertebral fractures were identified using VFA software with densitometry. The evidence is insufficient to determine the effects of the technology on health outcomes.

Policy History

Date	Action
11/2018	BCBSA National medical policy review. Description, summary and references updated. Policy statement unchanged.
10/2017	New references added from BCBSA National medical policy.
10/2016	New references added from BCBSA National medical policy.
1/2015	Clarified coding information.
2/2013	New references from BCBSA National medical policy.
11/2011-4/2012	Medical policy ICD 10 remediation: Formatting, editing and coding updates. No changes to policy statements.
9/2011	Reviewed - Medical Policy Group - Urology and Obstetrics/Gynecology. No changes to policy statements.
1/2011	BCBSA National medical policy review. No changes to policy statements.
10/2010	Reviewed - Medical Policy Group - Urology and Obstetrics/Gynecology. No changes to policy statements.
10/2009	Reviewed - Medical Policy Group - Urology and Obstetrics/Gynecology. No changes to policy statements.
10/2008	Reviewed - Medical Policy Group - Urology and Obstetrics/Gynecology. No changes to policy statements.
10/2007	Reviewed - Medical Policy Group - Urology and Obstetrics/Gynecology. No changes to policy statements.
8/2007	BCBSA National medical policy review. No changes to policy statements.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

References

1. Lee JH, Lee YK, Oh SH, et al. A systematic review of diagnostic accuracy of vertebral fracture assessment (VFA) in postmenopausal women and elderly men. *Osteoporos Int*. May 2016;27(5):1691-1699. PMID 26782682
2. Domiciano DS, Figueiredo CP, Lopes JB, et al. Vertebral fracture assessment by dual X-ray absorptiometry: a valid tool to detect vertebral fractures in community-dwelling older adults in a population-based survey. *Arthritis Care Res (Hoboken)*. May 2013;65(5):809-815. PMID 23212896

3. Ferrar L, Jiang G, Clowes JA, et al. Comparison of densitometric and radiographic vertebral fracture assessment using the algorithm-based qualitative (ABQ) method in postmenopausal women at low and high risk of fracture. *J Bone Miner Res.* Jan 2008;23(1):103-111. PMID 17892377
4. Binkley N, Krueger D, Gangnon R, et al. Lateral vertebral assessment: a valuable technique to detect clinically significant vertebral fractures. *Osteoporos Int.* Dec 2005;16(12):1513-1518. PMID 15834512
5. Cosman F, de Beur SJ, LeBoff MS, et al. Clinician's guide to prevention and treatment of osteoporosis. *Osteoporos Int.* Oct 2014;25(10):2359-2381. PMID 25182228
6. Centre for Metabolic Bone Diseases, University of Sheffield U. FRAX Fracture Risk Assessment Tool: Calculation Tool. n.d.; <https://www.sheffield.ac.uk/FRAX/tool.aspx?country=9>. Accessed August 31, 2017.
7. Kanterewicz E, Puigoriol E, Garcia-Barrionuevo J, et al. Prevalence of vertebral fractures and minor vertebral deformities evaluated by DXA-assisted vertebral fracture assessment (VFA) in a population-based study of postmenopausal women: the FRODOS study. *Osteoporos Int.* May 2014;25(5):1455-1464. PMID 24599272
8. Mrgan M, Mohammed A, Gram J. Combined vertebral assessment and bone densitometry increases the prevalence and severity of osteoporosis in patients referred to DXA scanning. *J Clin Densitom.* Oct-Dec 2013;16(4):549-553. PMID 23769657
9. Jager PL, Jonkman S, Koolhaas W, et al. Combined vertebral fracture assessment and bone mineral density measurement: a new standard in the diagnosis of osteoporosis in academic populations. *Osteoporos Int.* Apr 2011;22(4):1059-1068. PMID 20571773
10. Cummings SR, Black DM, Thompson DE, et al. Effect of alendronate on risk of fracture in women with low bone density but without vertebral fractures: results from the Fracture Intervention Trial. *Jama.* Dec 23-30 1998;280(24):2077-2082. PMID 9875874
11. Quandt SA, Thompson DE, Schneider DL, et al. Effect of alendronate on vertebral fracture risk in women with bone mineral density T scores of -1.6 to -2.5 at the femoral neck: the Fracture Intervention Trial. *Mayo Clin Proc.* Mar 2005;80(3):343-349. PMID 15757015
12. Kanis JA, Barton IP, Johnell O. Risedronate decreases fracture risk in patients selected solely on the basis of prior vertebral fracture. *Osteoporos Int.* May 2005;16(5):475-482. PMID 15875093
13. Bhoopalam N, Campbell SC, Moritz T, et al. Intravenous zoledronic acid to prevent osteoporosis in a veteran population with multiple risk factors for bone loss on androgen deprivation therapy. *J Urol.* Nov 2009;182(5):2257-2264. PMID 19758618
14. Greenspan SL, Nelson JB, Trump DL, et al. Effect of once-weekly oral alendronate on bone loss in men receiving androgen deprivation therapy for prostate cancer: a randomized trial. *Ann Intern Med.* Mar 20 2007;146(6):416-424. PMID 17371886
15. Rosen HN, Vokes TJ, Malabanan AO, et al. The official positions of the International Society for Clinical Densitometry: vertebral fracture assessment. *J Clin Densitom.* Oct-Dec 2013;16(4):482-488. PMID 24063846
16. International Society for Clinical Densitometry. 2015 ISCD Official Positions – Adult. 2015; <https://www.iscd.org/official-positions/2015-iscd-official-positions-adult/>. Accessed August 17, 2018.
17. Camacho PM, Petak SM, Binkley N, et al. American Association of Clinical Endocrinologists and American College of Endocrinology clinical practice guidelines for the diagnosis and treatment of postmenopausal osteoporosis - 2016. *Endocr Pract.* Sep 02 2016;22(Suppl 4):1-42. PMID 27662240
18. Watts NB, Adler RA, Bilezikian JP, et al. Osteoporosis in men: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab.* Jun 2012;97(6):1802-1822. PMID 22675062
19. Qaseem A, Forcica MA, McLean RM, et al. Treatment of low bone density or osteoporosis to prevent fractures in men and women: a clinical practice guideline update from the American College of Physicians. *Ann Intern Med.* Jun 06 2017;166(11):818-839. PMID 28492856
20. Management of osteoporosis in postmenopausal women: 2010 position statement of The North American Menopause Society. *Menopause.* Jan-Feb 2010;17(1):25-54; quiz 55-56. PMID 20061894
21. U.S. Preventive Services Task Force. Osteoporosis to Prevent Fractures: Screening. 2018; <https://www.uspreventiveservicestaskforce.org/Page/Document/UpdateSummaryFinal/osteoporosis-screening1>. Accessed August 17, 2018.
22. U. S. Preventive Services Task Force, Curry SJ, Krist AH, et al. Screening for osteoporosis to prevent fractures: US Preventive Services Task Force Recommendation Statement. *JAMA.* Jun 26 2018;319(24):2521-2531. PMID 29946735