



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an independent
Licensee of the Blue Cross and Blue Shield Association

Medical Policy

Dermatologic Applications of Photodynamic Therapy

Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

Policy Number: 463

BCBSA Reference Number: 2.01.44

NCD/LCD: N/A

Related Policies

- Light Therapy for Psoriasis, #[698](#)
- Oncologic Applications of Photodynamic Therapy, Including Barrett's Esophagus, #[454](#)
- Photodynamic Therapy for Choroidal Neovascularization, #[599](#)

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO BlueSM and Medicare PPO BlueSM Members

Photodynamic therapy may be considered **MEDICALLY NECESSARY** as a treatment of:

- Nonhyperkeratotic actinic keratoses of the face and scalp.
- Low-risk (eg superficial and nodular) basal cell skin cancer only when surgery and radiation are contraindicated.
- Cutaneous squamous cell carcinoma in situ (Bowen disease) only when surgery and radiation are contraindicated.

Photodynamic therapy is considered **INVESTIGATIONAL** for other dermatologic applications, including, but not limited to, acne vulgaris, high-risk basal cell carcinomas, hidradenitis suppurativa and mycoses.

Photodynamic therapy as a technique of skin rejuvenation, hair removal, or other cosmetic indications is considered **NOT MEDICALLY NECESSARY**.

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for situations where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	Prior authorization is not required .
Commercial PPO and Indemnity	Prior authorization is not required .
Medicare HMO BlueSM	Prior authorization is not required .
Medicare PPO BlueSM	Prior authorization is not required .

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

The above medical necessity criteria MUST be met for the following codes to be covered for Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:

CPT Codes

CPT codes:	Code Description
96567	Photodynamic therapy by external application of light to destroy premalignant lesions of the skin and adjacent mucosa <u>with application</u> and <u>illumination</u> /activation of photosensitive drug(s), <u>per day</u>
96573	Photodynamic therapy by external application of light to destroy premalignant lesions of the skin and adjacent mucosa with application and illumination/activation of photosensitizing drug(s) provided by a physician or other qualified health care professional, per day
96574	Debridement of premalignant hyperkeratotic lesion(s) (ie, targeted curettage, abrasion) followed with photodynamic therapy by external application of light to destroy premalignant lesions of the skin and adjacent mucosa with application and illumination/activation of photosensitizing drug(s) provided by a physician or other qualified health care professional, per day

HCPCS Codes

HCPCS codes:	Code Description
J7308	Aminolevulinic hydrochloric acid for topical administration, 20%, single unit dosage form (354 mg)
J7309	Methyl aminolevulinate (MAL) for topical administration, 16.8%, 1 gram
J7345	Aminolevulinic acid HCl for topical administration, 10% gel, 10 mg

The following ICD Diagnosis Codes are considered medically necessary when submitted with the CPT and HCPCS codes above if medical necessity criteria are met:

ICD- 10 Diagnosis Codes

ICD-10-CM Diagnosis codes:	Code Description

C44.01	Basal Cell Carcinoma Of Skin Of Lip
C44.111	Basal Cell Carcinoma Of Skin Of Unspecified Eyelid, Including Canthus
C44.112	Basal Cell Carcinoma Of Skin Of Right Eyelid, Including Canthus
C44.119	Basal Cell Carcinoma Of Skin Of Left Eyelid, Including Canthus
C44.211	Basal Cell Carcinoma Of Skin Of Unspecified Ear And External Auricular Canal
C44.212	Basal Cell Carcinoma Of Skin Of Right Ear And External Auricular Canal
C44.219	Basal Cell Carcinoma Of Skin Of Left Ear And External Auricular Canal
C44.310	Basal Cell Carcinoma Of Skin Of Unspecified Parts Of Face
C44.311	Basal Cell Carcinoma Of Skin Of Nose
C44.319	Basal Cell Carcinoma Of Skin Of Other Parts Of Face
C44.41	Basal Cell Carcinoma Of Skin Of Scalp And Neck
C44.510	Basal Cell Carcinoma Of Anal Skin
C44.511	Basal Cell Carcinoma Of Skin Of Breast
C44.519	Basal Cell Carcinoma Of Skin Of Other Part Of Trunk
C44.611	Basal Cell Carcinoma Of Skin Of Unspecified Upper Limb, Including Shoulder
C44.612	Basal Cell Carcinoma Of Skin Of Right Upper Limb, Including Shoulder
C44.619	Basal Cell Carcinoma Of Skin Of Left Upper Limb, Including Shoulder
C44.711	Basal Cell Carcinoma Of Skin Of Unspecified Lower Limb, Including Hip
C44.712	Basal Cell Carcinoma Of Skin Of Right Lower Limb, Including Hip
C44.719	Basal Cell Carcinoma Of Skin Of Left Lower Limb, Including Hip
C44.81	Basal Cell Carcinoma Of Overlapping Sites Of Skin
C44.91	Basal Cell Carcinoma Of Skin, Unspecified
D04.0	Carcinoma In Situ Of Skin Of Lip
D04.10	Carcinoma In Situ Of Skin Of Unspecified Eyelid, Including Canthus
D04.111	Carcinoma in situ of skin of right upper eyelid, including canthus
D04.112	Carcinoma in situ of skin of right lower eyelid, including canthus
D04.121	Carcinoma in situ of skin of left upper eyelid, including canthus
D04.122	Carcinoma in situ of skin of left lower eyelid, including canthus
D04.20	Carcinoma In Situ Of Skin Of Unspecified Ear And External Auricular Canal
D04.21	Carcinoma In Situ Of Skin Of Right Ear And External Auricular Canal
D04.22	Carcinoma In Situ Of Skin Of Left Ear And External Auricular Canal
D04.30	Carcinoma In Situ Of Skin Of Unspecified Part Of Face
D04.39	Carcinoma In Situ Of Skin Of Other Parts Of Face
D04.4	Carcinoma In Situ Of Skin Of Scalp And Neck
D04.5	Carcinoma In Situ Of Skin Of Trunk
D04.60	Carcinoma In Situ Of Skin Of Unspecified Upper Limb, Including Shoulder
D04.61	Carcinoma In Situ Of Skin Of Right Upper Limb, Including Shoulder
D04.62	Carcinoma In Situ Of Skin Of Left Upper Limb, Including Shoulder
D04.70	Carcinoma In Situ Of Skin Of Unspecified Lower Limb, Including Hip
D04.71	Carcinoma In Situ Of Skin Of Right Lower Limb, Including Hip
D04.72	Carcinoma In Situ Of Skin Of Left Lower Limb, Including Hip
D04.8	Carcinoma In Situ Of Skin Of Other Sites
D04.9	Carcinoma In Situ Of Skin, Unspecified
L57.0	Actinic Keratosis

Description

Photodynamic Therapy

PDT refers to light activation of a photosensitizer to generate highly reactive intermediaries, which ultimately cause tissue injury and necrosis. Two common photosensitizing agents are 5-aminolevulinic acid (5-ALA) and its methyl ester, methyl aminolevulinate. When applied topically, these agents pass

readily through abnormal keratin overlying the lesion and accumulate preferentially in dysplastic cells. The agents 5-ALA and methyl aminolevulinic acid are metabolized by underlying cells to photosensitizing concentrations of porphyrins. Subsequent exposure to photoactivation (maximum absorption at 404 to 420 nm and 635 nm) generates reactive oxygen species that are cytotoxic, ultimately destroying the lesion. PDT can cause erythema, burning, and pain. Healing occurs within 10 to 14 days, with generally acceptable cosmetic results. PDT with topical ALA has been investigated primarily as a treatment of actinic keratoses.

Summary

Photodynamic therapy (PDT) refers to light activation of a photosensitizer to generate highly reactive intermediaries, which ultimately cause tissue injury and necrosis. Photosensitizing agents are being proposed for use with dermatologic conditions such as actinic keratoses and nonmelanoma skin cancers.

For individuals who have nonhyperkeratotic actinic keratoses on the face or scalp who receive PDT, the evidence includes randomized controlled trials (RCTs). The relevant outcomes are symptoms, change in disease status, quality of life, and treatment-related morbidity. Evidence from multiple RCTs has found that PDT improves the net health outcome in patients with nonhyperkeratotic actinic keratoses on the face or scalp compared with placebo or other active interventions. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have low-risk basal cell carcinoma who receive PDT, the evidence includes RCTs and systematic reviews of RCTs. The relevant outcomes are symptoms, change in disease status, quality of life, and treatment-related morbidity. Systematic reviews of RCTs have found that PDT may not be as effective as surgery for superficial and nodular basal cell carcinoma. In the small number of trials available, PDT was more effective than placebo. The available evidence from RCTs has suggested that PDT has better cosmetic outcomes than surgery. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have squamous cell carcinoma in situ who receive PDT, the evidence includes RCTs. The relevant outcomes are symptoms, change in disease status, quality of life, and treatment-related morbidity. RCTs have found that PDT has similar or greater efficacy compared with cryotherapy and 5-fluorouracil. Additionally, adverse events and cosmetic outcomes appear to be better after PDT. Few RCTs have compared PDT with surgery or radiotherapy; as a result, conclusions cannot be drawn about PDT compared with these other standard treatments. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have nonmetastatic invasive squamous cell carcinoma who receive PDT, the evidence includes observational studies and a systematic review of observational studies. The relevant outcomes are overall survival, symptoms, change in disease status, quality of life, and treatment-related morbidity. Conclusions cannot be drawn from small, uncontrolled studies. RCTs are needed to determine the safety and efficacy of PDT for this condition. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have acne who receive PDT, the evidence includes RCTs and a systematic review. The relevant outcomes are symptoms, change in disease status, quality of life, and treatment-related morbidity. The available RCTs have not consistently found significantly better outcomes with PDT compared with other interventions, and a meta-analysis did not find significantly better results with PDT vs placebo. Several trials have found that PDT is associated with high rates of adverse events leading to the cessation of treatment. Trials tended to have relatively small sample sizes and used a variety of comparison interventions. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have noncancerous dermatologic skin conditions (eg, hidradenitis suppurativa, mycoses, port wine stain) who receive PDT, the evidence includes case series, systematic reviews of uncontrolled series, and an RCT for port wine stain. The relevant outcomes are symptoms, change in disease status, quality of life, and treatment-related morbidity. RCTs are needed to determine the safety

and efficacy of PDT for these conditions. The evidence is insufficient to determine the effects of the technology on health outcomes.

Policy History

Date	Action
2/2019	BCBSA National medical policy review. Description, summary and references updated. Policy statements unchanged.
10/2018	Clarified coding information.
5/2018	Clarified coding information.
3/2018	New references added from BCBSA National medical policy.
1/2018	Clarified coding information.
1/2017	New references added from BCBSA National medical policy.
3/2016	New references added from BCBSA National medical policy.
6/2015	BCBSA National medical policy review. Superficial basal cell carcinoma changed to low-risk (ie superficial or nodular) basal cell carcinoma. Non-superficial basal cell carcinoma changed to high-risk basal cell carcinoma. Dermatologic Applications of Photodynamic Therapy transferred from policy #068, Plastic Surgery. Effective 6/1/2015.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

References

1. Pariser DM, Lowe NJ, Stewart DM, et al. Photodynamic therapy with topical methyl aminolevulinate for actinic keratosis: results of a prospective randomized multicenter trial. *J Am Acad Dermatol.* Feb 2003;48(2):227-232. PMID 12582393
2. Morton C, Campbell S, Gupta G, et al. Intraindividual, right-left comparison of topical methyl aminolaevulinate-photodynamic therapy and cryotherapy in subjects with actinic keratoses: a multicentre, randomized controlled study. *Br J Dermatol.* Nov 2006;155(5):1029-1036. PMID 17034536
3. Hauschild A, Stockfleth E, Popp G, et al. Optimization of photodynamic therapy with a novel self-adhesive 5-aminolaevulinic acid patch: results of two randomized controlled phase III studies. *Br J Dermatol.* May 2009;160(5):1066-1074. PMID 19222455
4. Szeimies RM, Radny P, Sebastian M, et al. Photodynamic therapy with BF-200 ALA for the treatment of actinic keratosis: results of a prospective, randomized, double-blind, placebo-controlled phase III study. *Br J Dermatol.* Aug 2010;163(2):386-394. PMID 20518784
5. Szeimies RM, Stockfleth E, Popp G, et al. Long-term follow-up of photodynamic therapy with a self-adhesive 5-aminolaevulinic acid patch: 12 months data. *Br J Dermatol.* Feb 1 2010;162(2):410-414. PMID 19804593
6. Serra-Guillen C, Nagore E, Hueso L, et al. A randomized pilot comparative study of topical methyl aminolevulinate photodynamic therapy versus imiquimod 5% versus sequential application of both therapies in immunocompetent patients with actinic keratosis: clinical and histologic outcomes. *J Am Acad Dermatol.* Apr 2012;66(4):e131-137. PMID 22226430
7. Dirschka T, Radny P, Dominicus R, et al. Photodynamic therapy with BF-200 ALA for the treatment of actinic keratosis: results of a multicentre, randomized, observer-blind phase III study in comparison with a registered methyl-5-aminolaevulinate cream and placebo. *Br J Dermatol.* Jan 2012;166(1):137-146. PMID 21910711

8. Dirschka T, Radny P, Dominicus R, et al. Long-term (6 and 12 months) follow-up of two prospective, randomized, controlled phase III trials of photodynamic therapy with BF-200 ALA and methyl aminolaevulinate for the treatment of actinic keratosis. *Br J Dermatol.* Apr 2013;168(4):825-836. PMID 23252768
9. Giehl KA, Kriz M, Grahovac M, et al. A controlled trial of photodynamic therapy of actinic keratosis comparing different red light sources. *Eur J Dermatol.* May-Jun 2014;24(3):335-341. PMID 24876164
10. Rubel DM, Spelman L, Murrell DF, et al. Daylight photodynamic therapy with methyl aminolevulinate cream as a convenient, similarly effective, nearly painless alternative to conventional photodynamic therapy in actinic keratosis treatment: a randomized controlled trial. *Br J Dermatol.* Nov 2014;171(5):1164-1171. PMID 24861492
11. Neittaanmaki-Perttu N, Karppinen TT, Gronroos M, et al. Daylight photodynamic therapy for actinic keratoses: a randomized double-blinded nonsponsored prospective study comparing 5-aminolaevulinic acid nanoemulsion (BF-200) with methyl-5-aminolaevulinate. *Br J Dermatol.* Nov 2014;171(5):1172-1180. PMID 25109244
12. Zane C, Facchinetti E, Rossi MT, et al. A randomized clinical trial of photodynamic therapy with methyl aminolaevulinate vs. diclofenac 3% plus hyaluronic acid gel for the treatment of multiple actinic keratoses of the face and scalp. *Br J Dermatol.* May 2014;170(5):1143-1150. PMID 24506666
13. Reinhold U, Dirschka T, Ostendorf R, et al. A randomized, double-blind, phase III, multicentre study to evaluate the safety and efficacy of BF-200 ALA (Ameluz((R))) vs. placebo in the field-directed treatment of mild-to-moderate actinic keratosis with photodynamic therapy (PDT) when using the BF-RhodoLED((R)) lamp. *Br J Dermatol.* Oct 2016;175(4):696-705. PMID 26921093
14. Yazdanyar S, Zarchi K, Jemec GBE. Pain during topical photodynamic therapy - comparing methyl aminolevulinate (Metvix((R))) to aminolaevulinic acid (Ameluz((R))); an intra-individual clinical study. *Photodiagnosis Photodyn Ther.* Aug 02 2017. PMID 28780136
15. Patel G, Armstrong AW, Eisen DB. Efficacy of photodynamic therapy vs other interventions in randomized clinical trials for the treatment of actinic keratoses: a systematic review and meta-analysis. *JAMA Dermatol.* Dec 2014;150(12):1281-1288. PMID 25162181
16. Wang H, Xu Y, Shi J, et al. Photodynamic therapy in the treatment of basal cell carcinoma: a systematic review and meta-analysis. *Photodermatol Photoimmunol Photomed.* Jan 2015;31(1):44-53. PMID 25377432
17. Zou Y, Zhao Y, Yu J, et al. Photodynamic therapy versus surgical excision to basal cell carcinoma: meta-analysis. *J Cosmet Dermatol.* Dec 2016;15(4):374-382. PMID 27363535
18. Bath-Hextall FJ, Perkins W, Bong J, et al. Interventions for basal cell carcinoma of the skin. *Cochrane Database Syst Rev.* Jan 24 2007(1):CD003412. PMID 17253489
19. Roozeboom MH, Arits AH, Mosterd K, et al. Three-year follow-up results of photodynamic therapy vs. imiquimod vs. fluorouracil for treatment of superficial basal cell carcinoma: a single-blind, noninferiority, randomized controlled trial. *J Invest Dermatol.* Aug 2016;136(8):1568-1574. PMID 27113429
20. Szeimies RM, Ibbotson S, Murrell DF, et al. A clinical study comparing methyl aminolevulinate photodynamic therapy and surgery in small superficial basal cell carcinoma (8-20 mm), with a 12-month follow-up. *J Eur Acad Dermatol Venereol.* Nov 2008;22(11):1302-1311. PMID 18624836
21. Rhodes LE, de Rie M, Enstrom Y, et al. Photodynamic therapy using topical methyl aminolevulinate vs surgery for nodular basal cell carcinoma: results of a multicenter randomized prospective trial. *Arch Dermatol.* Jan 2004;140(1):17-23. PMID 14732655
22. Rhodes LE, de Rie MA, Leifsdottir R, et al. Five-year follow-up of a randomized, prospective trial of topical methyl aminolevulinate photodynamic therapy vs surgery for nodular basal cell carcinoma. *Arch Dermatol.* Sep 2007;143(9):1131-1136. PMID 17875873
23. Bath-Hextall FJ, Matin RN, Wilkinson D, et al. Interventions for cutaneous Bowen's disease. *Cochrane Database Syst Rev.* Jun 24 2013;6(6):CD007281. PMID 23794286
24. Morton C, Horn M, Leman J, et al. Comparison of topical methyl aminolevulinate photodynamic therapy with cryotherapy or Fluorouracil for treatment of squamous cell carcinoma in situ: Results of a multicenter randomized trial. *Arch Dermatol.* Jun 2006;142(6):729-735. PMID 16785375
25. Salim A, Leman JA, McColl JH, et al. Randomized comparison of photodynamic therapy with topical 5-fluorouracil in Bowen's disease. *Br J Dermatol.* Mar 2003;148(3):539-543. PMID 12653747

26. Lansbury L, Bath-Hextall F, Perkins W, et al. Interventions for non-metastatic squamous cell carcinoma of the skin: systematic review and pooled analysis of observational studies. *BMJ*. Nov 04 2013;347:f6153. PMID 24191270
27. Barbaric J, Abbott R, Posadzki P, et al. Light therapies for acne. *Cochrane Database Syst Rev*. Sep 27 2016;9:CD007917. PMID 27670126
28. Nicklas C, Rubio R, Cardenas C, et al. Comparison of efficacy of aminolaevulinic acid photodynamic therapy vs. adapalene gel plus oral doxycycline for treatment of moderate acne vulgaris-A simple, blind, randomized, and controlled trial. *Photodermatol Photoimmunol Photomed*. Jul 11 2018. PMID 29993146
29. Xu X, Zheng Y, Zhao Z, et al. Efficacy of photodynamic therapy combined with minocycline for treatment of moderate to severe facial acne vulgaris and influence on quality of life. *Medicine (Baltimore)*. Dec 2017;96(51):e9366. PMID 29390528
30. Pariser DM, Eichenfield LF, Bukhalo M, et al. Photodynamic therapy with methyl aminolaevulinate 80 mg g(-1) for severe facial acne vulgaris: a randomized vehicle-controlled study. *Br J Dermatol*. Apr 2016;174(4):770-777. PMID 26663215
31. Orringer JS, Sachs DL, Bailey E, et al. Photodynamic therapy for acne vulgaris: a randomized, controlled, split-face clinical trial of topical aminolevulinic acid and pulsed dye laser therapy. *J Cosmet Dermatol*. Mar 2010;9(1):28-34. PMID 20367670
32. Wiegell SR, Wulf HC. Photodynamic therapy of acne vulgaris using methyl aminolaevulinate: a blinded, randomized, controlled trial. *Br J Dermatol*. May 2006;154(5):969-976. PMID 16634903
33. Wu Q, Tu P, Zhou G, et al. A dose-finding study for hemoporphin in photodynamic therapy for port-wine stain: A multicenter randomized double-blind phase IIb trial. *Photodermatol Photoimmunol Photomed*. Sep 2018;34(5):314-321. PMID 29533491
34. Gold M, Bridges TM, Bradshaw VL, et al. ALA-PDT and blue light therapy for hidradenitis suppurativa. *J Drugs Dermatol*. Jan-Feb 2004;3(1 Suppl):S32-35. PMID 14964759
35. Schweiger ES, Riddle CC, Aires DJ. Treatment of hidradenitis suppurativa by photodynamic therapy with aminolevulinic acid: preliminary results. *J Drugs Dermatol*. Apr 2011;10(4):381-386. PMID 21455548
36. Calzavara-Pinton PG, Venturini M, Capezzer R, et al. Photodynamic therapy of interdigital mycoses of the feet with topical application of 5-aminolevulinic acid. *Photodermatol Photoimmunol Photomed*. Jun 2004;20(3):144-147. PMID 15144392
37. Mostafa D, Tarakji B. Photodynamic therapy in treatment of oral lichen planus. *J Clin Med Res*. Jun 2015;7(6):393-399. PMID 25883701
38. Yazdani Abyaneh MA, Falto-Aizpurua L, Griffith RD, et al. Photodynamic therapy for actinic cheilitis: a systematic review. *Dermatol Surg*. Feb 2015;41(2):189-198. PMID 25627629
39. Xiao Q, Li Q, Yuan KH, et al. Photodynamic therapy of port-wine stains: long-term efficacy and complication in Chinese patients. *J Dermatol*. Dec 2011;38(12):1146-1152. PMID 22032688
40. Zloty D, Guenther LC, Sapijaszko M, et al. Non-melanoma skin cancer in Canada. Chapter 4: Management of basal cell carcinoma. *J Cutan Med Surg*. May-Jun 2015;19(3):239-248. PMID 25986316
41. Poulin Y, Lynde CW, Barber K, et al. Non-melanoma skin cancer in Canada. Chapter 3: Management of actinic keratoses. *J Cutan Med Surg*. May-Jun 2015;19(3):227-238. PMID 25926621
42. National Comprehensive Cancer Network (NCCN). NCCN Practice Guidelines in Oncology: Basal cell skin cancer. Version 1.2019. http://www.nccn.org/professionals/physician_gls/pdf/nmsc.pdf. Accessed November 26, 2018.
43. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Squamous Cell Skin Cancer. Version 2.2019. https://www.nccn.org/professionals/physician_gls/pdf/squamous.pdf. Accessed November 26, 2018.
44. Centers for Medicare & Medicaid Services. National Coverage Determination (NCD) for Treatment of Actinic Keratosis (250.4). 2001; [https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCAId=238&NcaName=Allogeneic+Hematopoietic+Stem+Cell+Transplantation+\(HSCT\)+for+Myelodysplastic+Syndrome&ExpandComments=y&CommentPeriod=0&NCDId=129&ncdver=1&bc=AiAAAAAIEAAAA%3D%3D&](https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCAId=238&NcaName=Allogeneic+Hematopoietic+Stem+Cell+Transplantation+(HSCT)+for+Myelodysplastic+Syndrome&ExpandComments=y&CommentPeriod=0&NCDId=129&ncdver=1&bc=AiAAAAAIEAAAA%3D%3D&). Accessed November 26, 2018.