



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

## Medical Policy

# Implantable Bone-Conduction and Bone-Anchored Hearing Aids

### Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

### Policy Number: 479

BCBSA Reference Number: 7.01.03

NCD/LCD: N/A

### Related Policies

- Cochlear Implant, #[478](#)
- Semi-Implantable and Fully Implantable Middle Ear Hearing Aid #[480](#)

### Policy

#### Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO Blue<sup>SM</sup> and Medicare PPO Blue<sup>SM</sup> Members

Unilateral or bilateral fully or partially implantable bone-conduction (bone-anchored) hearing aid(s) may be **MEDICALLY NECESSARY** as an alternative to an air-conduction hearing aid in patients 5 years of age and older with conductive or mixed hearing loss who also meets at least one of the following medical criteria:

- Congenital or surgically induced malformations (e.g., atresia) of the external ear canal or middle ear
- Chronic external otitis or otitis media
- Tumors of the external canal and/or tympanic cavity, or
- Dermatitis of the external canal

**AND** meets the following audiologic criteria:

- A pure tone average bone-conduction threshold measured at 0.5, 1, 2, and 3 kHz of better than or equal to 45 dB (OBC and BP100 devices), 55 Db (Intenso device) or 65 dB (Cordele II device).

For bilateral implantation, patients should meet the above audiologic criteria and have symmetrically conductive or mixed hearing loss as defined by a difference between left- and right-side bone-conduction threshold of less than 10 dB on average measured at 0.5, 1, 2, and 3 kHz (4 kHz for OBC and Ponto Pro), or less than 15 dB at individual frequencies.

An implantable bone-conduction (bone-anchored) hearing aid may be **MEDICALLY NECESSARY** as an alternative to an air-conduction contralateral routing of signal hearing aid in patients 5 years of age and older with single-sided sensorineural deafness and normal hearing in the other ear. The pure tone

average air-conduction threshold of the normal ear should be better than 20 dB measured at 0.5, 1, 2, and 3 kHz.

Other uses of bone-conduction (bone-anchored) hearing aids, including use in patients with bilateral sensorineural hearing loss, are **INVESTIGATIONAL**.

### Prior Authorization Information

#### Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

#### Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	Prior authorization is <b>not required</b> .
Commercial PPO and Indemnity	Prior authorization is <b>not required</b> .
Medicare HMO Blue <sup>SM</sup>	Prior authorization is <b>not required</b> .
Medicare PPO Blue <sup>SM</sup>	Prior authorization is <b>not required</b> .

### CPT Codes / HCPCS Codes / ICD Codes

*Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.*

*Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.*

*The following codes are included below for informational purposes only; this is not an all-inclusive list.*

The above **medical necessity criteria** **MUST** be met for the following codes to be covered for **Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:**

#### CPT Codes

CPT codes:	Code Description
69710	Implantation or replacement of electromagnetic bone conduction hearing device in temporal bone
69714	Implantation, osseointegrated implant, temporal bone, with percutaneous attachment to external speech processor/cochlear stimulator; without mastoidectomy
69715	Implantation, osseointegrated implant, temporal bone, with percutaneous attachment to external speech processor/cochlear stimulator; with mastoidectomy

#### HCPCS Codes

HCPCS codes:	Code Description
L8690	Auditory osseointegrated device, includes all internal and external components

The following ICD Diagnosis Codes are considered medically necessary when submitted with the CPT/HCPCS codes above if **medical necessity criteria** are met:

## ICD-9 Diagnosis Codes

<b>ICD-9-CM diagnosis codes:</b>	<b>Code Description</b>
160.1	Malignant neoplasm of auditory tube, middle ear, and mastoid air cells
173.20	Unspecified malignant neoplasm of skin of ear and external auditory canal
380.15	Chronic mycotic otitis externa
380.16	Other chronic infective otitis externa
380.23	Other chronic otitis externa
380.52	Acquired stenosis of external ear canal secondary to surgery
381.10	Chronic serous otitis media, simple or unspecified
381.19	Other chronic serous otitis media
381.20	Chronic mucoid otitis media, simple or unspecified
381.29	Other chronic mucoid otitis media
381.3	Other and unspecified chronic nonsuppurative otitis media
382.1	Chronic tubotympanic suppurative otitis media
382.2	Chronic atticoantral suppurative otitis media
382.3	Unspecified chronic suppurative otitis media
389.00	Conductive hearing loss, unspecified
389.01	Conductive hearing loss, external ear
389.02	Conductive hearing loss, tympanic membrane
389.03	Conductive hearing loss, middle ear
389.04	Conductive hearing loss, inner ear
389.05	Conductive hearing loss, unilateral
389.06	Conductive hearing loss, bilateral
389.08	Conductive hearing loss of combined types
389.13	Neural hearing loss, unilateral
389.15	Sensorineural hearing loss, unilateral
389.17	Sensory hearing loss, unilateral
389.20	Mixed hearing loss, unspecified
389.21	Mixed hearing loss, unilateral
389.22	Mixed hearing loss, bilateral
744.02	Other anomalies of external ear with impairment of hearing
744.03	Anomaly of middle ear, except ossicles

## ICD-10 Diagnosis Codes

<b>ICD-10-CM Diagnosis codes:</b>	<b>Code Description</b>
C30.1	Malignant neoplasm of middle ear
C44.201	Unspecified malignant neoplasm of skin of unspecified ear and external auricular canal
C44.202	Unspecified malignant neoplasm of skin of right ear and external auricular canal
C44.209	Unspecified malignant neoplasm of skin of left ear and external auricular canal
H60.399	Other infective otitis externa, unspecified ear
H60.60	Unspecified chronic otitis externa, unspecified ear
H60.61	Unspecified chronic otitis externa, right ear
H60.62	Unspecified chronic otitis externa, left ear
H60.63	Unspecified chronic otitis externa, bilateral
H60.8x1	Other otitis externa, right ear
H60.8x2	Other otitis externa, left ear
H60.8x3	Other otitis externa, bilateral
H60.8x9	Other otitis externa, unspecified ear

H60.90	Unspecified otitis externa, unspecified ear
H60.91	Unspecified otitis externa, right ear
H60.92	Unspecified otitis externa, left ear
H60.93	Unspecified otitis externa, bilateral
H61.391	Other acquired stenosis of right external ear canal
H61.392	Other acquired stenosis of left external ear canal
H61.393	Other acquired stenosis of external ear canal, bilateral
H61.399	Other acquired stenosis of external ear canal, unspecified ear
H62.8x1	Other disorders of right external ear in diseases classified elsewhere
H62.8x2	Other disorders of left external ear in diseases classified elsewhere
H62.8x3	Other disorders of external ear in diseases classified elsewhere, bilateral
H62.8x9	Other disorders of external ear in diseases classified elsewhere, unspecified ear
H65.20	Chronic serous otitis media, unspecified ear
H65.21	Chronic serous otitis media, right ear
H65.22	Chronic serous otitis media, left ear
H65.23	Chronic serous otitis media, bilateral
H65.30	Chronic mucoid otitis media, unspecified ear
H65.31	Chronic mucoid otitis media, right ear
H65.32	Chronic mucoid otitis media, left ear
H65.33	Chronic mucoid otitis media, bilateral
H65.411	Chronic allergic otitis media, right ear
H65.412	Chronic allergic otitis media, left ear
H65.413	Chronic allergic otitis media, bilateral
H65.419	Chronic allergic otitis media, unspecified ear
H65.491	Other chronic nonsuppurative otitis media, right ear
H65.492	Other chronic nonsuppurative otitis media, left ear
H65.493	Other chronic nonsuppurative otitis media, bilateral
H65.499	Other chronic nonsuppurative otitis media, unspecified ear
H66.10	Chronic tubotympanic suppurative otitis media, unspecified
H66.11	Chronic tubotympanic suppurative otitis media, right ear
H66.12	Chronic tubotympanic suppurative otitis media, left ear
H66.13	Chronic tubotympanic suppurative otitis media, bilateral
H66.20	Chronic atticoantral suppurative otitis media, unspecified ear
H66.21	Chronic atticoantral suppurative otitis media, right ear
H66.22	Chronic atticoantral suppurative otitis media, left ear
H66.23	Chronic atticoantral suppurative otitis media, bilateral
H66.3X1	Other chronic suppurative otitis media, right ear
H66.3X2	Other chronic suppurative otitis media, left ear
H66.3X3	Other chronic suppurative otitis media, bilateral
H66.3X9	Other chronic suppurative otitis media, unspecified ear
H90.0	Conductive hearing loss, bilateral
H90.11	Conductive hearing loss, unilateral, right ear, with unrestricted hearing on the contralateral side
H90.12	Conductive hearing loss, unilateral, left ear, with unrestricted hearing on the contralateral side
H90.2	Conductive hearing loss, unspecified
H90.41	Sensorineural hearing loss, unilateral, right ear, with unrestricted hearing on the contralateral side
H90.42	Sensorineural hearing loss, unilateral, left ear, with unrestricted hearing on the contralateral side
H90.6	Mixed conductive and sensorineural hearing loss, bilateral

H90.71	Mixed conductive and sensorineural hearing loss, unilateral, right ear, with unrestricted hearing on the contralateral side
H90.72	Mixed conductive and sensorineural hearing loss, unilateral, left ear, with unrestricted hearing on the contralateral side
H90.8	Mixed conductive and sensorineural hearing loss, unspecified
H90.A11	Conductive hearing loss, unilateral, right ear with restricted hearing on the contralateral side
H90.A12	Conductive hearing loss, unilateral, left ear with restricted hearing on the contralateral side
H90.A21	Sensorineural hearing loss, unilateral, right ear, with restricted hearing on the contralateral side
H90.A22	Sensorineural hearing loss, unilateral, left ear, with restricted hearing on the contralateral side
H90.A31	Mixed conductive and sensorineural hearing loss, unilateral, right ear with restricted hearing on the contralateral side
H90.A32	Mixed conductive and sensorineural hearing, unilateral, left ear with restricted hearing on the contralateral side
Q16.1	Congenital absence, atresia and stricture of auditory canal (external)
Q16.4	Other congenital malformations of middle ear

## Description

### Hearing Loss

Hearing loss is described as conductive, sensorineural, or mixed, and can be unilateral or bilateral. Normal hearing detects sound at or below 20 decibels (dB). The American Speech Language Hearing Association has defined degree of hearing loss based on pure-tone average detection thresholds as mild (20-40 dB), moderate (40-60 dB), severe (60-80 dB), and profound ( $\geq 80$  dB). Pure-tone average is calculated by averaging hearing sensitivities (ie, the minimum volume that a patient hears) at multiple frequencies (perceived as pitch), typically within the range of 0.25 to 8 kHz.

Sound amplification using an air-conduction (AC) hearing aid can provide benefit to patients with sensorineural or mixed hearing loss. Contralateral routing of signal (CROS) is a system in which a microphone on the affected side transmits a signal to an AC hearing aid on the normal or less affected side.

### Treatment

External bone-conduction hearing devices function by transmitting sound waves through the bone to the ossicles of the middle ear. The external devices must be applied close to the temporal bone, with either a steel spring over the top of the head or a spring-loaded arm on a pair of spectacles. These devices may be associated with pressure headaches or soreness.

A bone-anchored implant system combines a vibrational transducer coupled directly to the skull via a percutaneous abutment that permanently protrudes through the skin from a small titanium implant anchored in the temporal bone. The system is based on osseointegration through which living tissue integrates with titanium in the implant over 3 to 6 months, conducting amplified and processed sound via the skull bone directly to the cochlea. The lack of intervening skin permits the transmission of vibrations at a lower energy level than required for external bone-conduction hearing aids. Implantable bone-conduction hearing systems are primarily indicated for people with conductive or mixed sensorineural or conductive hearing loss. They may also be used with CROS as an alternative to an AC hearing aid for individuals with unilateral sensorineural hearing loss.

Partially implantable magnetic bone-conduction hearing systems also referred to as transcutaneous bone-anchored systems, are an alternative to bone-conduction hearing systems that connect to bone percutaneously via an abutment. With this technique, acoustic transmission occurs transcutaneously via magnetic coupling of the external sound processor and the internally implanted device components. The bone-conduction hearing processor contains magnets that adhere externally to magnets implanted in shallow bone beds with the bone-conduction hearing implant. Because the processor adheres

magnetically to the implant, there is no need for a percutaneous abutment to physically connect the external and internal components. To facilitate greater transmission of acoustics between magnets, skin thickness may be reduced to 4 to 5 mm over the implant when it is surgically placed.

## Summary

Sensorineural, conductive, and mixed hearing loss may be treated with various devices, including conventional air-conduction or bone-conduction external hearing aids. Air-conduction hearing aids may not be suitable for patients with chronic middle ear and ear canal infections, atresia of the external canal, or an ear canal that cannot accommodate an ear mold. Bone-conduction hearing aids may be useful for individuals with conductive hearing loss, or (if used with contralateral routing of signal), for unilateral sensorineural hearing loss. Implantable, bone-anchored hearing aids (BAHAs) that use a percutaneous or transcutaneous connection to a sound processor have been investigated as alternatives to conventional bone-conduction hearing aids for patients with conductive or mixed hearing loss or for patients with unilateral single-sided sensorineural hearing loss.

For individuals who have conductive or mixed hearing loss who receive an implantable BAHA with a percutaneous abutment or a partially implantable BAHA with transcutaneous coupling to the sound processor, the evidence includes observational studies that have reported pre-post differences in hearing parameters after treatment with BAHAs. Relevant outcomes are functional outcomes, quality of life, and treatment-related morbidity. No prospective trials were identified. Observational studies reporting on within-subjects changes in hearing have generally reported hearing improvements with the devices. Given the objectively measured outcomes and the largely invariable natural history of hearing loss in individuals who would be eligible for an implantable bone-conduction device, the demonstrated improvements in hearing after device placement can be attributed to the device. Studies of partially implantable BAHAs have similarly demonstrated within-subjects' improvements in hearing. The single-arm studies have shown improvements in hearing in the device-aided state. No direct comparisons other than within-individual comparisons with external hearing aids were identified, but, for individuals unable to wear an external hearing aid, there may be few alternative treatments. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have unilateral sensorineural hearing loss who receive a fully or partially implantable BAHA with the contralateral routing of signal, the evidence includes an RCT, multiple prospective and retrospective case series, and a systematic review. Relevant outcomes are functional outcomes, quality of life, and treatment-related morbidity. Single-arm case series, with sample sizes ranging from 9 to 180 patients, have generally reported improvements in patient-reported speech quality, speech perception in noise, and satisfaction with bone-conduction devices with contralateral routing of the signal. However, a well-conducted systematic review of studies comparing bone-anchored devices with hearing aids using contralateral routing of signal found no evidence of improvement in speech recognition or hearing localization. The single RCT included in the systematic review was a pilot study enrolling only 10 patients and, therefore, does not provide definitive evidence. Quality RCTs on BAHA for unilateral sensorineural hearing loss are lacking. The evidence is insufficient to determine the effects of the technology on health outcomes.

For patients with single-sided sensorineural deafness, a binaural hearing benefit may be provided by way of contralateral routing of signals to the hearing ear. There is evidence that bilateral hearing assistance devices improve hearing to a greater degree than unilateral devices. BAHAs may be considered an alternative to external devices in patients who are not candidates for external devices. By extension, the use of an implantable bone-conduction device with contralateral routing of the signal may be considered medically necessary in patients with unilateral sensorineural deafness.

## Policy History

Date	Action
4/2020	BCBSA National medical policy review. Description, summary and references updated. Policy statements unchanged.
4/2019	BCBSA National medical policy review. Description, summary and references updated. Policy statements unchanged.

3/2017	New references added from BCBSA National medical policy.
12/2016	BCBSA National medical policy review. Policy statements changed to remove investigational statement for partially implantable devices. References added. Clarified coding information Effective 12/1/2016.
8/2016	New medically necessary indications described for partially implantable bone conduction hearing systems using magnetic coupling for acoustic transmission. Effective 8/1/2016.
4/2016	New references added from BCBSA National medical policy.
2/2016	Clarified coding information.
3/2015	New references added from BCBSA National medical policy.
7/2014	Updated Coding section with ICD10 procedure and diagnosis codes, effective 10/2015.
3/2014	BCBSA National medical policy review. Investigational statement clarified. Effective 3/1/2014.
11/2013	Coding information clarified.
5/2013	New references from BCBSA National medical policy.
2/2013	New references from BCBSA National medical policy.
11/2011-4/2012	Medical policy ICD 10 remediation: Formatting, editing and coding updates. No changes to policy statements.
12/2011	BCBSA National medical policy review. Changes to policy statements.
9/2011	BCBSA National medical policy review. Changes to policy statements.
8/2010	BCBSA National medical policy review. Changes to policy statements.
5/2010	Reviewed - Medical Policy Group - Pediatrics and Endocrinology. No changes to policy statements.
3/2010	Reviewed - Medical Policy Group - Allergy and ENT/Otolaryngology. No changes to policy statements.
5/2009	Reviewed - Medical Policy Group - Pediatrics and Endocrinology. No changes to policy statements.
3/2009	Reviewed - Medical Policy Group - Allergy and ENT/Otolaryngology. No changes to policy statements.
1/2009	BCBSA National medical policy review. No changes to policy statements.
5/2008	Reviewed - Medical Policy Group - Pediatrics and Endocrinology. No changes to policy statements.
3/2008	Reviewed - Medical Policy Group - Allergy and ENT/Otolaryngology. No changes to policy statements.
11/2007	BCBSA National medical policy review. Changes to policy statements.
5/2007	Reviewed - Medical Policy Group - Pediatrics and Endocrinology. No changes to policy statements.
3/2007	Reviewed - Medical Policy Group - Allergy and ENT/Otolaryngology. No changes to policy statements.

## Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)



## References

1. Colquitt JL, Loveman E, Baguley DM, et al. Bone-anchored hearing aids for people with bilateral hearing impairment: a systematic review. *Clin Otolaryngol*. Oct 2011;36(5):419-441. PMID 21816006
2. Colquitt JL, Jones J, Harris P, et al. Bone-anchored hearing aids (BAHAs) for people who are bilaterally deaf: a systematic review and economic evaluation. *Health Technol Assess*. Jul 2011;15(26):1-200, iii-iv. PMID 21729632
3. Kompis M, Kurz A, Pfiffner F, et al. Is complex signal processing for bone conduction hearing aids useful? *Cochlear Implants Int*. May 2014;15 Suppl 1:S47-50. PMID 24869443
4. Hill-Feltham P, Roberts SA, Gladdis R. Digital processing technology for bone-anchored hearing aids: randomised comparison of two devices in hearing aid users with mixed or conductive hearing loss. *J Laryngol Otol*. Feb 2014;128(2):119-127. PMID 24524414
5. Farnoosh S, Mitsinikos FT, Maceri D, et al. Bone-anchored hearing aid vs. reconstruction of the external auditory canal in children and adolescents with congenital aural atresia: a comparison study of outcomes. *Front Pediatr*. Jan 2014;2:5. PMID 24479110
6. Ramakrishnan Y, Marley S, Leese D, et al. Bone-anchored hearing aids in children and young adults: the Freeman Hospital experience. *J Laryngol Otol*. Feb 2011;125(2):153-157. PMID 20849670
7. den Besten CA, Harterink E, McDermott AL, et al. Clinical results of Cochlear BIA300 in children: Experience in two tertiary referral centers. *Int J Pediatr Otorhinolaryngol*. Dec 2015;79(12):2050-2055. PMID 26455259
8. McLarnon CM, Davison T, Johnson IJ. Bone-anchored hearing aid: comparison of benefit by patient subgroups. *Laryngoscope*. May 2004;114(5):942-944. PMID 15126761
9. Tringali S, Grayeli AB, Bouccara D, et al. A survey of satisfaction and use among patients fitted with a BAHA. *Eur Arch Otorhinolaryngol*. Dec 2008;265(12):1461-1464. PMID 18415113
10. Snik AF, Mylanus EA, Cremers CW. The bone-anchored hearing aid compared with conventional hearing aids. Audiologic results and the patients' opinions. *Otolaryngol Clin North Am*. Feb 1995;28(1):73-83. PMID 7739870
11. van der Pouw CT, Snik AF, Cremers CW. The BAHA HC200/300 in comparison with conventional bone conduction hearing aids. *Clin Otolaryngol Allied Sci*. Jun 1999;24(3):171-176. PMID 10384840
12. Wazen JJ, Caruso M, Tjellstrom A. Long-term results with the titanium bone-anchored hearing aid: the U.S. experience. *Am J Otol*. Nov 1998;19(6):737-741. PMID 9831146
13. Granstrom G, Tjellstrom A. The bone-anchored hearing aid (BAHA) in children with auricular malformations. *Ear Nose Throat J*. Apr 1997;76(4):238-240, 242, 244-237. PMID 9127523
14. Janssen RM, Hong P, Chadha NK. Bilateral bone-anchored hearing aids for bilateral permanent conductive hearing loss: a systematic review. *Otolaryngol Head Neck Surg*. Sep 2012;147(3):412-422. PMID 22714424
15. Bosman AJ, Snik AF, van der Pouw CT, et al. Audiometric evaluation of bilaterally fitted bone-anchored hearing aids. *Audiology*. May-Jun 2001;40(3):158-167. PMID 11465298
16. Priwin C, Stenfelt S, Granstrom G, et al. Bilateral bone-anchored hearing aids (BAHAs): an audiometric evaluation. *Laryngoscope*. Jan 2004;114(1):77-84. PMID 14709999
17. Snik AF, Mylanus EA, Proops DW, et al. Consensus statements on the BAHA system: where do we stand at present? *Ann Otol Rhinol Laryngol Suppl*. Dec 2005;195:2-12. PMID 16619473
18. Dun CA, de Wolf MJ, Mylanus EA, et al. Bilateral bone-anchored hearing aid application in children: the Nijmegen experience from 1996 to 2008. *Otol Neurotol*. Jun 2010;31(4):615-623. PMID 20393374
19. Ho EC, Monksfield P, Egan E, et al. Bilateral bone-anchored hearing aid: impact on quality of life measured with the Glasgow Benefit Inventory. *Otol Neurotol*. Oct 2009;30(7):891-896. PMID 19692937
20. Briggs R, Van Hasselt A, Luntz M, et al. Clinical performance of a new magnetic bone conduction hearing implant system: results from a prospective, multicenter, clinical investigation. *Otol Neurotol*. Jun 2015;36(5):834-841. PMID 25634465
21. Denoyelle F, Coudert C, Thierry B, et al. Hearing rehabilitation with the closed skin bone-anchored implant Sophono Alpha1: results of a prospective study in 15 children with ear atresia. *Int J Pediatr Otorhinolaryngol*. Mar 2015;79(3):382-387. PMID 25617189
22. Hol MK, Nelissen RC, Agterberg MJ, et al. Comparison between a new implantable transcutaneous bone conductor and percutaneous bone-conduction hearing implant. *Otol Neurotol*. Aug 2013;34(6):1071-1075. PMID 23598702



23. Nelissen RC, Agterberg MJ, Hol MK, et al. Three-year experience with the Sophono in children with congenital conductive unilateral hearing loss: tolerability, audiometry, and sound localization compared to a bone-anchored hearing aid. *Eur Arch Otorhinolaryngol.* Oct 2016;273(10):3149-3156. PMID 26924741
24. Iseri M, Orhan KS, Tuncer U, et al. Transcutaneous bone-anchored hearing aids versus percutaneous ones: multicenter comparative clinical study. *Otol Neurotol.* Jun 2015;36(5):849-853. PMID 25730451
25. Gerdes T, Salcher RB, Schwab B, et al. Comparison of audiological results between a transcutaneous and a percutaneous bone conduction instrument in conductive hearing loss. *Otol Neurotol.* Jul 2016;37(6):685-691. PMID 27093021
26. Dimitriadis PA, Farr MR, Allam A, et al. Three year experience with the cochlear Baha attract implant: a systematic review of the literature. *BMC Ear Nose Throat Disord.* Oct 2016;16:12. PMID 27733813
27. Reddy-Kolanu R, Gan R, Marshall AH. A case series of a magnetic bone conduction hearing implant. *Ann R Coll Surg Engl.* Nov 2016;98(8):552-553. PMID 27490984
28. Siegert R. Partially implantable bone conduction hearing aids without a percutaneous abutment (Otomag): technique and preliminary clinical results. *Adv Otorhinolaryngol.* 2011;71:41-46. PMID 21389703
29. Powell HR, Rolfe AM, Birman CS. A comparative study of audiological outcomes for two transcutaneous bone- anchored hearing devices. *Otol Neurotol.* Sep 2015;36(9):1525-1531. PMID 26375976
30. O'Neil MB, Runge CL, Friedland DR, et al. Patient outcomes in magnet-based implantable auditory assist devices. *JAMA Otolaryngol Head Neck Surg.* Jun 2014;140(6):513-520. PMID 24763485
31. Centric A, Chennupati SK. Abutment-free bone-anchored hearing devices in children: initial results and experience. *Int J Pediatr Otorhinolaryngol.* May 2014;78(5):875-878. PMID 24612554
32. Baker S, Centric A, Chennupati SK. Innovation in abutment-free bone-anchored hearing devices in children: Updated results and experience. *Int J Pediatr Otorhinolaryngol.* Oct 2015;79(10):1667-1672. PMID 26279245
33. Marsella P, Scorpecci A, Vallarino MV, et al. Sophono in pediatric patients: the experience of an Italian tertiary care center. *Otolaryngol Head Neck Surg.* Apr 8 2014;151(2):328-332. PMID 24714216
34. Bravo-Torres S, Der-Mussa C, Fuentes-Lopez E. Active transcutaneous bone conduction implant: audiological results in paediatric patients with bilateral microtia associated with external auditory canal atresia. *Int J Audiol.* 2018 Jan;57(1). PMID 28857620
35. Schmerber S, Deguine O, Marx M, et al. Safety and effectiveness of the Bonebridge transcutaneous active direct-drive bone-conduction hearing implant at 1-year device use. *Eur Arch Otorhinolaryngol.* Apr 2017;274(4):1835-1851. PMID 27475796
36. Rahne T, Seiwerth I, Gotze G, et al. Functional results after Bonebridge implantation in adults and children with conductive and mixed hearing loss. *Eur Arch Otorhinolaryngol.* Nov 2015;272(11):3263-3269. PMID 25425039
37. Laske RD, Roosli C, Pfiffner F, et al. Functional results and subjective benefit of a transcutaneous bone conduction device in patients with single-sided deafness. *Otol Neurotol.* Aug 2015;36(7):1151-1156. PMID 26111077
38. Riss D, Arnoldner C, Baumgartner WD, et al. Indication criteria and outcomes with the Bonebridge transcutaneous bone-conduction implant. *Laryngoscope.* Dec 2014;124(12):2802-2806. PMID 25142577
39. Manrique M, Sanhueza I, Manrique R, et al. A new bone conduction implant: surgical technique and results. *Otol Neurotol.* Feb 2014;35(2):216-220. PMID 24448280
40. Ihler F, Volbers L, Blum J, et al. Preliminary functional results and quality of life after implantation of a new bone conduction hearing device in patients with conductive and mixed hearing loss. *Otol Neurotol.* Feb 2014;35(2):211-215. PMID 24448279
41. Desmet J, Wouters K, De Bodt M, et al. Long-term subjective benefit with a bone conduction implant sound processor in 44 patients with single-sided deafness. *Otol Neurotol.* Jul 2014;35(6):1017-1025. PMID 24751733
42. Iseri M, Orhan KS, Kara A, et al. A new transcutaneous bone anchored hearing device - the Baha(R) Attract System: the first experience in Turkey. *Kulak Burun Bogaz Ihtis Derg.* Mar-Apr 2014;24(2):59-64. PMID 24835899

43. Peters JP, Smit AL, Stegeman I, et al. Review: Bone conduction devices and contralateral routing of sound systems in single-sided deafness. *Laryngoscope*. Jan 2015;125(1):218-226. PMID 25124297
44. Baguley DM, Bird J, Humphriss RL, et al. The evidence base for the application of contralateral bone anchored hearing aids in acquired unilateral sensorineural hearing loss in adults. *Clin Otolaryngol*. Feb 2006;31(1):6-14. PMID 16441794
45. den Besten CA, Monksfield P, Bosman A et al. Audiological and clinical outcomes of a transcutaneous bone conduction hearing implant: Six-month results from a multicentre study. *Clin Otolaryngol*. 2019 Mar;44(2). PMID 30358920
46. Leterme G, Bernardeschi D, Bensemman A, et al. Contralateral routing of signal hearing aid versus transcutaneous bone conduction in single-sided deafness. *Audiol Neurootol*. 2015;20(4):251-260. PMID 26021779
47. Snapp HA, Holt FD, Liu X, et al. Comparison of speech-in-noise and localization benefits in unilateral hearing loss subjects using contralateral routing of signal hearing aids or bone-anchored implants. *Otol Neurotol*. Jan 2017;38(1):11-18. PMID 27846038
48. Zeitler DM, Snapp HA, Telischi FF, et al. Bone-anchored implantation for single-sided deafness in patients with less than profound hearing loss. *Otolaryngol Head Neck Surg*. Jul 2012;147(1):105-111. PMID 22368043
49. Pai I, Kelleher C, Nunn T, et al. Outcome of bone-anchored hearing aids for single-sided deafness: a prospective study. *Acta Otolaryngol*. Jul 2012;132(7):751-755. PMID 22497318
50. Nicolas S, Mohamed A, Yoann P, et al. Long-term benefit and sound localization in patients with single-sided deafness rehabilitated with an osseointegrated bone-conduction device. *Otol Neurotol*. Jan 2013;34(1):111-114. PMID 23202156
51. Lin LM, Bowditch S, Anderson MJ, et al. Amplification in the rehabilitation of unilateral deafness: speech in noise and directional hearing effects with bone-anchored hearing and contralateral routing of signal amplification. *Otol Neurotol*. Feb 2006;27(2):172-182. PMID 16436986
52. Kunst SJ, Leijendeckers JM, Mylanus EA, et al. Bone-anchored hearing aid system application for unilateral congenital conductive hearing impairment: audiometric results. *Otol Neurotol*. Jan 2008;29(1):2-7. PMID 18199951
53. Kunst SJ, Hol MK, Mylanus EA, et al. Subjective benefit after BAHA system application in patients with congenital unilateral conductive hearing impairment. *Otol Neurotol*. Apr 2008;29(3):353-358. PMID 18494142
54. Gluth MB, Eager KM, Eikelboom RH, et al. Long-term benefit perception, complications, and device malfunction rate of bone-anchored hearing aid implantation for profound unilateral sensorineural hearing loss. *Otol Neurotol*. Dec 2010;31(9):1427-1434. PMID 20729779
55. Faber HT, Nelissen RC, Kramer SE, et al. Bone-anchored hearing implants in single-sided deafness patients: Long-term use and satisfaction by gender. *Laryngoscope*. Dec 2015;125(12):2790-2795. PMID 26152833
56. Monini S, Musy I, Filippi C, et al. Bone conductive implants in single-sided deafness. *Acta Otolaryngol*. Apr 2015;135(4):381-388. PMID 25720582
57. Amonoo-Kuofi K, Kelly A, Neeff M, et al. Experience of bone-anchored hearing aid implantation in children younger than 5 years of age. *Int J Pediatr Otorhinolaryngol*. Apr 2015;79(4):474-480. PMID 25680294
58. Marsella P, Scorpecci A, Pacifico C, et al. Pediatric BAHA in Italy: the "Bambino Gesù" Children's Hospital's experience. *Eur Arch Otorhinolaryngol*. Feb 2012;269(2):467-474. PMID 21739094
59. Davids T, Gordon KA, Clutton D, et al. Bone-anchored hearing aids in infants and children younger than 5 years. *Arch Otolaryngol Head Neck Surg*. Jan 2007;133(1):51-55. PMID 17224524
60. McDermott AL, Williams J, Kuo MJ, et al. The role of bone anchored hearing aids in children with Down syndrome. *Int J Pediatr Otorhinolaryngol*. Jun 2008;72(6):751-757. PMID 18433885
61. Verheij E, Bezdjian A, Grolman W, et al. A systematic review on complications of tissue preservation surgical techniques in percutaneous bone conduction hearing devices. *Otol Neurotol*. Aug 2016;37(7):829-837. PMID 27273402
62. Kiringoda R, Lustig LR. A meta-analysis of the complications associated with osseointegrated hearing aids. *Otol Neurotol*. Jul 2013;34(5):790-794. PMID 23739555
63. Dun CA, Faber HT, de Wolf MJ, et al. Assessment of more than 1,000 implanted percutaneous bone conduction devices: skin reactions and implant survival. *Otol Neurotol*. Feb 2012;33(2):192-198. PMID 22246385

64. Hobson JC, Roper AJ, Andrew R, et al. Complications of bone-anchored hearing aid implantation. *J Laryngol Otol*. Feb 2010;124(2):132-136. PMID 19968889
65. Wallberg E, Granstrom G, Tjellstrom A, et al. Implant survival rate in bone-anchored hearing aid users: long-term results. *J Laryngol Otol*. Nov 2011;125(11):1131-1135. PMID 21774847
66. Kraai T, Brown C, Neeff M, et al. Complications of bone-anchored hearing aids in pediatric patients. *Int J Pediatr Otorhinolaryngol*. Jun 2011;75(6):749-753. PMID 21470698
67. Allis TJ, Owen BD, Chen B, et al. Longer length Baha abutments decrease wound complications and revision surgery. *Laryngoscope*. Apr 2014;124(4):989-992. PMID 24114744
68. Calvo Bodnia N, Foghsgaard S, Nue Moller M, et al. Long-term results of 185 consecutive osseointegrated hearing device implantations: a comparison among children, adults, and elderly. *Otol Neurotol*. Dec 2014;35(10):e301-306. PMID 25122598
69. Rebol J. Soft tissue reactions in patients with bone anchored hearing aids. *Ir J Med Sci*. Jun 2015;184(2):487- 491. PMID 24913737
70. Larsson A, Tjellstrom A, Stafors J. Implant losses for the bone-anchored hearing devices are more frequent in some patients. *Otol Neurotol*. Feb 2015;36(2):336-340. PMID 24809279
71. den Besten CA, Nelissen RC, Peer PG, et al. A retrospective cohort study on the influence of comorbidity on soft tissue reactions, revision surgery, and implant loss in bone-anchored hearing implants. *Otol Neurotol*. Jun 2015;36(5):812-818. PMID 25811351
72. Mohamad S, Khan I, Hey SY, et al. A systematic review on skin complications of bone-anchored hearing aids in relation to surgical techniques. *Eur Arch Otorhinolaryngol*. Mar 2016;273(3):559-565. PMID 25503356
73. Fontaine N, Hemar P, Schultz P, et al. BAHA implant: implantation technique and complications. *Eur Ann Otorhinolaryngol Head Neck Dis*. Feb 2014;131(1):69-74. PMID 23835074
74. Hultcrantz M, Lanis A. A five-year follow-up on the osseointegration of bone-anchored hearing device implantation without tissue reduction. *Otol Neurotol*. Sep 2014;35(8):1480-1485. PMID 24770406
75. Nelissen RC, Stafors J, de Wolf MJ, et al. Long-term stability, survival, and tolerability of a novel osseointegrated implant for bone conduction hearing: 3-year data from a multicenter, randomized, controlled, clinical investigation. *Otol Neurotol*. Sep 2014;35(8):1486-1491. PMID 25080037
76. Singam S, Williams R, Saxby C, et al. Percutaneous bone-anchored hearing implant surgery without soft-tissue reduction: up to 42 months of follow-up. *Otol Neurotol*. Oct 2014;35(9):1596-1600. PMID 25076228
77. Roplekar R, Lim A, Hussain SS. Has the use of the linear incision reduced skin complications in bone-anchored hearing aid implantation? *J Laryngol Otol*. Jun 2016;130(6):541-544. PMID 27160014
78. American Academy of Otolaryngology-Head and Neck Surgery. Position Statement: Bone Conduction Hearing Devices. Position Statements 2016; <http://www.entnet.org/content/position-statement-bone-conduction-hearing-devices>. Accessed December 18, 2019.
79. Centers for Medicare & Medicaid Services. Medicare Policy Benefit Manual. Chapter 16 - General Exclusions from Coverage (Rev. 198). 2014; Rev. 189;<http://www.cms.gov/manuals/Downloads/bp102c16.pdf>. Accessed December 18, 2019.
80. Centers for Medicare & Medicaid Services. Fact sheets: CMS Updates Policies and Payment Rates for End- Stage Renal Disease Facilities for CY 2015 and Implementation of Competitive Bidding-Based Prices for Durable Medical Equipment, Prosthetics, Orthotics, and Supplies. 2014; <https://www.cms.gov/newsroom/fact-sheets/cms-updates-policies-and-payment-rates-end-stage-renal-disease-facilities-cy-2015-and-implementation>. Accessed December 18, 2019.