



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

Medical Policy

Ultrasonographic Measurement of Carotid Intima-Medial Thickness as an Assessment of Subclinical Atherosclerosis

Table of Contents

- [Policy: Commercial](#)
- [Coding Information](#)
- [Information Pertaining to All Policies](#)
- [Policy: Medicare](#)
- [Description](#)
- [References](#)
- [Authorization Information](#)
- [Policy History](#)

Policy Number: 547

BCBSA Reference Number: 2.02.16

NCD/LCD: National Coverage Determination (NCD) for Ultrasound Diagnostic Procedures (220.5)

Related Policies

- Novel Lipid Risk Factors in Risk Assessment and Management of Cardiovascular Disease, [#283](#)

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity

Ultrasonographic measurement of carotid artery intima-medial thickness (CIMT) as a technique of identifying subclinical atherosclerosis is **INVESTIGATIONAL** for use in the screening, diagnosis, or management of atherosclerotic disease.

Medicare HMO BlueSM and Medicare PPO BlueSM Members

BCBSMA does not cover ultrasound diagnostic procedures for the B-Scan for atherosclerotic narrowing of peripheral arteries for Medicare HMO Blue and Medicare PPO Blue members in accordance with CMS NCD.

Medicare HMO BlueSM and Medicare PPO BlueSM Members

Medical necessity criteria and coding guidance can be found through the link(s) below.

[National Coverage Determinations \(NCDs\)](#)

National Coverage Determination (NCD) for Ultrasound Diagnostic Procedures (220.5)

Note: To review the specific NCD, please remember to click “accept” on the CMS licensing agreement at the bottom of the CMS webpage.

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	This is not a covered service.
Commercial PPO and Indemnity	This is not a covered service.
Medicare HMO Blue SM	This is not a covered service.
Medicare PPO Blue SM	This is not a covered service.

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

The following CPT codes are considered investigational for Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:

CPT Codes

CPT codes:	Code Description
93895	Quantitative carotid intima media thickness and carotid atheroma evaluation, bilateral
0126T	Common carotid intima-media thickness (IMT) study for evaluation of atherosclerotic burden or coronary heart disease risk factor assessment

Description

Coronary Heart Disease

Coronary heart disease (CHD) accounts for 30.8% of all deaths in the United States.¹ Established major risk factors for CHD have been identified by the National Cholesterol Education Program Expert Panel. These risk factors include elevated serum levels of low-density lipoprotein cholesterol and total cholesterol, and reduced levels of high-density lipoprotein cholesterol. Other risk factors include a history of cigarette smoking, hypertension, family history of premature CHD, and age.

Diagnosis

The third report of the National Cholesterol Education Program Adult Treatment Panel established various treatment strategies to modify the risk of CHD, with emphasis on target goals of low-density lipoprotein cholesterol. Pathology studies have demonstrated that levels of traditional risk factors are associated with the extent and severity of atherosclerosis. The third report of the National Cholesterol Education Program Adult Treatment Panel recommended use of the Framingham criteria to further stratify those patients with 2 or more risk factors for more intensive lipid management.² However, at every level of risk factor exposure, there is substantial variation in the amount of atherosclerosis, presumably related to genetic susceptibility and the influence of other risk factors. Thus, there has been interest in identifying a technique that can improve the ability to diagnose those at risk of developing CHD, as well as to measure disease progression, particularly for those at intermediate risk.

The carotid arteries can be well-visualized by ultrasonography, and ultrasonographic measurement of the carotid intima-media thickness (IMT) has been investigated as a technique to identify and monitor subclinical atherosclerosis. B-mode ultrasound is most commonly used to measure the carotid IMT. IMT is measured and averaged over several sites in each carotid artery. Imaging the far wall of each common carotid artery yields more accurate and reproducible IMT measurements than imaging near wall. Two echogenic lines are produced, representing the lumen-intima interface and the media-adventitia interface. The distance between these 2 lines constitutes the IMT.

Summary

Ultrasonographic measurement of carotid intima-media (or intimal-medial) thickness (CIMT) refers to the use of B-mode ultrasound to determine the thickness of the 2 innermost layers of the carotid artery wall, the intima and the media. Detection and monitoring of intima-media thickening, which is a surrogate marker for atherosclerosis, may provide an opportunity to intervene earlier in atherogenic disease and/or monitor disease progression.

For individuals who are undergoing cardiac risk assessment who receive ultrasonic measurement of CIMT, the evidence includes a randomized controlled study, large cohort studies, case-control studies, and systematic reviews. Relevant outcomes are test accuracy and morbid events. Some studies have correlated increased CIMT with other commonly used markers for risk of coronary heart disease and with risk for future cardiovascular events. Lorenz et al (2012) found in their meta-analysis that CIMT was associated with increased cardiovascular events, although CIMT progression over time was not associated with increased cardiovascular event risk. Peters et al (2012) found that the added predictive value of CIMT was modest, and the ability to reclassify patients into clinically relevant categories was not demonstrated. The results from these reviews and other studies have demonstrated the predictive value of CIMT is uncertain and that the predictive ability for any level of population risk cannot be determined with precision. Also, available studies do not define how the use of CIMT in clinical practice improves outcomes. There is no scientific literature that directly tests the hypothesis that measurement of CIMT results in improved patient outcomes and no specific guidance on how measurements of CIMT should be incorporated into risk assessment and risk management. The objective of one study, however, was to define “normal” CIMT progression in low to moderate cardiovascular risk patients. Study results showed definite patterns related to various factors that could be used as a tool to earlier identify patients at increased cardiovascular risk, but patient outcomes were not assessed. The evidence is insufficient to determine the effects of the technology on health outcomes.

Policy History

Date	Action
7/2020	BCBSA National medical policy review. Description, summary and references updated. Policy statements unchanged.
6/2019	BCBSA National medical policy review. Description, summary and references updated. Policy statements unchanged.
6/2018	New references added from BCBSA National medical policy. Background and summary clarified.
2/2017	New references added from BCBSA National medical policy.
8/2015	New references added from BCBSA National medical policy.
1/2015	Clarified coding information.
9/2014	New references added from BCBSA National medical policy.
10/2013	New references from BCBSA National medical policy.
11/2011-4/2012	Medical policy ICD 10 remediation: Formatting, editing and coding updates. No changes to policy statements.
1/1/2012	New policy, effective 01/01/2012, describing ongoing non-coverage.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

References

1. Mozaffarian D, Benjamin EJ, Go AS, et al. Heart Disease and Stroke Statistics-2016 Update: A Report From the American Heart Association. *Circulation*. Jan 26 2016; 133(4): e38-360. PMID 26673558
2. Pasternak RC. Report of the Adult Treatment Panel III: the 2001 National Cholesterol Education Program guidelines on the detection, evaluation and treatment of elevated cholesterol in adults. *Cardiol Clin*. Aug 2003; 21(3): 393-8. PMID 14621453
3. Mookadam F, Moustafa SE, Lester SJ, et al. Subclinical atherosclerosis: evolving role of carotid intima-media thickness. *Prev Cardiol*. 2010; 13(4): 186-97. PMID 20860643
4. Den Ruijter HM, Peters SA, Anderson TJ, et al. Common carotid intima-media thickness measurements in cardiovascular risk prediction: a meta-analysis. *JAMA*. Aug 22 2012; 308(8): 796-803. PMID 22910757
5. Lorenz MW, Polak JF, Kavousi M, et al. Carotid intima-media thickness progression to predict cardiovascular events in the general population (the PROG-IMT collaborative project): a meta-analysis of individual participant data. *Lancet*. Jun 02 2012; 379(9831): 2053-62. PMID 22541275
6. van den Oord SC, Sijbrands EJ, ten Kate GL, et al. Carotid intima-media thickness for cardiovascular risk assessment: systematic review and meta-analysis. *Atherosclerosis*. May 2013; 228(1): 1-11. PMID 23395523
7. Plichart M, Celermajer DS, Zureik M, et al. Carotid intima-media thickness in plaque-free site, carotid plaques and coronary heart disease risk prediction in older adults. The Three-City Study. *Atherosclerosis*. Dec 2011; 219(2): 917-24. PMID 22005196
8. Keo HH, Baumgartner I, Hirsch AT, et al. Carotid plaque and intima-media thickness and the incidence of ischemic events in patients with atherosclerotic vascular disease. *Vasc Med*. Oct 2011; 16(5): 323-30. PMID 21908682
9. Nambi V, Chambless L, He M, et al. Common carotid artery intima-media thickness is as good as carotid intima-media thickness of all carotid artery segments in improving prediction of coronary heart disease risk in the Atherosclerosis Risk in Communities (ARIC) study. *Eur Heart J*. Jan 2012; 33(2): 183-90. PMID 21666250
10. Xie W, Liang L, Zhao L, et al. Combination of carotid intima-media thickness and plaque for better predicting risk of ischaemic cardiovascular events. *Heart*. Aug 2011; 97(16): 1326-31. PMID 21653216
11. Peters SA, den Ruijter HM, Bots ML, et al. Improvements in risk stratification for the occurrence of cardiovascular disease by imaging subclinical atherosclerosis: a systematic review. *Heart*. Feb 2012; 98(3): 177-84. PMID 22095617
12. Dobs AS, Nieto FJ, Szklo M, et al. Risk factors for popliteal and carotid wall thicknesses in the Atherosclerosis Risk in Communities (ARIC) Study. *Am J Epidemiol*. Nov 15 1999; 150(10): 1055-67. PMID 10568620
13. Chambless LE, Heiss G, Folsom AR, et al. Association of coronary heart disease incidence with carotid arterial wall thickness and major risk factors: the Atherosclerosis Risk in Communities (ARIC) Study, 1987-1993. *Am J Epidemiol*. Sep 15 1997; 146(6): 483-94. PMID 9290509
14. van der Meer IM, Bots ML, Hofman A, et al. Predictive value of noninvasive measures of atherosclerosis for incident myocardial infarction: the Rotterdam Study. *Circulation*. Mar 09 2004; 109(9): 1089-94. PMID 14993130
15. O'Leary DH, Polak JF, Kronmal RA, et al. Carotid-artery intima and media thickness as a risk factor for myocardial infarction and stroke in older adults. Cardiovascular Health Study Collaborative Research Group. *N Engl J Med*. Jan 07 1999; 340(1): 14-22. PMID 9878640
16. Lorenz MW, Schaefer C, Steinmetz H, et al. Is carotid intima media thickness useful for individual prediction of cardiovascular risk? Ten-year results from the Carotid Atherosclerosis Progression Study (CAPS). *Eur Heart J*. Aug 2010; 31(16): 2041-8. PMID 20530503
17. Folsom AR, Kronmal RA, Detrano RC, et al. Coronary artery calcification compared with carotid intima-media thickness in the prediction of cardiovascular disease incidence: the Multi-Ethnic Study of Atherosclerosis (MESA). *Arch Intern Med*. Jun 23 2008; 168(12): 1333-9. PMID 18574091

18. Paramsothy P, Knopp RH, Bertoni AG, et al. Association of combinations of lipid parameters with carotid intima-media thickness and coronary artery calcium in the MESA (Multi-Ethnic Study of Atherosclerosis). *J Am Coll Cardiol*. Sep 21 2010; 56(13): 1034-41. PMID 20846602
19. Blaha MJ, Rivera JJ, Budoff MJ, et al. Association between obesity, high-sensitivity C-reactive protein 2 mg/L, and subclinical atherosclerosis: implications of JUPITER from the Multi-Ethnic Study of Atherosclerosis. *Arterioscler Thromb Vasc Biol*. Jun 2011; 31(6): 1430-8. PMID 21474823
20. Patel J, Al Rifai M, Blaha MJ, et al. Coronary Artery Calcium Improves Risk Assessment in Adults With a Family History of Premature Coronary Heart Disease: Results From Multiethnic Study of Atherosclerosis. *Circ Cardiovasc Imaging*. Jun 2015; 8(6): e003186. PMID 26047825
21. Camhi SM, Katzmarzyk PT, Broyles ST, et al. Subclinical atherosclerosis and metabolic risk: role of body mass index and waist circumference. *Metab Syndr Relat Disord*. Apr 2011; 9(2): 119-25. PMID 21133775
22. Green D, Foiles N, Chan C, et al. An association between clotting factor VII and carotid intima-media thickness: the CARDIA study. *Stroke*. Jul 2010; 41(7): 1417-22. PMID 20466994
23. Bots ML, Palmer MK, Dogan S, et al. Intensive lipid lowering may reduce progression of carotid atherosclerosis within 12 months of treatment: the METEOR study. *J Intern Med*. Jun 2009; 265(6): 698-707. PMID 19298496
24. Baber U, Mehran R, Sartori S, et al. Prevalence, impact, and predictive value of detecting subclinical coronary and carotid atherosclerosis in asymptomatic adults: the Biolmage study. *J Am Coll Cardiol*. Mar 24 2015; 65(11): 1065-74. PMID 25790876
25. Geisel MH, Bauer M, Hennig F, et al. Comparison of coronary artery calcification, carotid intima-media thickness and ankle-brachial index for predicting 10-year incident cardiovascular events in the general population. *Eur Heart J*. Jun 14 2017; 38(23): 1815-1822. PMID 28379333
26. Villines TC, Hsu LL, Blackshear C, et al. Relation of Carotid Intima-Media Thickness to Cardiovascular Events in Black Americans (From the Jackson Heart Study). *Am J Cardiol*. Nov 01 2017; 120(9): 1528-1532. PMID 28844515
27. Johnson HM, Turke TL, Grossklaus M, et al. Effects of an office-based carotid ultrasound screening intervention. *J Am Soc Echocardiogr*. Jul 2011; 24(7): 738-47. PMID 21477989
28. Olmastroni E, Baragetti A, Casula M, et al. Multilevel Models to Estimate Carotid Intima-Media Thickness Curves for Individual Cardiovascular Risk Evaluation. *Stroke*. Jul 2019; 50(7): 1758-1765. PMID 31164073
29. Goff DC, Lloyd-Jones DM, Bennett G, et al. 2013 ACC/AHA guideline on the assessment of cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation*. Jun 24 2014; 129(25 Suppl 2): S49-73. PMID 24222018
30. Greenland P, Alpert JS, Beller GA, et al. 2010 ACCF/AHA guideline for assessment of cardiovascular risk in asymptomatic adults: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. Dec 14 2010; 56(25): e50-103. PMID 21144964
31. Jellinger PS, Handelsman Y, Rosenblit PD, et al. AMERICAN ASSOCIATION OF CLINICAL ENDOCRINOLOGISTS AND AMERICAN COLLEGE OF ENDOCRINOLOGY GUIDELINES FOR MANAGEMENT OF DYSLIPIDEMIA AND PREVENTION OF CARDIOVASCULAR DISEASE - EXECUTIVE SUMMARY Complete Appendix to Guidelines available at journals.aace.com. *Endocr Pract*. Apr 02 2017; 23(4): 479-497. PMID 28156151
32. Stein JH, Korcarz CE, Hurst RT, et al. Use of carotid ultrasound to identify subclinical vascular disease and evaluate cardiovascular disease risk: a consensus statement from the American Society of Echocardiography Carotid Intima-Media Thickness Task Force. Endorsed by the Society for Vascular Medicine. *J Am Soc Echocardiogr*. Feb 2008; 21(2): 93-111; quiz 189-90. PMID 18261694
33. Calonge N, Petitti DB, DeWitt TG, et al. Using nontraditional risk factors in coronary heart disease risk assessment: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. Oct 06 2009; 151(7): 474-82. PMID 19805770
34. Curry SJ, Krist AH, Owens DK, et al. Risk Assessment for Cardiovascular Disease With Nontraditional Risk Factors: US Preventive Services Task Force Recommendation Statement. *JAMA*. Jul 17 2018; 320(3): 272-280. PMID 29998297