



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an independent licensee of the Blue Cross and Blue Shield Association

Medical Policy Hippotherapy

Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

Policy Number: 560

BCBSA Reference Number: 8.03.12

NCD/LCD: N/A

Related Policies

None

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO BlueSM and Medicare PPO BlueSM Members

Hippotherapy is considered [INVESTIGATIONAL](#).

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	This is not a covered service.
Commercial PPO and Indemnity	This is not a covered service.
Medicare HMO BlueSM	This is not a covered service.
Medicare PPO BlueSM	This is not a covered service.

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

The following HCPCS code is considered investigational for **Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:**

HCPCS Codes

HCPCS codes:	Code Description
S8940	Equestrian/hippotherapy, per session

Description

Hippotherapy

Hippotherapy has been proposed as a technique to decrease the energy requirements and improve walking in patients with cerebral palsy. It is thought that the natural swaying motion of the horse induces a pelvic movement in the rider that simulates human ambulation. Also, variations in the horse's movements can prompt natural equilibrium movements in the rider.

Hippotherapy is also being evaluated in patients with multiple sclerosis and other causes of gait disorders, such as strokes.

As a therapeutic intervention, hippotherapy is typically conducted by a physical or occupational therapist and is aimed at improving impaired body function. Therapeutic horseback riding is typically conducted by riding instructors and is more frequently intended as social therapy. It is hoped that the multisensory environment may benefit children with profound social and communication deficits, such as autism spectrum disorder and schizophrenia. When considered together, hippotherapy and therapeutic riding are described as equine-assisted activities and therapies.

This evidence review addresses equine-assisted activities that focus on improving physical functions such as balance and gait.

Summary

Hippotherapy, also referred to as equine-assisted therapy, describes a treatment strategy that uses equine movement to engage sensory, neuromotor, and cognitive systems to achieve functional outcomes. Hippotherapy has been proposed as a therapy for patients with impaired walking or balance.

For individuals who have cerebral palsy, multiple sclerosis, stroke, or gait and balance disorders other than cerebral palsy, multiple sclerosis, and stroke who receive hippotherapy, the evidence includes systematic reviews, randomized trials, and case series. Relevant outcomes include symptoms and functional outcomes. Studies in cerebral palsy, multiple sclerosis, stroke, and other indications have had variable findings. The randomized trials are generally small and have significant methodologic problems. In the largest randomized trial conducted to date (72 children), which had blinding outcome assessment, hippotherapy had no clinically significant impact on children with cerebral palsy. There are no randomized controlled trials showing that hippotherapy is superior to alternative treatments for patients with multiple sclerosis. Hippotherapy for other indications has been compared primarily with no intervention and, although some benefits have been seen, it has not been shown to be more effective than other active therapies. The evidence is insufficient to determine the effects of the technology on health outcomes.

Policy History

Date	Action
5/2020	BCBSA National medical policy review. Description, summary and references updated. Policy statements unchanged.

4/2019	BCBSA National medical policy review. Description, summary and references updated. Policy statements unchanged.
4/2016	New references added from BCBSA National medical policy.
12/2014	New references added from BCBSA National medical policy.
11/2014	Medical policy ICD 10 remediation: Formatting, editing and coding updates. No changes to policy statements. Language transferred from Medical Policy 178, Complementary Medicine

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

References

1. Tseng SH, Chen HC, Tam KW. Systematic review and meta-analysis of the effect of equine assisted activities and therapies on gross motor outcome in children with cerebral palsy. *Disabil Rehabil*. Jan 2013;35(2):89-99. PMID 22630812
2. Zadnikar M, Kastrin A. Effects of hippotherapy and therapeutic horseback riding on postural control or balance in children with cerebral palsy: a meta-analysis. *Dev Med Child Neurol*. Aug 2011;53(8):684-691. PMID 21729249
3. Davis E, Davies B, Wolfe R, et al. A randomized controlled trial of the impact of therapeutic horse riding on the quality of life, health, and function of children with cerebral palsy. *Dev Med Child Neurol*. Feb 2009;51(2):111- 119; discussion 188. PMID 19191844
4. McGibbon NH, Benda W, Duncan BR, et al. Immediate and long-term effects of hippotherapy on symmetry of adductor muscle activity and functional ability in children with spastic cerebral palsy. *Arch Phys Med Rehabil*. Jun 2009;90(6):966-974. PMID 19480872
5. Benda W, McGibbon NH, Grant KL. Improvements in muscle symmetry in children with cerebral palsy after equine-assisted therapy (hippotherapy). *J Altern Complement Med*. Dec 2003;9(6):817-825. PMID 14736353
6. Kwon JY, Chang HJ, Yi SH, et al. Effect of hippotherapy on gross motor function in children with cerebral palsy: a randomized controlled trial. *J Altern Complement Med*. Jan 2015;21(1):15-21. PMID 25551626
7. Bronson C, Brewerton K, Ong J, et al. Does hippotherapy improve balance in persons with multiple sclerosis: a systematic review. *Eur J Phys Rehabil Med*. Sep 2010;46(3):347-353. PMID 20927000
8. Silkwood-Sherer D, Warmbier H. Effects of hippotherapy on postural stability, in persons with multiple sclerosis: a pilot study. *J Neurol Phys Ther*. Jun 2007;31(2):77-84. PMID 17558361
9. Munoz-Lasa S, Ferriero G, Valero R, et al. Effect of therapeutic horseback riding on balance and gait of people with multiple sclerosis. *G Ital Med Lav Ergon*. Oct-Dec 2011;33(4):462-467. PMID 22452106
10. Frevel D, Maurer M. Internet-based home training is capable to improve balance in multiple sclerosis: a randomized controlled trial. *Eur J Phys Rehabil Med*. Feb 2015;51(1):23-30. PMID 24755773
11. Lee CW, Kim SG, Yong MS. Effects of hippotherapy on recovery of gait and balance ability in patients with stroke. *J Phys Ther Sci*. Feb 2014;26(2):309-311. PMID 24648655
12. Wood WH, Fields BE. Hippotherapy: a systematic mapping review of peer-reviewed research, 1980 to 2018. *Disabil Rehabil*. 2019 Sep;1-25:1-25. PMID 31491353
13. Kim SG, Lee CW. The effects of hippotherapy on elderly persons' static balance and gait. *J Phys Ther Sci*. Jan 2014;26(1):25-27. PMID 24567669
14. Araujo TB, Silva NA, Costa JN, et al. Effect of equine-assisted therapy on the postural balance of the elderly. *Rev Bras Fisioter*. Sep-Oct 2011;15(5):414-419. PMID 22002189
15. de Araujo TB, de Oliveira RJ, Martins WR, et al. Effects of hippotherapy on mobility, strength and balance in elderly. *Arch Gerontol Geriatr*. May-Jun 2013;56(3):478-481. PMID 23290005

16. Homnick DN, Henning KM, Swain CV, et al. Effect of therapeutic horseback riding on balance in community- dwelling older adults with balance deficits. *J Altern Complement Med.* Jul 2013;19(7):622-626. PMID 23360659
17. Silkwood-Sherer DJ, Killian CB, Long TM, et al. Hippotherapy--an intervention to habilitate balance deficits in children with movement disorders: a clinical trial. *Phys Ther.* May 2012;92(5):707-717. PMID 22247403
18. Giagazoglou P, Arabatzi F, Dipla K, et al. Effect of a hippotherapy intervention program on static balance and strength in adolescents with intellectual disabilities. *Res Dev Disabil.* Nov 2012;33(6):2265-2270. PMID 22853887
19. Lechner HE, Kakebeeke TH, Hegemann D, et al. The effect of hippotherapy on spasticity and on mental well- being of persons with spinal cord injury. *Arch Phys Med Rehabil.* Oct 2007;88(10):1241-1248. PMID 17908564
20. American Hippotherapy Association, Inc. Statements of best practice for the use of hippotherapy by occupational therapy, physical therapy, and speech-language pathology professionals. Revised March 9, 2017. <https://americanhippotherapyassociation.org/wp-content/uploads/2018/09/Final-2017-Best-Practice-1.pdf>. Accessed January 20, 2020.
21. Bunketorp Kall L, Lundgren-Nilsson A, Blomstrand C, et al. The effects of a rhythm and music-based therapy program and therapeutic riding in late recovery phase following stroke: a study protocol for a three-armed randomized controlled trial. *BMC Neurol.* Nov 21 2012;12:141. PMID 23171380