



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an independent licensee of the Blue Cross and Blue Shield Association

Medical Policy Laser Treatment of Onychomycosis

Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

Policy Number: 562

BCBSA Reference Number: 2.01.89

NCD/LCD: NA

Related Policies

- Nonpharmacologic Treatment of Rosacea, #[462](#)
- Plastic Surgery, #[068](#)

Policy

**Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity
Medicare HMO BlueSM and Medicare PPO BlueSM Members**

Laser treatment of onychomycosis is [INVESTIGATIONAL](#).

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	This is not a covered service.
Commercial PPO and Indemnity	This is not a covered service.
Medicare HMO BlueSM	This is not a covered service.
Medicare PPO BlueSM	This is not a covered service.

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

CPT Codes

There is no specific CPT code for this service.

Description

Onychomycosis

Onychomycosis is a common chronic fungal infection of the nail. It is estimated to cause up to 50% of all nail disease and 33% of cutaneous fungal infections.¹ The condition can affect toenails or fingernails but is more frequently found in toenails. Primary infectious agents include dermatophytes (eg, *Trichophyton* species), yeasts (eg, *Candida albicans*), and nondermatophytic molds. In temperate Western countries, infections are generally caused by dermatophytes.

Aging is the most common risk factor for onychomycosis, most likely due to decreased blood circulation, longer exposure to fungi, and slower nail growth. Also, various medical conditions increase the risk of comorbid onychomycosis. They include diabetes, obesity, peripheral vascular disease, immunosuppression, and HIV infection. In certain populations, onychomycosis may lead to additional health problems. Although there is limited evidence of a causal link between onychomycosis and diabetic foot ulcers, at least one prospective study with diabetic patients found onychomycosis to be an independent predictor of foot ulcer.² Moreover, onychomycosis, especially more severe cases, may adversely impact the quality of life. Patients with onychomycosis have reported pain, uncomfortable nail pressure, embarrassment, and discomfort wearing shoes.^{3,4}

Diagnosis

The diagnosis of onychomycosis can be confirmed by potassium hydroxide preparation, culture, or histology.

Treatment

Treatments for onychomycosis include topical antifungals such as nail paints containing ciclopirox (ciclopiroxolamine) or amorolfine and oral antifungals such as terbinafine and itraconazole. These have low-to-moderate efficacy and a high relapse rate. Topical antifungals and some long-available oral medications (eg, griseofulvin) require a long course of treatment, which presents issues for patient compliance. Moreover, oral antifungal medications have been associated with adverse effects such as a risk of hepatotoxicity.

Several types of device-based therapies are under investigation for the treatment of onychomycosis, including ultrasound, iontophoresis, photodynamic therapy, and laser systems. A potential advantage of lasers is that they have greater tissue penetration than antifungal medication and thus may be more effective at treating infection embedded within the nail. Another potential advantage is that laser treatments are provided in a clinical setting in only one or several sessions and, thus, requires less long-term patient compliance.

Laser treatment of onychomycosis uses the principle of selective photothermolysis. This is defined as the precise targeting of tissue using a specific wavelength of light. The premise is that light is absorbed into the target area and heat generated by that energy is sufficient to damage the target area while sparing the surrounding area. The aim of laser treatment for onychomycosis is to heat the nail bed to temperatures required to disrupt fungal growth (approximately 40°-60°C) and at the same time avoid pain and necrosis to surrounding tissues.⁵

Characteristics of laser systems used to treat onychomycosis are listed in Table 1.⁵

Table 1. Characteristics of Lasers for Treating Onychomycosis

Variables	Characteristics
Wavelength	Lasers are single-wavelength light sources. There needs to be sufficient tissue penetration to adequately treat nail fungus. The near-infrared spectrum tends to be used because this part of the spectrum has maximum tissue penetrance in the dermis and epidermis and the nail plate is similar to the epidermis. To date, most laser systems for treating onychomycosis have been Neodymium yttrium aluminum garnet (Nd:YAG) lasers that typically operate at 1064 nm; 940- to 1320-nm and 1440-nm wavelengths are also options.
Pulse Duration	Pulses need to be short to avoid damaging the tissue surrounding the target area. For example, short-pulse systems have microsecond pulse durations and Q-switched lasers have nanosecond pulse durations.
Repetition rate (frequency of pulses, in hertz)	Spot size to the diameter of the laser beam. For treating onychomycosis, laser spot sizes range from 1 to 10 mm.
Fluence (in J/cm ²)	Fluence refers to the amount of energy delivered into the area.

Summary

Onychomycosis is a common fungal infection of the nail. Currently available treatments for onychomycosis, including systemic and topical antifungal medications, have relatively low efficacy and require a long course of treatment. Laser systems are proposed as another treatment option.

For individuals who have onychomycosis who receive treatment with laser therapy, the evidence includes small, randomized controlled trials. The relevant outcomes are symptoms, change in disease status, medication use, and treatment-related morbidity. The randomized controlled trials reported inconsistent results and had methodologic limitations. Clinical and mycologic outcomes differed across the trials, lacked consistent blinding of outcome assessments, and often reported outcomes on a per-nail basis without accounting for correlated measurements. The published evidence to date does not permit determining whether laser treatment improves health outcomes in patients with onychomycosis. Additional well-designed, adequately powered, and well-conducted randomized controlled trials are needed. The evidence is insufficient to determine the effects of the technology on health outcomes.

Policy History

Date	Action
2/2019	BCBSA National medical policy review. Description, summary and references updated. Policy statements unchanged.
1/2017	New references added from BCBSA National medical policy.
7/2015	New references added from BCBSA National medical policy.
9/2014	New references added from BCBSA National medical policy.
11/2013	New medical policy describing investigational indications. Effective 11/1/2013.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

References

1. [Rodgers P, Bassler M. Treating onychomycosis. Am Fam Physician. Feb 15 2001;63\(4\):663-672, 677-668. PMID 11237081](#)
2. [Boyko EJ, Ahroni JH, Cohen V, et al. Prediction of diabetic foot ulcer occurrence using commonly available clinical information: the Seattle Diabetic Foot Study. Diabetes Care. Jun 2006;29\(6\):1202-1207. PMID 16731996](#)
3. [Drake LA, Scher RK, Smith EB, et al. Effect of onychomycosis on quality of life. J Am Acad Dermatol. May 1998;38\(5 Pt 1\):702-704. PMID 9591814](#)
4. [Elewski BE. Onychomycosis. Treatment, quality of life, and economic issues. Am J Clin Dermatol. Jan-Feb 2000;1\(1\):19-26. PMID 11702301](#)
5. [Gupta A, Simpson F. Device-based therapies for onychomycosis treatment. Skin Therapy Lett. Oct 2012;17\(9\):4-9. PMID 23032936](#)
6. [Bristow IR. The effectiveness of lasers in the treatment of onychomycosis: a systematic review. J Foot Ankle Res. Aug 2014;7:34. PMID 25104974](#)
7. [Karsai S, Jager M, Oesterhelt A, et al. Treating onychomycosis with the short-pulsed 1064-nm-Nd:YAG laser: results of a prospective randomized controlled trial. J Eur Acad Dermatol Venereol. Jan 2017;31\(1\):175-180. PMID 27521028](#)
8. [Kim TI, Shin MK, Jeong KH, et al. A randomised comparative study of 1064 nm Neodymium-doped yttrium aluminium garnet \(Nd:YAG\) laser and topical antifungal treatment of onychomycosis. Mycoses. Jul 12 2016;59\(12\):803-810. PMID 27402466](#)
9. [El-Tatawy RA, Abd El-Naby NM, El-Hawary EE, et al. A comparative clinical and mycological study of Nd-YAG laser versus topical terbinafine in the treatment of onychomycosis. J Dermatolog Treat. Feb 11 2015:1-4. PMID 25669435](#)
10. [Xu Y, Miao X, Zhou B, et al. Combined oral terbinafine and long-pulsed 1,064-nm Nd: YAG laser treatment is more effective for onychomycosis than either treatment alone. Dermatol Surg. Nov 2014;40\(11\):1201-1207. PMID 25322165](#)
11. [Landsman AS, Robbins AH, Angelini PF, et al. Treatment of mild, moderate, and severe onychomycosis using 870- and 930-nm light exposure. J Am Podiatr Med Assoc. May-Jun 2010;100\(3\):166-177. PMID 20479446](#)
12. [Landsman AS, Robbins AH. Treatment of mild, moderate, and severe onychomycosis using 870- and 930-nm light exposure: some follow-up observations at 270 days. J Am Podiatr Med Assoc. Mar-Apr 2012;102\(2\):169-171. PMID 22461276](#)
13. [Ameen M, Lear JT, Madan V, et al. British Association of Dermatologists' guidelines for the management of onychomycosis 2014. Br J Dermatol. Nov 2014;171\(5\):937-958. PMID 25409999](#)