



## MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

### Medical Policy

## Whole Body Dual X-Ray Absorptiometry to Determine Body Composition

### Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

### Policy Number: 577

BCBSA Reference Number: 6.01.40

NCD/LCD: NA

### Related Policies

Vertebral Fracture Assessment with Densitometry, #[449](#)

### Policy

#### Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO Blue<sup>SM</sup> and Medicare PPO Blue<sup>SM</sup> Members

Dual x-ray absorptiometry (DXA) body composition studies are considered [INVESTIGATIONAL](#).

### Prior Authorization Information

#### Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

#### Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

|                                       | Outpatient                            |
|---------------------------------------|---------------------------------------|
| Commercial Managed Care (HMO and POS) | This is <b>not</b> a covered service. |
| Commercial PPO and Indemnity          | This is <b>not</b> a covered service. |
| Medicare HMO Blue <sup>SM</sup>       | This is <b>not</b> a covered service. |
| Medicare PPO Blue <sup>SM</sup>       | This is <b>not</b> a covered service. |

### CPT Codes / HCPCS Codes / ICD Codes

*Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.*

*Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.*

## **CPT Codes**

There is no specific CPT code for this service.

## **ICD Diagnosis Codes**

Investigational for all diagnoses.

## **Description**

### **BODY COMPOSITION MEASUREMENT**

Measurements of body composition have been used to study how lean body mass and body fat change during health and disease and have provided a research tool to study the metabolic effects of aging, obesity, and various wasting conditions such as occur with AIDS or after bariatric surgery. A variety of techniques has been researched, including most commonly, anthropomorphic measures, bioelectrical impedance, and dual x-ray absorptiometry (DXA). All of these techniques are based in part on assumptions about the distribution of different body compartments and their density, and all rely on formulas to convert the measured parameter into an estimate of body composition. Therefore, all techniques will introduce variation based on how the underlying assumptions and formulas apply to different populations of subjects (ie, different age groups, ethnicities, or underlying conditions). Techniques using anthropomorphics, bioelectrical impedance, underwater weighing, and DXA are briefly reviewed below.

### **Anthropomorphic Techniques**

Anthropomorphic techniques for the estimation of body composition include measurements of skinfold thickness at various sites, bone dimensions, and limb circumference. These measurements are used in various equations to predict body density and body fat. Due to its ease of use, measurement of skinfold thickness is one of the most common techniques. The technique is based on the assumption that the subcutaneous adipose layer reflects total body fat, but this association may vary with age and sex.

### **Bioelectrical Impedance**

Bioelectrical impedance is based on the relation among the volume of the conductor (ie, human body), the conductor's length (ie, height), the components of the conductor (ie, fat and fat-free mass), and its impedance. Estimates of body composition are based on the assumption that the overall conductivity of the human body is closely related to lean tissue. The impedance value is then combined with anthropomorphic data to give body compartment measures. The technique involves attaching surface electrodes to various locations on the arm and foot. Alternatively, the patient can stand on pad electrodes.

### **Underwater Weighing**

Underwater weighing requires the use of a specially constructed tank in which the subject is seated on a suspended chair. The subject is then submerged in the water while exhaling. While valued as a research tool, weighing people underwater is obviously not suitable for routine clinical use. This technique is based, on the assumption the body can be divided into 2 compartments with constant densities: adipose tissue, with a density of 0.9 g/cm<sup>3</sup>, and lean body mass (ie, muscle and bone), with a density of 1.1 g/cm<sup>3</sup>. One limitation of the underlying assumption is the variability in density between muscle and bone; eg, bone has a higher density than muscle, and bone mineral density varies with age and other conditions. Also, the density of body fat may vary, depending on the relative components of its constituents (eg, glycerides, sterols, glycolipids).

### **Dual X-Ray Absorptiometry**

While the cited techniques assume 2 body compartments, DXA can estimate 3 body compartments consisting of fat mass, lean body mass, and bone mass. DXA systems use a source that generates x-rays at 2 energies. The differential attenuation of the 2 energies is used to estimate the bone mineral content and the soft tissue composition. When 2 x-ray energies are used, only 2 tissue compartments can be measured; therefore, soft tissue measurements (ie, fat and lean body mass) can only be measured in

areas in which no bone is present. DXA can also determine body composition in defined regions (ie, the arms, legs, and trunk). DXA measurements are based in part on the assumption that the hydration of fat-free mass remains constant at 73%. Hydration, however, can vary from 67% to 85% and can vary by disease state. Other assumptions used to derive body composition estimates are considered proprietary by DXA manufacturers.

## Summary

Using low-dose x-rays of 2 different energy levels, whole body dual x-ray absorptiometry (DXA) measures lean tissue mass, total and regional body fat, as well as bone density. DXA scans have become a tool for research on body composition (eg, as a more convenient replacement for underwater weighing). This evidence review addresses potential applications in clinical care rather than research use of the technology.

For individuals who have a clinical condition associated with abnormal body composition who receive DXA body composition studies, the evidence includes several cross-sectional studies comparing DXA with other techniques. Relevant outcomes are symptoms and change in disease status. The available studies were primarily conducted in research settings and often used DXA body composition studies as a reference standard; these studies do not permit conclusions about the accuracy of DXA for measuring body composition. More importantly, no studies were identified in which DXA body composition measurements were actively used in patient management. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have a clinical condition managed by monitoring changes in body composition over time who receive serial DXA body composition studies, the evidence includes several prospective studies monitoring patients over time. Relevant outcomes are symptoms and change in disease status. The studies used DXA as a tool to measure body composition and were not designed to assess the accuracy of DXA. None of the studies used DXA findings to make patient management decisions or addressed how serial body composition assessment might improve health outcomes. The evidence is insufficient to determine the effects of the technology on health outcomes.

## Policy History

| Date           | Action   |
|----------------|--|
| 10/2018        | BCBSA National medical policy review. Description, summary and references updated. Policy statement unchanged. |
| 1/2016         | New references added from BCBSA National medical policy.   |
| 2/2016         | BCBSA National medical policy review. Abbreviation in policy statement changed to DXA.                         |
| 2/2015         | New references added from BCBSA National medical policy.   |
| 3/2014         | New references added from BCBSA National medical policy.   |
| 11/2011-4/2012 | Medical policy ICD 10 remediation: Formatting, editing and coding updates. No changes to policy statements.    |
| 12/2011        | New policy describing ongoing non-coverage. Effective 12/2011.   |

## Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

## References

1. Alves FD, Souza GC, Biolo A, et al. Comparison of two bioelectrical impedance devices and dual-energy X-ray absorptiometry to evaluate body composition in heart failure. *J Hum Nutr Diet*. Dec 2014;27(6):632-638. PMID 24684316
2. Ziai S, Coriati A, Chabot K, et al. Agreement of bioelectric impedance analysis and dual-energy X-ray absorptiometry for body composition evaluation in adults with cystic fibrosis. *J Cyst Fibros*. Sep 2014;13(5):585-588. PMID 24522087
3. Elkan AC, Engvall IL, Tengstrand B, et al. Malnutrition in women with rheumatoid arthritis is not revealed by clinical anthropometrical measurements or nutritional evaluation tools. *Eur J Clin Nutr*. Oct 2008;62(10):1239-1247. PMID 17637600
4. Jensky-Squires NE, Dieli-Conwright CM, Rossuello A, et al. Validity and reliability of body composition analysers in children and adults. *Br J Nutr*. Oct 2008;100(4):859-865. PMID 18346304
5. Kullberg J, Brandberg J, Angelhed JE, et al. Whole-body adipose tissue analysis: comparison of MRI, CT and dual energy X-ray absorptiometry. *Br J Radiol*. Feb 2009;82(974):123-130. PMID 19168691
6. Liem ET, De Lucia Rolfe E, L'Abee C, et al. Measuring abdominal adiposity in 6 to 7-year-old children. *Eur J Clin Nutr*. Jul 2009;63(7):835-841. PMID 19127281
7. Bedogni G, Agosti F, De Col A, et al. Comparison of dual-energy X-ray absorptiometry, air displacement plethysmography and bioelectrical impedance analysis for the assessment of body composition in morbidly obese women. *Eur J Clin Nutr*. Nov 2013;67(11):1129-1132. PMID 24022260
8. Monteiro PA, Antunes Bde M, Silveira LS, et al. Body composition variables as predictors of NAFLD by ultrasound in obese children and adolescents. *BMC Pediatr*. Jan 29 2014;14:25. PMID 24476029
9. Tompuri TT, Lakka TA, Hakulinen M, et al. Assessment of body composition by dual-energy X-ray absorptiometry, bioimpedance analysis and anthropometrics in children: the Physical Activity and Nutrition in Children study. *Clin Physiol Funct Imaging*. Jan 2015;35(1):21-33. PMID 24325400
10. Bazzocchi A, Ponti F, Cariani S, et al. Visceral fat and body composition changes in a female population after RYGBP: a two-year follow-up by DXA. *Obes Surg*. Mar 2015;25(3):443-451. PMID 25218013
11. Franzoni E, Ciccarese F, Di Pietro E, et al. Follow-up of bone mineral density and body composition in adolescents with restrictive anorexia nervosa: role of dual-energy X-ray absorptiometry. *Eur J Clin Nutr*. Feb 2014;68(2):247-252. PMID 24346474
12. International Society for Clinical Densitometry. 2015 ISCD Official Positions – Adult. 2015; <https://www.iscd.org/official-positions/2015-iscd-official-positions-adult/>. Accessed August 22, 2018.