



# MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an independent licensee of the Blue Cross and Blue Shield Association

## Medical Policy Surgery for Groin Pain in Athletes

### Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

### Policy Number: 695

BCBSA Reference Number: 7.01.142

NCD/LCD: N/A

### Related Policies

- Orthopedic Applications of Platelet-Rich Plasma, #[737](#)
- Surgical Treatment of Femoroacetabular Impingement, #[145](#)

### Policy

#### Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO Blue<sup>SM</sup> and Medicare PPO Blue<sup>SM</sup> Members

Surgical treatment of groin pain (also known as athletic pubalgia, Gilmore groin, osteitis pubis, pubic inguinal pain syndrome, inguinal disruption, slap shot gut, sportsmen groin, footballers groin injury complex, hockey groin syndrome, athletic hernia, sports hernia or core muscle injury) is considered [INVESTIGATIONAL](#).

### Prior Authorization Information

#### Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

#### Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	This is <b>not</b> a covered service.
Commercial PPO and Indemnity	This is <b>not</b> a covered service.
Medicare HMO Blue <sup>SM</sup>	This is <b>not</b> a covered service.
Medicare PPO Blue <sup>SM</sup>	This is <b>not</b> a covered service.

## CPT Codes / HCPCS Codes / ICD Codes

*Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.*

*Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.*

## CPT Codes

No specific CPT codes.

## Description

### Diagnosis

A diagnosis of groin pain in athletes is based primarily on history, physical exam, and imaging. The clinical presentation will generally be a gradual onset of progressive groin pain associated with the activity. A physical exam will not reveal any evidence for a standard inguinal hernia or groin muscle strain. Imaging with magnetic resonance imaging (MRI) or ultrasound is generally done as part of the workup. In addition to the exclusion of other sources of lower abdominal and groin pain (eg, stress fractures, femoroacetabular impingement, labral tears), imaging may identify injury to the soft tissues of the groin and abdominal wall.<sup>1</sup>

### Treatment

#### Conservative

Many injuries will heal with conservative treatment, which includes rest, icing, nonsteroidal anti-inflammatory drugs, and rehabilitation exercises. A physical therapy (PT) program that focuses on strength and coordination of core muscles acting on the pelvis may improve recovery. In a 1999 study, 68 athletes with chronic adductor-related groin pain were randomized to 8 to 12 weeks of an active training PT program that focused on strength and coordination of core muscles, particularly adductors, or to standard PT without active training.<sup>2</sup> At 4 months posttreatment, 68% of patients in the active training group had returned to sports without groin pain compared with 12% in the standard PT group. At 8- to 12-year follow-up, 50% of athletes in the active training group rated their outcomes as excellent compared with 22% in the standard PT group.<sup>3</sup> For in-season professional athletes, injections of corticosteroid or platelet-rich plasma (see policy #[737](#)), or a short corticosteroid burst with taper have also been used.

#### Surgical

Surgical treatment is typically reserved for patients who have failed at least 3 months of conservative treatment. One approach consists of open or laparoscopic sutured hernia repair with mesh reinforcement of the posterior wall of the inguinal canal. Laparoscopic procedures may use either a transabdominal preperitoneal or an extraperitoneal approach. A variety of musculotendinous defects, nerve entrapments, and inflammatory conditions have been observed with surgical exploration. Meyers et al. (2008) have proposed that any of the 17 soft tissues that attach or cross the pubic symphysis can be involved, leading to as many as 26 surgical procedures and 121 different combinations of procedures that address the various core muscle injuries.<sup>4</sup> The objective is to stabilize the pubic joint by tightening or broadening the attachments of various structures to the pubic symphysis and/or by loosening the attachments or other supporting structures via epimysiotomy or detachment.

Because various surgical procedures used to treat sports-related groin pain have reported success, it has been proposed that general fibrosis from any surgery may act to stabilize the anterior pelvis and thus play a role in improved surgical outcomes.

## Summary

Sports-related groin pain, commonly known as athletic pubalgia or sports hernia, is characterized by disabling activity-dependent lower abdominal and groin pain not attributable to any other cause.

Athletic pubalgias most frequently diagnosed in high-performance male athletes, particularly those who participate in sports that involve rapid twisting and turning such as soccer, hockey, and football. For patients who fail conservative therapy, surgical repair of any defects identified in the muscles, tendons, or nerves has been proposed.

For individuals who have sports-related groin pain who receive mesh reinforcement or who receive surgical repair and release of soft tissue, the evidence includes 2 randomized controlled trials (RCT's) and a number of case series. Relevant outcomes are symptoms, functional outcomes, and treatment-related morbidity. The evidence on mesh reinforcement for inguinal-related groin pain includes 2 RCTs and a large prospective series. Results of the RCTs have suggested that, in carefully selected patients, mesh reinforcement results in an earlier return to play. However, a large prospective series from 2016 indicated that only about 20% of patients with chronic groin pain benefit from inguinal surgery. Further study is needed to define the patient population that would benefit from this treatment approach. An alternative approach to the treatment of groin pain in athletes involves repair or release of soft tissue. This approach has been reported in a large series. It included a 2008 review of medical records spanning 2 decades and over 5000 cases. More recent reports on these procedures from other institutions are needed. The evidence is insufficient to determine the effects of the technology on health outcomes.

## Policy History

Date	Action
4/2020	BCBSA National medical policy review. Description, summary and references updated. Policy statements unchanged.
4/2019	BCBSA National medical policy review. Description, summary and references updated. Policy statements unchanged.
3/2018	New references added from BCBSA National medical policy. Background and summary clarified.
4/2017	BCBSA National medical policy review. Athletic pubalgia" changed to "groin pain in athletes". Title changed to "Surgery for Groin Pain in Athletes." 4/1/2017.
12/2014	New policy describing investigational indications. Effective 12/1/2014.

## Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

## References

1. Khan W, Zoga AC, Meyers WC. Magnetic resonance imaging of athletic pubalgia and the sports hernia: current understanding and practice. *Magn Reson Imaging Clin N Am*. Feb 2013;21(1):97-110. PMID 23168185
2. Holmich P, Uhrskou P, Ulnits L, et al. Effectiveness of active physical training as treatment for long-standing adductor-related groin pain in athletes: randomised trial. *Lancet*. Feb 6 1999;353(9151):439-443. PMID 9989713
3. Holmich P, Nyvold P, Larsen K. Continued significant effect of physical training as treatment for overuse injury: 8- to 12-year outcome of a randomized clinical trial. *Am J Sports Med*. Nov 2011;39(11):2447-2451. PMID 21813441
4. Meyers WC, McKechnie A, Philippon MJ, et al. Experience with sports hernia spanning two decades. *Ann Surg*. Oct 2008;248(4):656-665. PMID 18936579
5. Litwin DE, Sneider EB, McEnaney PM, et al. Athletic pubalgia (sports hernia). *Clin Sports Med*. Apr 2011;30(2):417-434. PMID 21419964
6. Weir A, Brukner P, Delahunt E, et al. Doha agreement meeting on terminology and definitions in groin pain in athletes. *Br J Sports Med*. Jun 2015;49(12):768-774. PMID 26031643

7. Munegato D, Bigoni M, Gridavilla G, et al. Sports hernia and femoroacetabular impingement in athletes: A systematic review. *World J Clin Cases*. Sep 16 2015;3(9):823-830. PMID 26380829
8. Thorborg K, Holmich P, Christensen R, et al. The Copenhagen Hip and Groin Outcome Score (HAGOS): development and validation according to the COSMIN checklist. *Br J Sports Med*. May 2011;45(6):478-491. PMID 21478502
9. Paajanen H, Brinck T, Hermunen H, et al. Laparoscopic surgery for chronic groin pain in athletes is more effective than nonoperative treatment: a randomized clinical trial with magnetic resonance imaging of 60 patients with sportsman's hernia (athletic pubalgia). *Surgery*. Jul 2011;150(1):99-107. PMID 21549403
10. Ekstrand J, Ringborg S. Surgery versus conservative treatment in soccer players with chronic groin pain: A prospective randomised study in soccer players. *Eur J Sports Traumatol Rel Res*. 2001;23:141-145.
11. Ahumada LA, Ashruf S, Espinosa-de-los-Monteros A, et al. Athletic pubalgia: definition and surgical treatment. *Ann Plast Surg*. Oct 2005;55(4):393-396. PMID 16186706
12. Steele P, Annear P, Grove JR. Surgery for posterior inguinal wall deficiency in athletes. *J Sci Med Sport*. Dec 2004;7(4):415-421; discussion 422-413. PMID 15712496
13. Paajanen H, Syvahuoko I, Airo I. Totally extraperitoneal endoscopic (TEP) treatment of sportsman's hernia. *Surg Laparosc Endosc Percutan Tech*. Aug 2004;14(4):215-218. PMID 15472551
14. Kumar A, Doran J, Batt ME, et al. Results of inguinal canal repair in athletes with sports hernia. *J R Coll Surg Edinb*. Jun 2002;47(3):561-565. PMID 12109611
15. Irshad K, Feldman LS, Lavoie C, et al. Operative management of hockey groin syndrome: 12 years of experience in National Hockey League players. *Surgery*. Oct 2001;130(4):759-764; discussion 764-756. PMID 11602909
16. Kopelman D, Kaplan U, Hatoum OA, et al. The management of sportsman's groin hernia in professional and amateur soccer players: a revised concept. *Hernia*. Feb 2016;20(1):69-75. PMID 25380561
17. American Academy of Orthopaedic Surgeons, Wilkerson R. OrthoInfo: Sports Hernia (Athletic Pubalgia). 2017; <http://orthoinfo.aaos.org/topic.cfm?topic=A00573>. Accessed December 28, 2019.