



## MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an independent Licensee of the Blue Cross and Blue Shield Association

### Medical Policy

## Multitarget Polymerase Chain Reaction Testing for Diagnosis of Bacterial Vaginosis

### Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

### Policy Number: 711

BCBSA Reference Number: 2.04.127

NCD/LCD: N/A

### Related Policies

- Identification of Microorganisms Using Nucleic Acid Probe, [#555](#)

### Policy

#### Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO Blue<sup>SM</sup> and Medicare PPO Blue<sup>SM</sup> Members

Multitarget polymerase chain reaction (PCR) testing for diagnosis of bacterial vaginosis is considered **INVESTIGATIONAL**.

### Prior Authorization Information

#### Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

#### Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	This is <b>not</b> a covered service.
Commercial PPO and Indemnity	This is <b>not</b> a covered service.
Medicare HMO Blue <sup>SM</sup>	This is <b>not</b> a covered service.
Medicare PPO Blue <sup>SM</sup>	This is <b>not</b> a covered service.

## CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

### CPT codes

There is no specific CPT code for this testing.

### Description

#### Bacterial Vaginosis

BV is a condition caused by an imbalance in the normal bacteria vaginal flora. It is common, especially in women of reproductive age. While there is no single known etiologic agent, there is a shift in vaginal flora that involves a depletion of hydrogen peroxide-producing *Lactobacillus* species with a rise in vaginal pH and overgrowth of other bacteria, including *Gardnerella vaginalis*, *Mycoplasma hominis*, *Peptostreptococcus*, *Mobiluncus* species, and other anaerobic gram-negative rods.

Vaginal culture is not an appropriate diagnostic method to identify BV because BV is not caused by the presence of a particular bacterial species.

Various commercial tests provide rapid and accurate pH evaluation and amine detection. For example, automated devices that measure the volatile gases produced from vaginal samples and a colorimetric pH test are commercially available.

Nucleic acid probes of DNA fragments are available to detect and quantify specific bacteria in vaginal fluid samples. Polymerase chain reaction (PCR) methods extract and amplify the DNA fragments using either universal or specific primers. The result can be qualitative (to assess whether a specific microorganism is present) or quantitative (to assess how many microorganisms are present). The technology can be used to measure multiple organisms (eg, those known to be associated with BV) at the same time and is commercially available as multitarget PCR testing.

(Policy #555 addresses the use of nucleic acid probes to detect other microorganisms of clinical significance. This policy includes identification of *G. vaginalis* which is a single microorganism associated with BV.)

#### Proposed Multitarget PCR Test

Examples of commercially available multitarget PCR tests and the organisms in the panels are shown in Table 1; this may not be an exhaustive list of all commercially available tests.

Table 1. Components of Commercially Available Multitarget PCR Tests

Organism	SureSwab	BD Max	MDL Panel	NuSwab	GenPath BV Test
<i>Atopobium vaginae</i>	X	X	X	X	X
<i>Gardnerella vaginalis</i>	X	X	X		X
<i>Lactobacillus</i> species	X <sup>a</sup>	X	X		X
<i>Megasphaera</i> (type 1, type 2, and/or species)	X	X	X	X	X

BVAB (type 1 and/or type 2)		X	X	X	X
-----------------------------	--	---	---	---	---

BVAB: bacterial vaginosis-associated bacteria; MDL: Medical Diagnostics Laboratory; PCR: polymerase chain reaction.

<sup>a</sup> *Lactobacillus crispatus* and *Lactobacillus jensenii*.

The SureSwab Total (Quest Diagnostics) test involves obtaining vaginal swab specimens, extracting total DNA, and quantitating the four types of bacteria using PCR. Results are reported as log cells per milliliter for each organism and concentrations of all *Lactobacilli* species are reported together then classified into one of the following three categories: not supportive, equivocal, and supportive.

A classification of *not supportive* of BV diagnosis is based on:

- The presence of *Lactobacillus* species, *G. vaginalis* levels <6.0 log cells/mL, and absence of *Atopobium vaginae* and *Megasphaera* species; or
- The absence of *Lactobacillus* species, *G. vaginalis* levels <6.0 log cells/mL, and absence of *A. vaginae* and *Megasphaera* species; or
- The absence of all targeted organisms.
- A classification of equivocal is based on:
- The presence of *Lactobacillus* species, plus *G. vaginalis* at least 6.0 log cells/mL, and/or presence of *A. vaginae* and/or *Megasphaera* species.

A classification of supportive of BV diagnosis is based on the absence of *Lactobacillus* species, and presence of *G. vaginalis* levels of at least 6.0 log cells/mL, and presence of *A. vaginae* and/or *Megasphaera* species.

Another product, the BD Max (Becton, Dickinson), tests for markers of BV and vaginitis. The test uses a similar process to that described for SureSwab. Vaginal swab specimens are collected, DNA is extracted, and real-time PCR is used to quantitate targeted organisms. Results of BV marker tests are not reported for individual organisms. Instead, qualitative BV results are reported as positive or negative for BV based on the relative quantity of the various organisms.

Medical Diagnostics Laboratory offers a Bacterial Vaginosis Panel. Markers are shown above in Table 1 and are assessed using real-time PCR and *Lactobacillus* is profiled using quantitative PCR. GenPath Diagnostics also offers a bacterial vaginosis test.

The NuSwab® Select BV test (Laboratory Corporation of American) uses semiquantitative PCR analysis of three predictive marker organisms of vaginal dysbiosis to generate a total score that is associated with the presence or absence of BV. In this test system, samples with a total score of 0 to 1 are considered negative for BV, samples with a score of 3 to 6 are positive for BV, and samples with a score of 2 are indeterminate for BV.

Several of the manufacturers of the BV tests also have extensions that include other causes of vaginitis such as *Trichomonas vaginalis* and *Candidiasis* species.

## Summary

Bacterial vaginosis (BV) is a common medical condition resulting from an imbalance in the normal vaginal flora. Although identification of *Gardnerella vaginalis* has traditionally been associated with BV, there is no single etiologic agent. Most cases are asymptomatic, and most symptomatic cases can be diagnosed

using clinical and microscopic evaluation. Multitarget polymerase chain reaction (PCR) testing is proposed as an alternative to currently available laboratory tests to diagnose BV. This test may improve outcomes if it is a more accurate and reliable method to diagnose BV.

In individuals who have signs or symptoms of BV who receive multitarget PCR testing, the evidence includes several prospective studies on technical performance and diagnostic accuracy. The relevant outcomes are test validity, symptoms, and change in disease status. Several studies have evaluated the diagnostic accuracy of multitarget PCR tests for BV, including two studies evaluating commercially available tests. The studies found sensitivities between 90% and 95% and specificities between 85% and 90% compared with standard methods of diagnosis. Most studies used a combination of the Amsel criteria and Nugent scoring as the reference standard. There is a lack of direct evidence on the clinical utility of PCR testing for BV (ie, studies showing that testing leads to better patient management decisions and/or better health outcomes than current approaches). Moreover, a chain of evidence does not currently support multitarget testing because most symptomatic women can be diagnosed with a standard workup. The evidence is insufficient to determine the effects of the technology on health outcomes.

## Policy History

Date	Action
2/2019	BCBSA National medical policy review. Description, summary and references updated. Policy statements unchanged.
8/2017	New references added from BCBSA National medical policy.
1/2016	New references added from BCBSA National medical policy.
3/2015	New medical policy describing investigational indications. Effective 3/1/2015.

## Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

## References

1. Lamont RF, Sobel JD, Akins RA, et al. The vaginal microbiome: new information about genital tract flora using molecular based techniques. *BJOG*. Apr 2011;118(5):533-549. PMID 21251190
2. Centers for Disease Control and Prevention (CDC). Bacterial Vaginosis (BV) Statistics. 2015; <http://www.cdc.gov/std/bv/stats.htm>. Accessed November 27, 2018.
3. Amsel R, Totten PA, Spiegel CA, et al. Nonspecific vaginitis. Diagnostic criteria and microbial and epidemiologic associations. *Am J Med*. Jan 1983;74(1):14-22. PMID 6600371
4. Landers DV, Wiesenfeld HC, Heine RP, et al. Predictive value of the clinical diagnosis of lower genital tract infection in women. *Am J Obstet Gynecol*. Apr 2004;190(4):1004-1010. PMID 15118630
5. Nugent RP, Krohn MA, Hillier SL. Reliability of diagnosing bacterial vaginosis is improved by a standardized method of gram stain interpretation. *J Clin Microbiol*. Feb 1991;29(2):297-301. PMID 1706728
6. Ison CA, Hay PE. Validation of a simplified grading of Gram stained vaginal smears for use in genitourinary medicine clinics. *Sex Transm Infect*. Dec 2002;78(6):413-415. PMID 12473800
7. Hilbert DW, Smith WL, Chadwick SG, et al. Development and validation of a highly accurate quantitative real-time PCR assay for diagnosis of bacterial vaginosis. *J Clin Microbiol*. Apr 2016;54(4):1017-1024. PMID 26818677
8. Food and Drug Administration. Evaluation of Automatic Class III Designation For BD Max Vaginal Panel: Decision Summary. 2016; [https://www.accessdata.fda.gov/cdrh\\_docs/reviews/DEN160001.pdf](https://www.accessdata.fda.gov/cdrh_docs/reviews/DEN160001.pdf). Accessed November 27, 2018.

9. Gaydos CA, Beqaj S, Schwebke JR, et al. Clinical Validation of a Test for the Diagnosis of Vaginitis. *Obstet Gynecol.* Jul 2017;130(1):181-189. PMID 28594779
10. Cartwright CP, Pherson AJ, Harris AB, et al. Multicenter study establishing the clinical validity of a nucleic-acid amplification-based assay for the diagnosis of bacterial vaginosis. *Diagn Microbiol Infect Dis.* Jun 1 2018. PMID 29937222
11. Cartwright CP, Lembke BD, Ramachandran K, et al. Development and validation of a semiquantitative, multitarget PCR assay for diagnosis of bacterial vaginosis. *J Clin Microbiol.* Jul 2012;50(7):2321-2329. PMID 22535982
12. Kusters JG, Reuland EA, Bouter S, et al. A multiplex real-time PCR assay for routine diagnosis of bacterial vaginosis. *Eur J Clin Microbiol Infect Dis.* Sep 2015;34(9):1779-1785. PMID 26143346
13. Rumyantseva T, Shipitsyna E, Guschin A, et al. Evaluation and subsequent optimizations of the quantitative AmpliSens Florocenosis/Bacterial vaginosis-FRT multiplex real-time PCR assay for diagnosis of bacterial vaginosis. *Apmis.* Dec 2016;124(12):1099-1108. PMID 27714844
14. Rumyantseva TA, Bellen G, Romanuk TN, et al. Utility of microscopic techniques and quantitative real-time polymerase chain reaction for the diagnosis of vaginal microflora alterations. *J Low Genit Tract Dis.* Jul 11 2014. PMID 25023332
15. van der Veer C, van Houdt R, van Dam A, et al. Accuracy of a commercial multiplex PCR for the diagnosis of bacterial vaginosis. *J Med Microbiol.* Sep 2018;67(9):1265-1270. PMID 29985123
16. Workowski KA, Bolan GA, Centers for Disease Control and Prevention, et al. Sexually transmitted diseases treatment guidelines, 2015. *MMWR Recomm Rep.* Jun 5 2015;64(RR-03):1-137. PMID 26042815
17. Committee on Practice Bulletins-Obstetrics, The American College of Obstetricians Gynecologists. Practice bulletin no. 130: prediction and prevention of preterm birth. *Obstet Gynecol.* Oct 2012;120(4):964-973. PMID 22996126
18. U.S. Preventive Services Task Force. Bacterial Vaginosis in Pregnancy to Prevent Preterm Delivery: Screening. 2008; <https://www.uspreventiveservicestaskforce.org/Page/Document/UpdateSummaryFinal/bacterial-vaginosis-in-pregnancy-to-prevent-preterm-delivery-screening>. Accessed November 27, 2018.