



## MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

# Medical Policy Dry Needling of Myofascial Trigger Points

### Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

### Policy Number: 792

BCBSA Reference Number: 2.01.100

NCD/LCD: N/A

### Related Policies

None

### Policy

#### Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO Blue<sup>SM</sup> and Medicare PPO Blue<sup>SM</sup> Members

Dry needling of trigger points for the treatment of myofascial pain is considered [INVESTIGATIONAL](#).

### Prior Authorization Information

#### Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

#### Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	This is <b>not</b> a covered service.
Commercial PPO and Indemnity	This is <b>not</b> a covered service.
Medicare HMO Blue <sup>SM</sup>	This is <b>not</b> a covered service.
Medicare PPO Blue <sup>SM</sup>	This is <b>not</b> a covered service.

### CPT Codes / HCPCS Codes / ICD Codes

*Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.*

*Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.*

*The following codes are included below for informational purposes only; this is not an all-inclusive list.*

## **CPT Codes**

There are not any CPT codes for this procedure

## **Description**

### **Dry Needling**

Dry needling refers to a procedure in which a fine needle is inserted into the skin and muscle at a site of myofascial pain. The needle may be moved in an up-and-down motion, rotated, and/or left in place for as long as 30 minutes. The intent is to stimulate underlying myofascial trigger points, muscles, and connective tissues to manage myofascial pain. Dry needling may be performed with acupuncture needles or standard hypodermic needles but is performed without the injection of medications (eg, anesthetics, corticosteroids). Dry needling is proposed to treat dysfunctions in skeletal muscle, fascia, and connective tissue; diminish persistent peripheral pain, and reduce impairments of body structure and function.

The physiologic basis for dry needling depends on the targeted tissue and treatment objectives. The most studied targets are trigger points. Trigger points are discrete, focal, hyperirritable spots within a taut band of skeletal muscle fibers that produce local and/or referred pain when stimulated. Trigger points are associated with local ischemia and hypoxia, a significantly lowered pH, local and referred pain and altered muscle activation patterns.<sup>1</sup> Trigger points can be visualized by magnetic resonance imaging and elastography. The reliability of manual identification of trigger points has not been established.

Deep dry needling is believed to inactivate trigger points by eliciting contraction and subsequent relaxation of the taut band via a spinal cord reflex. This local twitch response is defined as a transient visible or palpable contraction or dimpling of the muscle, and has been associated with alleviation of spontaneous electrical activity; reduction of numerous nociceptive, inflammatory, and immune system related chemicals; and relaxation of the taut band.<sup>1</sup> Deep dry needling of trigger points is believed to reduce local and referred pain, improve range of motion, and decrease trigger point irritability.

Superficial dry needling is thought to activate mechanoreceptors and have an indirect effect on pain by inhibiting C-fiber pain impulses. The physiologic basis for dry needling treatment of excessive muscle tension, scar tissue, fascia, and connective tissues is not as well described in the literature.<sup>1</sup>

## **Summary**

Trigger points are discrete, focal, hyperirritable spots within a taut band of skeletal muscle fibers that produce local and/or referred pain when stimulated. Dry needling refers to a procedure whereby a fine needle is inserted into the trigger point to induce a twitch response and relieve the pain.

For individuals who have myofascial trigger points associated with neck and/or shoulder pain who receive dry needling of trigger points, the evidence includes randomized controlled trials (RCTs) and a systematic review. The relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. As reported in the systematic review of literature published through 2013, only 1 of 8 studies found significantly greater reductions in pain with dry needling compared with other treatments. Two more recent RCTs comparing dry needling with manual therapy did not find significantly better outcomes after dry needling. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have myofascial trigger points associated with plantar heel pain who receive dry needling of trigger points, the evidence includes RCTs, quasi-experimental studies, and a systematic review. The relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. The systematic review, which included three quasi-experimental studies, rated study quality as

poor. One RCT was double-blind and sham-controlled; it found a statistically significant greater reduction in pain in the dry needling group than in the sham group but the difference was not clinically significant (ie, it did not meet the prespecified minimally important difference). The other RCT, a single-blind trial comparing dry needling with usual care, found a significantly greater reduction in pain at the end of active treatment but not at follow-up one month later. Moreover, range of motion outcomes did not differ significantly between groups at either time point. To date, the studies have not demonstrated a statistical or a clinical benefit for dry needling. Additional RCTs, especially those with a sham-control group, would strengthen the evidence base. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have myofascial trigger points associated with temporomandibular myofascial pain who receive dry needling of trigger points, the evidence includes an RCT. The relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. One double-blind, sham-controlled randomized trial was identified; it found that one week after completing the intervention, there were no statistically significant differences between groups in pain scores or function (unassisted jaw opening without pain). There was a significantly higher pain pressure threshold in the treatment group. Additional RCTs, especially those with a sham-control group, are needed. The evidence is insufficient to determine the effects of the technology on health outcomes.

## Policy History

Date	Action
5/2019	BCBSA National medical policy review. Description, summary and references updated. Policy statements unchanged.
5/2017	New references added from BCBSA National medical policy.
9/2016	Policy implementation delayed from 7/1/2016 to 9/1/2016. Coding information clarification. Effective 9/1/2016.
7/2016	New medical policy describing investigational indications. Effective 7/1/2016.

## Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

## References

1. American Physical Therapy Association (APTA). Educational resource paper: Description of Dry Needling in Clinical Practice. 2013; <http://www.apta.org/Statelssues/DryNeedling/ClinicalPracticeResourcePaper/>. Accessed March 13, 2019.
2. Alvarez DJ, Rockwell PG. Trigger points: diagnosis and management. *Am Fam Physician*. Feb 15 2002;65(4):653-660. PMID 11871683.
3. Cagnie B, Castelein B, Pollie F, et al. Evidence for the use of ischemic compression and dry needling in the management of trigger points of the upper trapezius in patients with neck pain: a systematic review. *Am J Phys Med Rehabil*. Jul 2015;94(7):573-583. PMID 25768071.
4. Llamas-Ramos R, Pecos-Martin D, Gallego-Izquierdo T, et al. Comparison of the short-term outcomes between trigger point dry needling and trigger point manual therapy for the management of chronic mechanical neck pain: a randomized clinical trial. *J Orthop Sports Phys Ther*. Nov 2014;44(11):852-861. PMID 25269764.
5. De Meulemeester KE, Castelein B, Coppieters I, et al. Comparing trigger point dry needling and manual pressure technique for the management of myofascial neck/shoulder pain: a randomized clinical trial. *J Manipulative Physiol Ther*. Jan 2017;40(1):11-20. PMID 28017188.

6. Perez-Palomares S, Oliván-Blázquez B, Perez-Palomares A, et al. Contribution of dry needling to individualized physical therapy treatment of shoulder pain: a randomized clinical trial. *J Orthop Sports Phys Ther.* Jan 2017;47(1):11-20. PMID 27937046.
7. Cerezo-Tellez E, Lacomba MT, Fuentes-Gallardo I, et al. Dry needling of the trapezius muscle in office workers with neck pain: a randomized clinical trial. *J Man Manip Ther.* Sep 2016;24(4):223-232. PMID 27582622.
8. Cerezo-Tellez E, Torres-Lacomba M, Fuentes-Gallardo I, et al. Effectiveness of dry needling for chronic nonspecific neck pain: a randomized, single-blinded, clinical trial. *Pain.* Sep 2016;157(9):1905-1917. PMID 27537209.
9. Cotchett MP, Landorf KB, Munteanu SE. Effectiveness of dry needling and injections of myofascial trigger points associated with plantar heel pain: a systematic review. *J Foot Ankle Res.* Sep 1 2010;3:18. PMID 20807448.
10. Cotchett MP, Munteanu SE, Landorf KB. Effectiveness of trigger point dry needling for plantar heel pain: a randomized controlled trial. *Phys Ther.* Aug 2014;94(8):1083-1094. PMID 24700136.
11. Eftekharsadat B, Babaei-Ghazani A, Zeinolabedinzadeh V. Dry needling in patients with chronic heel pain due to plantar fasciitis: A single-blinded randomized clinical trial. *Med J Islam Repub Iran.* Sep 2016;30:401. PMID 27683642.
12. Diracoglu D, Vural M, Karan A, et al. Effectiveness of dry needling for the treatment of temporomandibular myofascial pain: a double-blind, randomized, placebo controlled study. *J Back Musculoskelet Rehabil.* Dec 2012;25(4):285-290. PMID 23220812.
13. Brady S, McEvoy J, Dommerholt J, et al. Adverse events following trigger point dry needling: a prospective survey of chartered physiotherapists. *J Man Manip Ther.* Aug 2014;22(3):134-140. PMID 25125935.
14. American Physical Therapy Association (APTA). Physical Therapists and the Performance of Dry Needling. 2012; <http://www.apta.org/Statelssues/DryNeedling/ResourcePaper/>. Accessed March 13, 2019.
15. American Academy of Orthopaedic Physical Therapists. AAOMPT position statement on dry needling. 2009; [http://aaompt.org/Main/About\\_Us/Position\\_Statements/Main/About\\_Us/Position\\_Statements.aspx?hkey=03f5a333-3-f28d-4715-b355-cb25fa9bac2c](http://aaompt.org/Main/About_Us/Position_Statements/Main/About_Us/Position_Statements.aspx?hkey=03f5a333-3-f28d-4715-b355-cb25fa9bac2c). Accessed March 13, 2019.