



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an independent licensee of the Blue Cross and Blue Shield Association

Medical Policy Surgical Deactivation of Headache Trigger Sites

Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

Policy Number: 801

BCBSA Reference Number: 7.01.135

NCD/LCD: N/A

Related Policies

- Biofeedback for the Treatment of Headache, #[152](#)
- Botulinum Toxin Injections, #[006](#)
- Occipital Nerve Stimulation, #[237](#)

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO BlueSM and Medicare PPO BlueSM Members

Surgical deactivation of trigger sites is considered **INVESTIGATIONAL** for the treatment of migraine and non-migraine headache.

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	This is not a covered service.
Commercial PPO and Indemnity	This is not a covered service.
Medicare HMO Blue SM	This is not a covered service.
Medicare PPO Blue SM	This is not a covered service.

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

CPT Codes

There is no specific SPT code for this service.

Description

MIGRAINE HEADACHE

Migraine is a common headache disorder with a prevalence in the United States of approximately 18% in women and 6% in men.¹ According to the International Headache Society (2013), migraine headache is a recurrent disorder with attacks lasting 4 to 72 hours.² Typical features of migraine headaches include unilateral location, pulsating quality, moderate or severe intensity, and associated symptoms such as nausea, photophobia, and/or phonophobia.

Treatment

A variety of medications are used to treat acute migraine episodes. They include medications taken at the onset of an attack to abort the attack (triptans, ergotamines), and medications to treat the pain and other symptoms of migraines once they are established (nonsteroidal anti-inflammatory drugs, narcotic analgesics, antiemetics). Prophylactic medication therapy may be appropriate for people with migraines that occur more than two days per week. In addition to medication, behavioral treatments such as relaxation and cognitive therapy are used to manage migraine headache. Moreover, botulinum toxin type A injections are a U.S. Food and Drug Administration–approved treatment for chronic migraine (migraines occurring on at least 15 days a month for at least 3 months).

Surgical Deactivation

Surgical deactivation of trigger sites is another proposed treatment of migraine headache. The procedure was developed by a plastic surgeon (Bahman Guyuron, MD), following observations that some patients who had cosmetic forehead lifts reported improvement or elimination of migraine symptoms postsurgery.^{3,4} The procedure is based on the theory that migraine headaches arise due to inflammation of trigeminal nerve branches in the head and neck caused by irritation of the surrounding musculature, bony foramen, and perhaps fascia bands. Accordingly, surgical treatment of migraines involves removing the relevant nerve sections, muscles, fascia, and/or vessels. The treatment is also based on the theory there are specific migraine trigger sites and that these sites can be located in individual patients. In studies conducted by Guyuron's research group, clinical evaluation and diagnostic injections of botulinum toxin have been used to locate trigger sites. The specific surgical procedure varies according to the patient's migraine trigger site. The surgical procedures are performed under general anesthesia in an ambulatory care setting and take an average of one hour.

Surgical procedures have been developed at four trigger sites: frontal, temporal, rhinogenic, and occipital. Frontal headaches are believed to be activated by irritation of the supratrochlear and suborbital nerves by glabellar muscles or vessels. The surgical procedure involves the removal of the glabellar muscles encasing these nerves. Fat from the upper eyelid is used to fill the defect in the muscles and shield the nerve. Temporal headaches may be activated by inflammation of the zygomatico-temporal branch of the trigeminal nerve by the temporalis muscles or vessels adjacent to the nerve. To treat migraines located at this trigger site, a segment (≈ 2.5 cm) of the zygomatico-temporal branch of the trigeminal nerve is removed endoscopically. Rhinogenic headaches may involve intranasal abnormalities (eg, deviated septum), which may irritate the end branches of the trigeminal nerve. Surgical treatment includes septoplasty and turbinectomy. Finally, occipital headaches may be triggered by irritation of the occipital nerve caused by the semispinalis capitis muscle or the occipital artery. Surgery consists of removal of a segment of the semispinalis capitis muscle medial to the greater occipital nerve approximately 1 cm wide

and 2.5 cm long, followed by insertion of a subcutaneous flap between the nerve and the muscle to avoid nerve impingement.

Non-Migraine Headache

It has been proposed that other types of headaches (eg, tension headaches) may also be triggered by irritation of the trigeminal nerve.

Treatment

Although the mechanism of action is less well established for headaches other than migraine, it is possible that surgical treatment of trigger sites may also be beneficial for some non–migraine headaches.

Summary

Migraine is a common headache disorder that is treated using various medications, which can be taken at the onset of an attack and/or for migraine prophylaxis. Other treatments include behavioral treatments and botulinum toxin injections. Surgical deactivation of trigger sites is another proposed treatment. Surgical deactivation is based on the theory that migraine headaches arise due to inflammation of the trigeminal nerve branches in the head and neck and that specific trigger sites can be identified in individual patients. Surgical deactivation has also been proposed for other types of headaches (eg, tension headaches).

For individuals who have migraine headaches who receive surgical deactivation of headache trigger sites, the evidence includes randomized controlled trials. The relevant outcomes are symptoms, change in disease status, morbid events, and treatment-related morbidity. Three randomized controlled trials have been published; only one used a sham control and blinded patients to the treatment group. All 3 reported statistically significantly better outcomes at 12 months in patients who received decompression surgery for migraine headache than the control intervention. However, the trials were subject to methodologic limitations (eg, unclear and variable patient selection processes, variability in surgical procedures depending on trigger site). In addition, findings from two trials not blinded or sham-controlled were subject to the placebo effect. Additional sham-controlled randomized studies are needed. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have non–migraine headaches who receive surgical deactivation of headache trigger sites, the evidence includes no published studies. The relevant outcomes are symptoms, change in disease status, morbid events, and treatment-related morbidity. The evidence is insufficient to determine the effects of the technology on health outcomes.

Policy History

Date	Action
4/2019	BCBSA National medical policy review. Description, summary and references updated. Policy statements unchanged.
11/2015	New references added from BCBSA National medical policy.
10/2014	New references added from BCBSA National medical policy.
10/2013	BCBSA National medical policy review. Policy statement clarified.
2/2013	New medical policy describing investigational indications.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

References

1. Bigal ME, Lipton RB. The epidemiology, burden, and comorbidities of migraine. *Neurol Clin.* May 2009;27(2):321-334. PMID 19289218
2. Headache Classification Committee of the International Headache Society. The International Classification of Headache Disorders, 3rd edition (beta version). *Cephalalgia.* Jul 2013;33(9):629-808. PMID 23771276
3. Guyuron B, Reed D, Kriegler JS, et al. A placebo-controlled surgical trial of the treatment of migraine headaches. *Plast Reconstr Surg.* Aug 2009;124(2):461-468. PMID 19644260
4. Liu MT, Armijo BS, Guyuron B. A comparison of outcome of surgical treatment of migraine headaches using a constellation of symptoms versus botulinum toxin type A to identify the trigger sites. *Plast Reconstr Surg.* Feb 2012;129(2):413-419. PMID 21987048
5. Guyuron B, Kriegler JS, Davis J, et al. Comprehensive surgical treatment of migraine headaches. *Plast Reconstr Surg.* Jan 2005;115(1):1-9. PMID 15622223
6. Guyuron B, Kriegler JS, Davis J, et al. Five-year outcome of surgical treatment of migraine headaches. *Plast Reconstr Surg.* Feb 2011;127(2):603-608. PMID 20966820
7. Mathew PG. A critical evaluation of migraine trigger site deactivation surgery. *Headache.* Jan 2014;54(1):142-152. PMID 24116941
8. Omranifard M, Abdali H, Ardakani MR, et al. A comparison of outcome of medical and surgical treatment of migraine headache: In 1 year follow-up. *Adv Biomed Res.* Aug 2016;5:121. PMID 27563631
9. Loder E, Weizenbaum E, Frishberg B, et al. Choosing wisely in headache medicine: the American Headache Society's list of five things physicians and patients should question. *Headache.* Nov-Dec 2013;53(10):1651-1659. PMID 24266337